



Skreden, M., Hillesund, E. R., Wills, A. K., Brantsæter, A. L., Bere, E., & Øverby, N. C. (2018). Adherence to the New Nordic Diet during pregnancy and subsequent maternal weight development: a study conducted in the Norwegian Mother and Child Cohort Study (MoBa). *British Journal of Nutrition*, 119(11), 1286-1294.
<https://doi.org/10.1017/S0007114518000776>

Peer reviewed version

Link to published version (if available):
[10.1017/S0007114518000776](https://doi.org/10.1017/S0007114518000776)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Cambridge University Press at <https://doi.org/10.1017/S0007114518000776> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/pure/user-guides/explore-bristol-research/ebr-terms/>




Figure 2: Predicted marginal mean BMI trajectory for each category of NND adherence score. The plot on the left is from the unadjusted model; the middle plot on the confounder adjusted model (a)*; and the right plot the confounder plus mediator model (b)*. For clarity, only the 95% CI for NND tertile 1 (lowest) is shown (grey).

*Model a: adjusted for maternal age, parity, education, pre-pregnant smoking, physical activity, energy intake and offspring birthweight
 Model b: model a + exclusive breast feeding duration.