



Kjøllestad, M. K. R., Smith, G. D., Ariansen, I., Kinge, J. M., Degerud, E., & Næss, Ø. (2018). The association between BMI and mortality using early adulthood BMI as an instrumental variable for midlife BMI. *Scientific Reports*, 8, Article 11499. <https://doi.org/10.1038/s41598-018-29089-z>

Publisher's PDF, also known as Version of record

License (if available):
CC BY

Link to published version (if available):
[10.1038/s41598-018-29089-z](https://doi.org/10.1038/s41598-018-29089-z)

[Link to publication record on the Bristol Research Portal](#)
PDF-document

This is the final published version of the article (version of record). It first appeared online via Springer Nature at <https://www.nature.com/articles/s41598-018-29089-z>. Please refer to any applicable terms of use of the publisher.

University of Bristol – Bristol Research Portal

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available: <http://www.bristol.ac.uk/red/research-policy/pure/user-guides/brp-terms/>

Original research article

The association between BMI and mortality using early adulthood BMI as an instrumental variable for midlife BMI.

Marte K.R. Kjøllestadal^{1*}, George Davey Smith², Inger Ariansen³, Jonas Minet Kinge^{1,3}, Eirik Degerud³, Øyvind Næss^{1,3}

- 1) Institute of Health and Society, University of Oslo, Norway
- 2) MRC Integrative Epidemiology Unit (IEU), at the University of Bristol, UK
- 3) Domain for Mental and Physical Health, Norwegian Institute of Public Health, Oslo, Norway

Corresponding author: Marte K.R.Kjøllestadal, E-mail: m.k.kjollesdal@medisin.uio.no. Phone: 0047 22 85 06 78. Institute of Health and society, University of Oslo. Postbox 1130 Blindern, 0318 Oslo, Norway.

Supplementary table A. Associations between Body Mass Index (BMI) measured in midlife and in early adulthood (dependent variable) and risk factors for cardiovascular disease in midlife, in three weight categories. From crude linear regressions.

| | BMI, early adulthood (18-20 years) β (95 % CI) | BMI, midlife (40-50 years) β (95 % CI) |
|--|--|--|
| Normal- and underweight | | |
| Cholesterol (mmol/L) | -0.02 (-0.02, -0.02) | 0.08 (0.08, 0.09) |
| Systolic Blood Pressure (mmHg) | -0.27 (-0.33, -0.22) | 1.48 (1.42, 1.54) |
| Diastolic Blood Pressure (mmHg) | -0.17 (-0.21, -0.14) | 0.63 (0.59, 0.67) |
| Heart Rate (beats/min) | -0.30 (-0.35, -0.25) | -0.98 (-1.03, -0.92) |
| Smoking, ref: no | 1.02 (1.01, 1.03) | 0.87 (0.86, 0.87) |
| Current treatment for hypertension, ref: no | 1.10 (0.97, 1.05) | 1.17 (1.11, 1.24) |
| BMI, midlife (kg/m ²) | | 0.35 (0.34, 0.36) |
| Overweight | | |
| Cholesterol (mmol/L) | -0.02 (-0.03, -0.02) | 0.60 (0.06, 0.08) |
| Systolic Blood Pressure (mmHg) | -0.25 (-0.30, -0.20) | 1.09 (1.01, 1.16) |

| | | |
|--|----------------------|---------------------|
| Diastolic Blood Pressure (mmHg) | -0.15 (-0.18, -0.11) | 0.08 (0.80, 0.91) |
| Heart Rate (beats/min) | -0.17 (-0.21, -0.12) | 0.62 (0.54, 0.69) |
| Smoking, ref: no | 1.09 (1.08, 0.10) | 0.96 (0.95, 0.97) |
| Current treatment for hypertension, ref: no | 1.01 (0.99, 1.04) | 1.26 (1.21, 1.31) |
| BMI, midlife (kg/m ²) | | 0.39 (0.38, 0.40) |
| Obese | | |
| Cholesterol (mmol/L) | -0.01 (-0.02, -0.01) | -0.00 (-0.01, 0.00) |
| Systolic Blood Pressure (mmHg) | 0.05 (-0.03, 0.12) | 0.59 (0.50, 0.67) |
| Diastolic Blood Pressure (mmHg) | 0.02 (-0.03, 0.08) | 0.34 (0.27, 0.40) |
| Heart Rate (beats/min) | 0.05 (-0.02, 0.12) | 0.67 (0.59, 0.75) |
| Smoking, ref: no | 1.09 (1.07, 1.10) | 0.98 (0.97, 1.00) |
| Current treatment for hypertension, ref: no | 1.04 (1.02, 1.06) | 1.11 (1.09, 1.13) |
| BMI, midlife (kg/m ²) | | 0.43 (0.41, 0.44) |

Supplementary Table B. Hazard ratios for cardiovascular (CVD) and all-cause mortality with body mass

index (BMI) percentiles from cox regressions.

| | HR (95 % CI) | | | | | |
|------------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| | Model 1 | Model 2 | Model 3 | Model 1 | Model 2 | Model 3 |
| CVD | Men | | | Women | | |
| Midlife | | | | | | |
| 10 percentile | 1.45 (0.91, 1.43) | 1.11 (0.89, 1.39) | 1.11 (0.89, 1.40) | 0.88 (0.60, 1.30) | 0.88 (0.60, 1.30) | 0.91 (0.61, 1.34) |
| 10-50 percentile | 1 | 1 | 1 | 1 | 1 | 1 |
| 50-90 percentile | 1.12 (0.99, 1.26) | 1.07 (0.95, 1.21) | 0.93 (0.82, 1.05) | 1.22 (0.86, 1.74) | 1.14 (0.80, 1.62) | 0.91 (0.64, 1.31) |
| 90 percentile | 2.43 (2.10, 2.81) | 2.19 (1.89, 2.54) | 1.47 (1.25, 1.72) | 2.20 (1.45, 3.33) | 1.91 (0.26, 2.89) | 1.20 (0.76, 1.88) |
| Early adulthood | | | | | | |
| 10 percentile | 1.21 (1.00, 1.46) | 1.17 (0.97, 1.41) | 1.17 (0.96, 1.41) | 1.02 (0.62, 1.70) | 0.96 (0.58, 1.58) | 0.98 (0.59, 1.60) |
| 10-50 percentile | 1 | 1 | 1 | 1 | 1 | 1 |
| 50-90 percentile | 1.17 (1.04, 1.32) | 1.16 (1.03, 1.31) | 1.08 (0.96, 1.22) | 1.02 (0.62, 1.70) | 1.06 (0.72, 1.40) | 0.91 (0.65, 1.26) |
| 90 percentile | 2.22 (1.90, 2.58) | 2.06 (1.77, 2.40) | 1.57 (1.34, 1.84) | 1.56 (1.05, 2.32) | 1.41 (0.95, 2.09) | 1.06 (0.71, 1.59) |

All cause


Midlife

| | | | | | | |
|------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| 10 percentile | 1.53 (1.39, 1.69) | 1.50 (1.36, 1.65) | 1.38 (0.95, 1.25) | 1.24 (1.10, 1.40) | 1.25 (1.11, 1.41) | 1.24 (1.12, 1.38) |
| 10-50 percentile | 1 | 1 | 1 | 1 | 1 | 1 |
| 50-90 percentile | 1.03 (0.97, 1.09) | 0.99 (0.94, 1.05) | 0.95 (0.90, 1.01) | 1.10 (0.97, 1.25) | 1.06 (0.94, 1.20) | 1.01 (0.89, 1.15) |
| 90 percentile | 1.60 (1.47, 1.73) | 1.47 (1.35, 1.59) | 1.25 (1.14, 1.37) | 1.66 (1.42, 1.96) | 1.54 (1.31, 1.81) | 1.39 (1.18, 1.64) |

Early

adulthood

| | | | | | | |
|------------------|-------------------|-------------------|-------------------|--------------------|-------------------|-------------------|
| 10 percentile | 1.17 (1.06, 1.28) | 1.13 (1.11, 1.48) | 1.10 (1.00, 1.21) | 1.07 (0.91, 1.26) | 1.04 (0.88, 1.22) | 1.02 (0.86, 1.20) |
| 10-50 percentile | 1 | 1 | 1 | 1 | 1 | 1 |
| 50-90 percentile | 1.12 (1.05, 1.19) | 1.11 (1.05, 1.18) | 1.05 (1.03, 1.16) | 0.97 (0.87, 1.08) | 0.96 (0.86, 1.07) | 0.93 (0.83, 1.04) |
| 90 percentile | 1.57 (1.45, 1.71) | 1.48 (1.36, 1.56) | 1.32 (1.21, 1.44) | 1.30 (1.213, 1.50) | 1.24 (1.07, 1.42) | 1.12 (0.97, 1.29) |



Supplementary Figure A. Covariate balance (with 95 % confidence intervals) by levels of Body Mass Index (black) and instrument (grey).

