



Whitehouse, M., Berstock, J. R., Kelly, M., Gregson, C., Judge, A., Sayers, A., & Chesser, T. J. (2019). Higher 30-day mortality associated with the use of intramedullary nails compared with sliding hip screws for the treatment of trochanteric hip fractures: a prospective national registry study. *Bone and Joint Journal*, 101-B(1), 83-91. <https://doi.org/10.1302/0301-620X.101B1.BJJ-2018-0601.R2>

Peer reviewed version

Link to published version (if available):
[10.1302/0301-620X.101B1.BJJ-2018-0601.R2](https://doi.org/10.1302/0301-620X.101B1.BJJ-2018-0601.R2)

[Link to publication record in Explore Bristol Research](#)
PDF-document


This is the author accepted manuscript (AAM). The final published version (version of record) is available online via SBJs at <https://online.boneandjoint.org.uk/doi/full/10.1302/0301-620X.101B1.BJJ-2018-0601.R2>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available: <http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Figure 1: Patient inclusions/exclusions into the study



= fracture; SHS = sliding hip screw; L Nail = long intramedullary nail; S Nail = short intramedullary nail; ASA = American Society of Anesthesiologists grade; Pre-op = pre-operative; AMTS = Abbreviated Mental Test Score; MDT = multi-disciplinary team; LSOA = lower layer super output area