



Baritussio, A., Gately, A., Pawade, J., Marks, D. I., & Bucciarelli-Ducci, C. (2017). Extensive cardiac infiltration in acute T-cell lymphoblastic leukemia: occult extra-medullary relapse and remission after salvage chemotherapy. *European Heart Journal*, 38(24), 1933. Article ehw393. <https://doi.org/10.1093/eurheartj/ehw393>

Peer reviewed version

Link to published version (if available):
[10.1093/eurheartj/ehw393](https://doi.org/10.1093/eurheartj/ehw393)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the accepted author manuscript (AAM). The final published version (version of record) is available online via Oxford Academic at <https://doi.org/10.1093/eurheartj/ehw393> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Extensive Cardiac Infiltration in Acute T-Cell Lymphoblastic Leukemia: Occult Extra-Medullary Relapse and Remission after Salvage Chemotherapy.

A. Baritussio MD¹, A. Gately MBChB², J. Pawade FRCPATH³, D. I. Marks MBBS², C. Bucciarelli-Ducci MD¹

¹Bristol Heart Institute, Bristol NIHR Cardiovascular Biomedical Research Unit (BRU), University Hospitals Bristol and University of Bristol, Bristol, United Kingdom

²Bristol Haematology and Oncology Centre, University Hospitals Bristol, Bristol, United Kingdom

³Haemato-Oncology Diagnostic Service, Bristol, United Kingdom

Short Title Extensive cardiac infiltration in acute T-cell lymphoblastic leukemia

Corresponding author:

Dr Chiara Bucciarelli-Ducci

NIHR Bristol Cardiovascular Biomedical Research Unit,

CMR Unit, Bristol Heart Institute

Upper Maudlin Street

Bristol, BS2 8HW, United Kingdom

Email: c.bucciarelli-ducci@bristol.ac.uk

Telephone: 01173425888

A 38 year-old man was admitted with vasculitic rash and horizontal diplopia nine months post allogenic stem cell transplantation for T-cell acute lymphoblastic leukemia (ALL). Magnetic resonance (MR) of the orbits showed enlarged extra-ocular muscles and an echocardiogram, to exclude cardiac embolic sources, showed bi-ventricular hypertrophy and speckled myocardium. A 12-lead ECG showed diffuse T-wave inversion (Panel 1A). A cardiovascular MR (CMR) showed preserved bi-ventricular systolic function, septal left ventricular (LV) hypertrophy (max 21 mm, normal <12mm) (Panel 1B) with increased LV mass (88 g/m², normal 48-77 g/m²). Multiple areas of myocardial and pericardial infiltration (n=9, largest 18x50mm) were noted on advanced tissue characterisation, showing markedly increased T1 values on native T1 mapping, and non-enhancing after contrast administration (Panel 1C and D, white arrows). The CMR findings raised the suspicion of acute cardiac involvement of ALL. A MR-guided biopsy of the swollen ocular left rectus muscle confirmed the diagnosis of relapsed acute T-cell ALL (Panel 2A and B), further supported by evidence of 0.09% T-lymphoblasts on bone marrow flow cytometry and low level central nervous system (CNS) disease on lumbar puncture. As all the investigations confirmed acute multi-organ ALL relapse (heart, extra-ocular muscles, bone marrow and CNS), systemic chemotherapy with *nelarabine* and intra-thecal cytosine *arabioside* were started. Diplopia rapidly improved, and a repeat orbit MR 14 days after commencement of chemotherapy showed complete resolution of extra-ocular muscle enlargement. A 1-month repeat CMR to assess cardiac response demonstrated complete resolution of all nodular lesions with normalization of LV wall thickness (10 mm) and mass (60 g/m²) (Panel 3 B, C and D), and of the widespread T-wave inversion on ECG (Panel 3A).

Declaration of Interest

None of the authors has disclosure to declare.

Role of the Funding Source

No funding source was involved.