



Williamson, J. S., Ambler, G. K., Twine, C. P., Williams, I. M., & Williams, G. L. (2018). Elective Repair of Abdominal Aortic Aneurysm and the Risk of Colonic Ischaemia: Systematic Review and Meta-Analysis. *European Journal of Vascular and Endovascular Surgery*, 56(1), 31-39. <https://doi.org/10.1016/j.ejvs.2018.03.005>

Peer reviewed version

License (if available):  
CC BY-NC-ND

Link to published version (if available):  
[10.1016/j.ejvs.2018.03.005](https://doi.org/10.1016/j.ejvs.2018.03.005)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Elsevier at <https://www.sciencedirect.com/science/article/pii/S1078588418301400> . Please refer to any applicable terms of use of the publisher.

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available: <http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Table 2. Outcome data for each study. Abbreviations. IMA (inferior mesenteric artery), IIA (internal iliac artery), NS (Not stated in the manuscript), CI (colonic ischaemia), CA(Coeliac artery), SMA (Superior mesenteric artery).

Author (year)	Intervention	CI rate % (EVAR)	CI rate % (OR)	CI mortality rate	Reintervention rate EVAR	Reintervention rate OR	Time to CI	Preoperative visceral arterial status	IMA intervention	IIA intervention
Schermerhorn 2008	Both	1	2.1	NS	'bowel resection' 3% p=0.02	Bowel resection 3.4% p=0.02	NS	NS		
Miller 2009	EVAR	1.4		4 of 11	3 of 11		7/11 within 30 days, 4/11 after 30 days	IMA and IIA patency	10/11 with CI had pre-existing IMA occlusion	Unilateral IIA embolization increased risk of CI. No comparable IMA data.
Valentine 1998	OR		0.83	0 of 1		1 of 1	11 days	CA and SMA not IMA, IIA	No difference in 'GI' complications	
Dadian 2001	EVAR	2.9		3 of 8	2 of 8		7 of 8 within 30 days, 1 within 6 months	IMA and IIA patency/embolization	5/8 with CI had preop occluded IMA. No comparable data without CI..	No effect on CI rates with uni or bilateral IIA embolization
Geraghty 2004	EVAR	1.7		2 of 4	3 of 4		2 +/- 1.4 days	IMA not reported. IIA patency/embolization reported		No effect on CI rates with uni or bilateral IIA embolisation
Maldonado 2004	EVAR	1.2		2 of 4	3 of 4		3<12h, 1<7 days	IMA and IIA patency/embolization	All with CI had patent IMA preop.	No effect on CI with preop uni or bilateral IIA embolization.
Perry 2008	Both	0.5	2.2	37.8%	27% colectomy p<0.01)	31% colectomy p<0.01	NS	NS		
Chiesa 2012	OR		3	NS				NS		
Cruz 2001	OR		1.6	0 of 2				NS		
Bonardelli 2012	Both	0	0.3	0 of 1	0 of 0	0 of 1	After 30 days	NS		
Mehta 2005	Both	0.6	2.6	NS	1 of 1	6 of 6	NS	Data could not be extracted		No patients developed CI where bilateral IIA sacrifice performed
Utlee 2016	Both	0.6	3.6	25/107 (23%)	14/26 (54%)	37/78 (47%)	NS	IMA and IIA patency/embolization/reimplantation	Higher risk of CI if IMA reimplantation performed in OR	.Higher risk of CI if unilateral IIA ligation/embolization in OR and EVAR
Hynes 2017	Both	0.3%	0.7%		11/13	10/11	9/24< 24h, 18/19<7d	NS		