



Hamilton, H. B., & Wass, D. F. (2017). How Important Are Radical Mechanisms in Frustrated Lewis Pair Chemistry? *Chem*, 3(2), 198-199. <https://doi.org/10.1016/j.chempr.2017.07.003>

Publisher's PDF, also known as Version of record

License (if available):
Other

Link to published version (if available):
[10.1016/j.chempr.2017.07.003](https://doi.org/10.1016/j.chempr.2017.07.003)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the final published version of the article (version of record). It first appeared online via Elsevier at <https://www.sciencedirect.com/science/article/pii/S2451929417303108?via%3Dihub>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available: <http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Preview

How Important Are Radical Mechanisms in Frustrated Lewis Pair Chemistry?

Hugh B. Hamilton¹ and Duncan F. Wass^{1,*}

Reaction mechanism is the foundation of understanding chemical reactivity. In this issue of *Chem*, Stephan and co-workers overturn long-held assumptions regarding the reactivity of frustrated Lewis pairs by providing compelling evidence that radical mechanisms might operate in some cases.

Frustrated Lewis pair (FLP) chemistry has been perhaps the most significant development in molecular main-group chemistry over the past decade. The realization that certain sterically encumbered combinations of Lewis acids and Lewis bases can act cooperatively to activate small molecules such as hydrogen has led to a wealth of new chemistry, not least in the use of such systems in catalysis.¹ One of the features that has caught the imagination of the community is that this area builds on one of the very first concepts we are taught in studying chemistry, namely Lewis's description of acids and bases as electron-pair acceptors or donors. It is inherently exciting to have something original to say about a nearly 100-year-old concept, particularly when it leads to such rich, varied, and useful chemistry.² The received wisdom in FLP chemistry is that, on the basis of Lewis's original insight for acids and bases, it involves the transfer of electron pairs. Previous work in this area has always corroborated, or assumed, that the activation of small molecules with FLPs is a two-electron process.

The work published in this issue of *Chem* by Stephan and co-workers³ suggests that we need to revisit this assumption; a purely heterolytic, two-electron model of FLP-mediated small-molecule activa-

tion might not always be applicable. This work provides compelling evidence that, in some circumstances, a single-electron transfer (SET) mechanism is operating. This challenge to the previous assumption is all the more remarkable in that the systems studied are some of the most well-known FLPs based on boron or aluminum Lewis acids and triarylphosphine Lewis bases. The subtleties of FLP chemistry clearly require more sophisticated rationales than has hitherto been the case, and this will enable a deeper understanding of the mechanistic processes involved.

Starting with the well-established FLP combinations of ^tBu₃P and E(C₆F₅)₃ (E = B, Al), Stephan and co-workers first demonstrate the inherent "frustration" of these FLPs before presenting their reactivity with tetrachloro-1,4-benzoquinone (*p*-O₂C₆Cl₄). Equimolar reactions afford the expected products ^tBu₃POC₆Cl₄OE(C₆F₅)₃.

E(C₆F₅)₃ shows only weak interactions when combined with Ph₃SnH, but each ^tBu₃P/E(C₆F₅)₃ FLP results in the formation of the (stannyl)phosphonium salt [^tBu₃PSnPh₃][HB(C₆F₅)₃] or [^tBu₃PSnPh₃][(μ -H)(Al(C₆F₅)₃)₂]. The structures of these were elucidated through nuclear magnetic resonance (NMR) spectroscopic and crystallographic analysis. These heterolytic-cleavage and FLP-

addition reactions portrayed "classic" FLP behavior—the acceptance and donation of an electron pair between the substrate and the two components of the FLP. However, when the phosphine was changed from ^tBu₃P to Mes₃P, still a well-known FLP combination, the results were surprising. Instant color changes were seen when Mes₃P was mixed with the two Lewis acids, even though there was little NMR evidence to suggest any change from the starting reactants. Experiments with electron paramagnetic resonance (EPR) spectroscopy were the key to unlocking the nature of this system—the results clearly pertain to the existence of the [Mes₃P·]⁺ radical cation for the Mes₃P/Al(C₆F₅)₃ FLP. This insinuates the presence of the [·Al(C₆F₅)₃]⁻ counterion, the short lifetime of which explains its absence in the EPR spectrum. Weaker signals were seen in the EPR spectrum for the Mes₃P/B(C₆F₅)₃ FLP, making any conclusions less definitive; however, the presence of a single-electron transfer (SET) equilibrium between Mes₃P/Al(C₆F₅)₃ and the respective charged radical species is certainly an intriguing suggestion.

Reactions with 0.5 equiv of *p*-O₂C₆Cl₄ provide further evidence for the presence of radical species, such that UV-visible spectroscopy supports the existence of the [Mes₃P·]⁺ cation after the Mes₃P/E(C₆F₅)₃ (E = B, Al) FLP was combined with the quinone. It should be noted that this was not the case when the phosphine and quinone were analyzed together in the absence of E(C₆F₅)₃. [Mes₃P·]₂[(C₆F₅)₃EOC₆Cl₄OE(C₆F₅)₃] is the expected product of these FLP reactions with quinone. Crystallographic data were obtainable only upon the addition of Ph₃SnH, which afforded [Mes₃PH]₂[(C₆F₅)₃BOC₆Cl₄OB(C₆F₅)₃].

¹School of Chemistry, University of Bristol, Cantock's Close, Bristol BS8 1TS, UK

*Correspondence: duncan.wass@bristol.ac.uk
<http://dx.doi.org/10.1016/j.chempr.2017.07.003>




Figure 1. Electron-Pair Transfer or SET Mechanisms in FLP Chemistry

The addition of 1 equiv of *p*-O₂C₆Cl₄ to the FLPs produced a purple solution, which changed to pale yellow after a short amount of time. This suggests a two-step process whereby [Mes₃P•]₂ [(C₆F₅)₃EOC₆Cl₄OE(C₆F₅)₃] (E = B, Al) is initially formed and then converted to [(Mes₃POC₆Cl₄OE(C₆F₅)₃]. NMR spectroscopic and crystallographic data support this chain of events. This argument is further supported by the absence of any reaction between [Mes₃P•][(μ-HO)(Al(C₆F₅)₃)₂] and *p*-O₂C₆Cl₄, showing that [•OC₆Cl₄OE(C₆F₅)₃][−] is not formed in this way.⁴

Additional evidence for this mechanism was obtained through the addition of Ph₃SnH to the Mes₃P/E(C₆F₅)₃

(E = B, Al) FLPs; NMR spectra indicated the formation of [Mes₃PH][HB(C₆F₅)₃] and [Mes₃PH][(μ-H)(Al(C₆F₅)₃)₂] and the release of Ph₃SnSnPh₃ as a side product. This is indicative of H• abstraction from Ph₃SnH, demonstrating homolytic cleavage of the Sn–H bond, which was cleaved heterolytically by the ^tBu₃P/E(C₆F₅)₃ (E = B, Al) FLPs.

The crucial point here is that although both Mes₃P/E(C₆F₅)₃ and ^tBu₃P/E(C₆F₅)₃ give analogous products upon activation of H₂,^{1,5} the route by which this is achieved is likely to be different: either a standard, two-electron transfer process in the case of ^tBu₃P/E(C₆F₅)₃ or SET in the case of Mes₃P/E(C₆F₅)₃ (Figure 1). This

has important implications for the scope and selectivity of these systems.

These results should inspire many future studies, and two aspects immediately come to mind. First, it is intriguing that the FLPs described here are well known but that some of the key observations (for example, the color change in mixing Mes₃P and E(C₆F₅)₃) have previously been either ignored or not deemed worthy of further investigation. How many previous FLP systems actually operated via a SET mechanism? A systematic study would be valuable and would greatly aid a more sophisticated description of FLP processes. Analysis of these systems via computational methods should also aid understanding. Second, using this more detailed understanding of the factors that lead to electron-pair or single-electron pathways should enable the design of FLPs that specifically operate via one or the other mechanism. This should allow further expansion of an already compelling area of chemistry.

- Stephan, D.W., and Erker, G. (2015). *Angew. Chem. Int. Ed.* 54, 6400–6441.
- Welch, G.C., San Juan, R.R., Masuda, J.D., and Stephan, D.W. (2006). *Science* 314, 1124–1126.
- Liu, L., Cao, L.L., Shao, Y., Ménard, G., and Stephan, D.W. (2017). *Chem* 3, this issue, 259–267.
- Ménard, G., Hatnean, J.A., Cowley, H.J., Lough, A.J., Rawson, J.M., and Stephan, D.W. (2013). *J. Am. Chem. Soc.* 135, 6446–6449.
- Ménard, G., Tran, L., and Stephan, D.W. (2013). *Dalton Trans.* 42, 13685–13691.