



Walker, S., & Maskell, N. (2017). Pneumothorax management—chest drain or needle aspiration? *Journal of Thoracic Disease*, 23(4), 339-345. <https://doi.org/10.21037/jtd.2017.08.155>

Peer reviewed version

License (if available):
Other

Link to published version (if available):
[10.21037/jtd.2017.08.155](https://doi.org/10.21037/jtd.2017.08.155)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the accepted author manuscript (AAM). The final published version (version of record) is available online via AME at <https://doi.org/10.21037/jtd.2017.08.155> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available: <http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Despite pneumothoraces being described in medical literature as far back as 15th century(1), the optimal way to remove air from the pleural cavity has not yet been ascertained, nor indeed have we determined whether it is necessary to do so at all. The well conducted randomised control trial by Thelle et al(2) has added to the evidence base concerning one of the fundamental questions: whether needle aspiration (NA) or chest tube drainages (CTD) is superior in evacuating a pneumothorax.

It is an important question, particularly as expert consensus remains divided, with differing advice from national advisory bodies. The American College of Chest Physicians Delphi consensus (2001) (3) does not advise NA, instead supporting proceeding directly with CTD insertion, whilst the BTS guidelines (2010)(4) suggests NA as 1st line in primary spontaneous pneumothorax (PSP), and as an option in small sub-centimetre secondary spontaneous pneumothorax (SSP). There have been several studies(5-10) over the last 25 years which have attempted to address the issue, although interpretation has been made difficult with heterogeneous inclusion criteria, methodologies and definitions of success. However, taken together, they suggest that whilst hospital duration is typically shorter for patients treated with NA(5-7, 9, 10), whilst immediate success rates is generally higher with CTD insertion (5, 7, 8).

Thelle et al study randomised 127 patients presenting with a spontaneous pneumothorax, both primary and secondary, to either NA or CTD. The primary outcome was length of stay, with secondary outcomes including immediate and one-week success rates. Patients were allowed two aspirations in the NA cohort, before proceeding to CTD insertion if these failed. The study demonstrated hospital length of stay which were significantly shorter in all patients treated with NA pathway, including each sub-group (PSP and SSP). Immediate success with NA was almost twice that of CTD. Whilst with rates in NA arm not dissimilar to preceding studies, for CTD it was much lower (32% vs 64-93%) (5, 7, 8). It is also important to note that the high immediate success rate (73.8%) with needle aspiration in SP represents a management pathway consisting of a subsequent second aspiration and then chest drain if required. The success of the initial NA was a less encouraging 50%. The low incidence of adverse event in NA in this study also supported previous literature.

The success of NA in both PSP and SSP patients provides interesting insight into pneumothorax research. CTD has been favoured in patients where persistent air leak is suspected, as it is felt that NA would be more likely to fail in these circumstances. This reasoning is also why CTD is preferred in SSP, with the reasonable supposition that these patients will be more likely to have ongoing air-leak due to bullae rupture. With NA associated with twice the immediate success rates of CTD in both PSP and SSP, this suggests either that persistent air leak is uncommon in both these cohorts, or less likely a chest tube is not required to manage the persistent air leak. Additionally, there has been supposition, with increasing evidence that the underlying lung in PSP is abnormal, that PSP and SSP are not two distinct entities, but differing ends of the same spectrum(11). If this study's findings are correct, and the initial treatment pathway should be the same, then there is less reason to differentiate between them.

There may have been an opportunity with the study in helping us predict which patients will develop pneumothorax recurrence. Results from 6 and 12 months would have helped clarify some uncertainties about the long-term recurrence in SP, and more whether particularly the method of treatment influences recurrence rates.

If the findings of this study are verified by further studies, particularly on SSP patients, needle aspiration, with its associated fewer bed-days and adverse events is an attractive first line option and should be offered to patients with a PSP or SSP. However, patients must be counselled that there is a 50:50 chance of their initial NA failing, and requiring further intervention. The study authors should be congratulated on a well-designed study which, by including both PSP and SSP, progresses the understanding of pneumothoraces and its management further.

1. Kaya SO, Karatepe M, Tok T, Onem G, Dursunoglu N, Goksin I. Were pneumothorax and its management known in 15th-century anatolia? *Texas Heart Institute Journal*. 2009;36(2):152.
2. Thelle A, Gjerdevik M, SueChu M, Hagen O, Bakke P. Randomised comparison of needle aspiration and chest tube drainage in spontaneous pneumothorax. *European Respiratory Journal*. 2017;49(4):1601296.
3. Baumann MH, Strange C, Heffner JE, Light R, Kirby TJ, Klein J, et al. Management of spontaneous pneumothorax: an American College of Chest Physicians Delphi consensus statement. *CHEST Journal*. 2001;119(2):590-602.
4. MacDuff A, Arnold A, Harvey J, Group BTSPDG. Management of spontaneous pneumothorax: British Thoracic Society Pleural Disease Guideline 2010. *Thorax*. 2010;65 Suppl 2(Suppl 2):ii18-31.
5. Ayed AK, Chandrasekaran C, Sukumar M. Aspiration versus tube drainage in primary spontaneous pneumothorax: a randomised study. *Eur Respir J*. 2006;27(3):477-82.
6. Harvey J, Prescott RJ. Simple aspiration versus intercostal tube drainage for spontaneous pneumothorax in patients with normal lungs. *Bmj*. 1994;309(6965):1338-9.
7. Noppen M, Alexander P, Driesen P, Slabbynck H, Verstraeten A. Manual aspiration versus chest tube drainage in first episodes of primary spontaneous pneumothorax: a multicenter, prospective, randomized pilot study. *American journal of respiratory and critical care medicine*. 2002;165(9):1240-4.
8. Andrivet P, Djedaini K, Teboul J-L, Brochard L, Dreyfuss D. Spontaneous pneumothorax: comparison of thoracic drainage vs immediate or delayed needle aspiration. *Chest*. 1995;108(2):335-9.
9. Korczyński P, Górska K, Nasiłowski J, Chazan R, Krenke R. Comparison of small bore catheter aspiration and chest tube drainage in the management of spontaneous pneumothorax. *Noncommunicable Diseases: Springer*; 2015. p. 15-23.
10. Parlak M, Uil SM, van den Berg JW. A prospective, randomised trial of pneumothorax therapy: manual aspiration versus conventional chest tube drainage. *Respiratory medicine*. 2012;106(11):1600-5.
11. Bintcliffe OJ, Hallifax RJ, Edey A, Feller-Kopman D, Lee YC, Marquette CH, et al. Spontaneous pneumothorax: time to rethink management? *The Lancet Respiratory medicine*. 2015;3(7):578-88.