



Hudson, B. E., Ameneshoa, K., Gopfert, A., Goddard, R., Forbes, K., Verne, J., Collins, P., Gordon, F., Portal, A. J., Reid, C., & McCune, C. A. (2017). Integration of palliative and supportive care in the management of advanced liver disease: development and evaluation of a prognostic screening tool and supportive care intervention. *Frontline Gastroenterology*, 8(1), 45-52.
<https://doi.org/10.1136/flgastro-2016-100734>

Other version

License (if available):
CC BY-NC

Link to published version (if available):
[10.1136/flgastro-2016-100734](https://doi.org/10.1136/flgastro-2016-100734)

[Link to publication record on the Bristol Research Portal](#)
PDF-document

This is the submitted manuscript. The final published version (version of record) is available online via BMJ at <http://dx.doi.org/10.1136/flgastro-2016-100734> . Please refer to any applicable terms of use of the publisher.

University of Bristol – Bristol Research Portal

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/brp-terms/>

FIGURE 1 – Rapid Cycle Plan-Do-Study-Act methodology [16]




FIGURE 2 - Integration of prognostic screening tool and development of supportive care intervention using rapid-cycle PDSA methodology




FIGURE 3 - Integration of prognostic screening into weekly hepatology MDT proforma, completed weekly for each hepatology inpatient at University Hospitals Bristol Trust (front and reverse of sheet)

LIVER MDT Date: _____

Patient name: _____
 Patient number: _____
 Date of admission: _____
 Consultant in charge: _____

Primary Diagnosis:

Cirrhosis (UKELD score = __)
 Varices
 Encephalopathy
 Ascites
 Addiction issues

Current Issues:

Discussion and plan:

Ceiling of care:

Full
 Ward-based
 Symptomatic

Calculating the Child Pugh Score for Cirrhosis Mortality	Parameter	Points assigned		
		1	2	3
	Ascites	Absent	Mild	Moderate-Severe
	Encephalopathy	None	Grade 1-2	Grade 3-4
	Bilirubin (micromol/L)	<34.2	34.2 – 51.3	>51.3
	Albumin (g/L)	>35	28-35	<28
	INR	<1.7	1.7-2.3	>2.3

Child Pugh A	5 – 6 points	100% 1 year survival
Child Pugh B	6 – 9 points	81% 1 year survival
Child Pugh C	> 10 points	45% 1 year survival

West Haven Grading of Encephalopathy	
Grade	Criteria
1	Trivial lack of awareness
	Euphoria or anxiety
	Shortened attention span
	Impaired performance of addition
2	Lethargy or apathy
	Minimal disorientation of time or place
	Subtle personality changes
	Inappropriate behavior
	Impaired performance of subtraction
3	Somnolence to semi-stupor but responsive to verbal stimuli
	Confusion
	Gross disorientation
4	Coma

WHO performance status	
0	Fully active, able to carry out all pre-disease performance without restriction
1	Restricted in physically strenuous activity but ambulatory and able to carry out work of a light or sedentary nature
2	Ambulatory and capable of all self-care but unable to carry out any work activities. Up and about >50% of the time
3	Capable of only limited self-care. Confined to bed or chair >50% of the time
4	Completely disabled. Cannot carry out self-care. Totally confined to bed or chair

Suitability for liver transplant assessment

A patient's current suitability for liver transplant assessment and work up is multifactorial and complex. Decisions regarding this are made at consultant level with support from the MDT.

There are however some clear factors which, when present, render patients unsuitable at the current time, and for whom "unsuitable for transplant work up" can be ticked on the poor prognosis scoring criteria:

- Ongoing alcohol use in the context of previously diagnosed alcohol related liver disease
- Ongoing disruptive substance abuse
- Untreated malignancy (not including HCC)
- Life expectancy < 1 year due to non-hepatic co-morbidity
- Age >75 (unless exceptional circumstances)

Poor prognosis screening: cirrhotic patients only	
Criteria	Tick
Child Pugh Grade C	
> 2 liver-related admissions last 6 months	
Ongoing alcohol use (ARLD patients)	
Unsuitable for transplant work-up	
WHO performance status 3-4	

Total score: _____

If total score > 2, consider:

Poor prognosis discussion with patient/family
 Poor prognosis letter to GP
 Advance care planning discussions
 Specialist palliative care referral
 Allocation of hepatology specialist nurse

COMPLETED BY: _____
 SIGNED: _____
 GMC NUMBER: _____