



Supic, I., Skrzypczyk, P., & Cavalcanti, D. (2019). Methods to estimate entanglement in teleportation experiments. *Physical Review A*, 99(3), [032334]. <https://doi.org/10.1103/PhysRevA.99.032334>

Publisher's PDF, also known as Version of record

License (if available):
Other

Link to published version (if available):
[10.1103/PhysRevA.99.032334](https://doi.org/10.1103/PhysRevA.99.032334)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the final published version of the article (version of record). It first appeared online via APS at <https://doi.org/10.1103/PhysRevA.99.032334> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available: <http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Methods to estimate entanglement in teleportation experiments

Ivan Šupić,¹ Paul Skrzypczyk,² and Daniel Cavalcanti³

¹*Département de Physique Appliquée, Université de Genève, 1211 Genève, Switzerland*

²*H. H. Wills Physics Laboratory, University of Bristol, Tyndall Avenue, Bristol BS8 1TL, United Kingdom*

³*ICFO-Institut de Ciències Fòniques, The Barcelona Institute of Science and Technology, 08860 Castelldefels (Barcelona), Spain*



(Received 29 October 2018; published 22 March 2019)

Quantum state teleportation is a protocol where a shared entangled state is used as a quantum channel to transmit quantum information between distinct locations. Here we consider the task of estimating entanglement of the underlying shared state in teleportation protocols. We show that the data accessible in a teleportation experiment allows us to put a lower bound on some entanglement measures, such as entanglement negativity and robustness. Furthermore, we show cases in which the lower bounds are tight. The introduced lower bounds can also be interpreted as quantifiers of the nonclassicality of a teleportation experiment. Thus, our findings provide a quantitative relation between teleportation and entanglement.

DOI: [10.1103/PhysRevA.99.032334](https://doi.org/10.1103/PhysRevA.99.032334)

I. INTRODUCTION

The seminal work by Bennett *et al.* [1] from 1993 demonstrated the possibility to faithfully transfer the quantum state of a system to a spatially distant one, without having to physically send it. Named quantum teleportation, this protocol made a huge impact on the development of quantum information processing, being a building block for more advanced protocols such as cryptographic tasks [2], quantum repeaters [3], quantum computing [4,5], and many others.

Ideally, in order to realize teleportation, two parties, Alice and Bob, need to share a pair of particles in a maximally entangled state $|\Phi^+\rangle = \sum_{i=0}^{d-1} |ii\rangle/\sqrt{d}$, where d is the local Hilbert space dimension of the system. Then, Alice applies a joint Bell state measurement (a measurement where all measurement operators are maximally entangled) on a third system in state $|\omega\rangle$ and her share of the maximally entangled state and communicates the result to Bob. Bob, upon receiving the message from Alice, applies a unitary operation on his system, which ends up in the desired state $|\omega\rangle$. A very important feature of quantum teleportation is that Alice does not need to know the state $|\omega\rangle$.

In realistic conditions it is impossible to achieve perfect quantum teleportation. There has been a lot of effort in describing imperfect teleportation as well as the role of generic entangled states in the protocol. In a teleportation experiment where a set of N_x states $\{\omega_x\}_{x=1}^{N_x}$ —which need not necessarily be pure states—is teleported, the most common benchmark between classical and quantum teleportation is the average fidelity of teleportation [6]

$$\bar{F}_{\sigma_{a|\omega_x}} = \frac{1}{N_x} \sum_{a,x} p(a|\omega_x) F(U_a \rho_{a|\omega_x}^B U_a^\dagger, \omega_x), \quad (1)$$

where $F(\rho, \sigma) = \|\sqrt{\rho}\sqrt{\sigma}\|_1$ is the fidelity, and

$$\rho_{a|\omega_x}^B = \frac{\text{tr}_{\text{VA}}[(M_a^{\text{VA}} \otimes \mathbb{1}^{\text{B}})(\omega_x^{\text{V}} \otimes \rho^{\text{AB}})]}{p(a|\omega_x)} \quad (2)$$

are the states Bob obtains, conditioned on the input state ω_x and Alice's measurement output a , while $p(a|\omega_x)$ is the probability for Alice to obtain the outcome a when the input state is ω_x . The teleportation process is considered to be quantum (i.e., nonclassical) if the fidelity of teleportation is higher than the fidelity that could be obtained using solely classical resources (i.e., no entanglement preshared between Alice and Bob). Based on this figure of merit, not all entangled states are useful for achieving nonclassical teleportation [6,7], among them the bound entangled states [8].

Notice that the average fidelity of teleportation is a coarse-grained measure, reducing all the information available in a teleportation experiment to a single number. Thus, it could happen that, even though an entangled state could not achieve an average fidelity higher than a separable state, a deeper analysis of the relation between input and output states, summarized by the states (2) and observed statistics $p(a|\omega_x)$, could lead to a better assessment on the nonclassical nature of the teleportation process. Motivated by this, in our recent work [9] we proved that the information available in a teleportation experiment allows us to prove that every entangled state can be used to demonstrate a nonclassical teleportation process, if a suitable set of input states and measurement are chosen. The method was moreover used to experimentally demonstrate nonclassical teleportation stemming from a state unable to outperform separable states in terms of the average fidelity of teleportation [10].

Our main goal is to go beyond detection of nonclassicality, and to show that the teleportation data can also be used to estimate the amount of entanglement of the state shared between Alice and Bob. This can be done in two ways: by considering the full teleportation data (the correlations between input and output states) or through the violation of teleportation witnesses (linear functions of teleportation data).

Let us recall the formalism for describing quantum state teleportation introduced in [9]. Quantum teleportation manifests in the nonlocal correlations between Alice's joint measurement outputs and states prepared for Bob. A teleportation

experiment is nonclassical if it excludes a “local-hidden-channel model,” which would, in a classical way, correlate Alice’s outputs with Bob’s reduced states. In order to see exactly the form of such classical teleportation channels let us see what data Alice and Bob can observe if they share a separable state. The set of reduced states of Bob forms a teleportation assemblage (teleportage in [11]). In a teleportation experiment, Bob’s (unnormalized) state, given Alice’s input state is ω_x^V , is given by

$$\begin{aligned} \sigma_{a|\omega_x}^B &= \text{tr}_{VA} [(M_a^{VA} \otimes \mathbb{1}^B)(\omega_x^V \otimes \rho^{AB})] \\ &= \text{tr}_V [\tilde{M}_a^{VB}(\omega_x^V \otimes \mathbb{1}^B)], \end{aligned} \quad (3)$$

where M_a^{VA} are the operators describing the measurement happening inside Alice’s box, and

$$\tilde{M}_a^{VB} = \text{tr}_A [(M_a^{VA} \otimes \mathbb{1}^B)(\mathbb{1}^V \otimes \rho^{AB})]. \quad (4)$$

Notice that the normalization of $\sigma_{a|\omega_x}^B$ gives the probabilities of Alice’s outcomes $p(a|\omega_x)$.

If ρ^{AB} is separable, i.e., $\rho^{AB} = \sum_\lambda p_\lambda \rho_\lambda^A \otimes \rho_\lambda^B$, then [9]

$$\tilde{M}_a^{VB} = \sum_\lambda p_\lambda M_{a|\lambda}^V \otimes \rho_\lambda^B, \quad (5)$$

where

$$M_{a|\lambda}^V = \text{tr}_A [M_a^{VA}(\mathbb{1}^V \otimes \rho_\lambda^A)]. \quad (6)$$

We call this case a local-hidden-channel model for the teleportation experiment, since it can be understood in the following way: at each round of the experiment a classical variable λ is sent to Alice and Bob. Upon reading the value of λ Alice’s device output a with probability $p(a|\omega_x, \lambda) = \text{tr}[M_{a|\lambda}^V \omega_x^V]$ and Bob’s device generates the state ρ_λ .

Given an observed teleportation data, i.e., $\{\sigma_{a|\omega_x}^B\}$, such classical model can be tested by solving the following optimization problem:

$$\begin{aligned} \text{given} \quad & \{\sigma_{a|\omega_x}^B\}_{a,x}, \{\omega_x^V\}_x \\ \text{find} \quad & \{\tilde{M}_a^{VB}\}_a \\ \text{s.t.} \quad & \sigma_{a|\omega_x}^B = \text{tr} [\tilde{M}_a^{VB}(\omega_x^V \otimes \mathbb{1}^B)] \quad \forall a, x, \\ & \tilde{M}_a^{VB} \in \mathcal{S} \quad \forall a, \end{aligned} \quad (7)$$

where \mathcal{S} denotes the set of separable operators, i.e., operators of the form $\sum_\lambda \tau_\lambda \otimes \chi_\lambda$, with $\tau_\lambda \geq 0$ and $\chi_\lambda \geq 0$ for all λ . This set cannot be characterized in a simple way, but its relaxations to the set of operators with positive partial transpose (PPT) or the set of k -shareable operators [12] can be characterized through linear SDP constraints [13].

In Ref. [9] we proved that with a suitable set of input states and a suitable measurement for Alice, every entangled state leads to a nonclassical teleportation data, in the sense of not having the models (3) and (5).

To conclude the Introduction, let us note that there are two elements contributing to the constraint that the channel operators (4) are separable: the shared state ρ^{AB} is separable; or Alice’s measurement operators M_a^{VA} are not entangling operators. Indeed, taking $M_a^{VA} = \sum_\lambda \tau_{\lambda,a}^V \otimes \chi_{\lambda,a}^A$ makes the

channel operators separable:

$$\begin{aligned} \tilde{M}_a^{VB} &= \text{tr}_A \left[\left(\sum_\lambda \tau_{\lambda,a}^V \otimes \chi_{\lambda,a}^A \otimes \mathbb{1}^B \right) (\mathbb{1}^V \otimes \rho^{AB}) \right] \\ &= \sum_\lambda \tau_{\lambda,a}^V \otimes \pi_{\lambda,a}^B, \end{aligned} \quad (8)$$

where $\pi_{\lambda,a}^B = \text{tr}[(\chi_{\lambda,a}^A \otimes \mathbb{1}^B)\rho^{AB}]$. With this in mind, infeasibility of the problem (7) certifies two things simultaneously:

- (1) the state ρ^{AB} is entangled,
- (2) the measurements M_a^{VA} are entangling.

In the rest of this paper we show that entanglement not only leads to qualitatively different teleportation experiments than separable states, but it can also be quantified from a teleportation experiment. In Sec. II we discuss possibilities to quantify entanglement negativity from a teleportation experiment. In Sec. III we put a lower bound on robustness of entanglement and define various robustness-based teleportation quantifiers. Finally, in Sec. IV we introduce teleportation weight and relate it to the best separable approximation of an entangled state.

II. ESTIMATING ENTANGLEMENT NEGATIVITY FROM A TELEPORTATION EXPERIMENT

Negativity of entanglement [14] is a widely used entanglement measure, largely due to the fact that it can be computed efficiently. In the original paper introducing entanglement negativity [14], the authors examined relation between “teleportation capacity” of a quantum state and its negativity. More precisely, the authors proved that the average fidelity of teleportation puts a lower bound on the entanglement negativity of the shared state. Since we introduced a way to characterize teleportation experiment beyond average fidelity, we expect that our method can give some further insight into the role of entanglement negativity in teleportation. In this section we prove that, indeed, by using all the accessible information in a teleportation experiment one can place lower bound on the entanglement negativity of the shared state ρ^{AB} .

A. Estimating entanglement negativity from teleportation data

The entanglement negativity of a state ρ^{AB} can be expressed in the following way:

$$\begin{aligned} \mathcal{N}(\rho_{AB}) &= \min_{\rho_+, \rho_-} \text{tr}(\rho_-) \\ \text{s.t.} \quad & \rho^{AB} = \rho_+ - \rho_- \\ & \rho_\pm^{TA} \geq 0. \end{aligned} \quad (9)$$

Let us see how this optimization problem can be rephrased in terms of the available teleportation data, i.e., the teleportation assemblage $\{\sigma_{a|\omega_x}\}_{a,x}$. Analogously to (3) we can introduce the auxiliary teleportation assemblages

$$\sigma_{a|\omega_x}^\pm = \text{tr}_{VA} [(M_a^{VA} \otimes \mathbb{1}^B)(\omega_x^V \otimes \rho_\pm^{AB})].$$

With this notation the first constraint from (9) can be written as

$$\sigma_{a|\omega_x} = \sigma_{a|\omega_x}^+ - \sigma_{a|\omega_x}^-.$$

The objective function is easily identified as

$$\sum_a \text{tr}[\sigma_{a|\omega_x}^-] = \text{tr}[\omega_x^V \otimes \rho_{-}^{AB}] = \text{tr}[\rho_{-}^{AB}]. \quad (10)$$

Finally we have to characterize the effective measurements $M_{a,\pm}^{*VB}$ which can arise from a PPT state. From (4) and the identity

$$\text{tr}_B[(\mathbb{1}^A \otimes M^{BC})(\Phi^{+AB} \otimes \mathbb{1}^C)] = \frac{1}{d}(M^{AC})^{T_A} \quad (11)$$

we obtain

$$d(M_a^{*VB})^T = \text{tr}_{V_1A}[(\mathbb{1}^V \otimes M_a^{V_1A} \otimes \mathbb{1}^B)(\Phi^{+VV_1} \otimes \rho^{ABT_B})]. \quad (12)$$

If the state ρ^{AB} is PPT the left-hand side of the last equation represents an unnormalized quantum state, which means that the operators $(M_a^{*VB})^T$ and hence (M_a^{*VB}) are also positive, which justifies the last constraint from (13). Now we are ready to construct a semidefinite program whose solution represents a lower bound on the negativity of the shared state ρ^{AB} :

$$\begin{aligned} \min_{\{M_{a,\pm}^*\}_a, \rho_{\pm}} \quad & \sum_a \text{tr}[\sigma_{a|\omega_x}^-] \\ \text{s.t.} \quad & \sigma_{a|\omega_x} = \sigma_{a|\omega_x}^+ - \sigma_{a|\omega_x}^-, \\ & \sigma_{a|\omega_x}^{\pm} = \text{tr}_V[M_{a,\pm}^{*VB}(\omega_x^V \otimes \mathbb{1}^B)], \\ & \sum_a M_{a,\pm}^{*VB} = \mathbb{1}^V \otimes \rho_{\pm}^B, \\ & M_{a,\pm}^{*VB} \geq 0. \end{aligned} \quad (13)$$

The next step is to prove that the solution of (13) lower bounds the negativity of entanglement of the state ρ^{AB} , given by the solution of (9). First let us note that in the case the set of input states is tomographically complete and Alice applies the Bell state measurement, the solutions to (9) and (13) coincide. To see that, let us rewrite the optimization problem (13) for $a = 0$ given that $M_0^{VA} = \Phi^{+VA}$:

$$\begin{aligned} \min_{\rho_{\pm}} \quad & \text{tr}[\rho_-] \\ \text{s.t.} \quad & \text{tr}_{VA}[(\Phi^{+VA} \otimes \mathbb{1}^B)(\omega_x^V \otimes \rho^{AB})] \\ & = \text{tr}_{VA}\{(\Phi^{+VA} \otimes \mathbb{1}^B)[\omega_x^V \otimes (\rho_+^{AB} - \rho_-^{AB})]\}, \\ & \sigma_{0|\omega_x}^{\pm} = \text{tr}_V[M_{0,\pm}^{*VB}(\omega_x^V \otimes \mathbb{1}^B)], \\ & M_{0,\pm}^{*VB} = \text{tr}_A[(\Phi^{+VA} \otimes \mathbb{1}^B)(\mathbb{1}^V \otimes \rho_{\pm}^{AB})] \\ & = \frac{1}{d}\rho_{\pm}^{TB} \geq 0. \end{aligned} \quad (14)$$

The first constraint in case of a tomographically complete set of inputs is satisfied if and only if

$$\rho_{AB} = \rho_+ - \rho_-,$$

which finally reduces (14) to (9). For the other values of a the constraints from (14) are automatically satisfied. The first constraint can be rewritten as

$$\begin{aligned} \text{tr}_{VA}[(U_a^V \Phi^{+VA} U_a^{\dagger V} \otimes \mathbb{1}^B)(\omega_x^V \otimes \rho^{AB})] \\ = \text{tr}_{VA}\{(U_a^V \Phi^{+VA} U_a^{\dagger V} \otimes \mathbb{1}^B)[\omega_x^V \otimes (\rho_+^{AB} - \rho_-^{AB})]\}, \end{aligned}$$

which is equivalent to

$$\begin{aligned} \text{tr}_{VA}[(\Phi^{+VA} \otimes \mathbb{1}^B)(U_a^{\dagger V} \omega_x^V U_a^V \otimes \rho^{AB})] \\ = \text{tr}_{VA}\{(\Phi^{+VA} \otimes \mathbb{1}^B)[U_a^{\dagger V} \omega_x^V U_a^V \otimes (\rho_+^{AB} - \rho_-^{AB})]\}. \end{aligned}$$

If the set $\{\omega_x\}_x$ is tomographically complete, so is $\{U_a^{\dagger} \omega_x U_a\}_x$, and thus the last statement is equivalent to $\rho_{AB} = \rho_+ - \rho_-$. Similarly the last constraint from (14) reduces to $U_a \rho_{\pm}^{TB} U_a^{\dagger} \geq 0$, which is satisfied if $\rho_{\pm}^{TB} \geq 0$. Thus, we see that when Alice applies the full Bell state measurement and has access to a tomographically complete set of input states, the optimization problems (13) and (9) are equivalent.

In the general case, note that the states ρ'_{\pm} leading to the optimal solution of (9) by forming $\sigma_{a|\omega_x}^{\pm} = \text{tr}_{VA}[(M_a^{VA} \otimes \mathbb{1}^B)(\omega_x^V \otimes \rho'^{AB})]$ with arbitrary measurements $\{M_a^{VA}\}$ and input states $\{\omega_x\}_x$ satisfy all the constraints of (13). The equivalence between the objective functions follows from (10) and the last constraint is satisfied due to (12). This means that the solution to (13) cannot be larger than $\mathcal{N}(\rho^{AB})$, i.e., it places a lower bound on the entanglement negativity of ρ^{AB} .

B. Estimating entanglement negativity from violations of nonclassical teleportation witnesses

In [9] we introduced an SDP optimization problem which determines if observed teleportation data can be reproduced with classical teleportation channels. The dual form of the introduced SDP gives a nonclassical teleportation witness, i.e., an operator which is positive whenever evaluated on Bob's states obtained from a classical teleportation experiment, but can take a negative value when evaluated on states resulting from a nonclassical teleportation. Violation of nonclassical teleportation witness was used to experimentally certify nonclassicality of a teleportation protocol in [10]. In this subsection we show that teleportation witnesses, besides certifying nonclassical teleportation, can be used to put a lower bound on entanglement negativity of the shared state. The use of teleportation witnesses may be favorable since knowing the full teleportation assemblage requires performing quantum state tomography which is often a costly task. Contrarily, one can obtain the violation of a teleportation witness without performing the full tomography of the teleported states.

Let us observe that the average teleportation fidelity represents a particular type of teleportation witness, and remind again that the authors of [14] proved that it lower bounds entanglement negativity. Here we provide an SDP which allows us to estimate entanglement negativity from violation of an arbitrary teleportation witness $F_{a|\omega_x}$. Assuming that the observed violation of the witness is w , the following SDP provides a lower bound on the negativity of the shared state

$$\begin{aligned} \mathcal{N}(\rho^{AB}) \geq f(w) = \min_{\{M_{a,\pm}^*\}_a, \rho_{\pm}^B} \quad & \text{tr}[\rho_-^B] \\ \text{s.t.} \quad & w = \sum_{a,x} \text{tr}[(\omega_x^V \otimes F_{a|\omega_x}^B)(M_{a+}^{*VB} - M_{a-}^{*VB})], \\ & \sum_a M_{a,\pm}^{*VB} = \mathbb{1}^V \otimes \rho_{\pm}^B, \\ & M_{a,\pm}^{*VB} \geq 0 \quad \forall a. \end{aligned} \quad (15)$$

This optimization problem is equivalent to (13). It looks for the state with minimum negativity that could have led to the given violation of the nonclassical teleportation witness.

III. ESTIMATING ENTANGLEMENT ROBUSTNESS FROM TELEPORTATION EXPERIMENTS

Entanglement negativity is not the only entanglement measure that can be estimated from a teleportation experiment. In this section we turn our attention to entanglement robustness and show how it can be inferred from full data accessible in a teleportation experiment. Moreover, we show that the lower bound can have a meaning of a quantifier of nonclassicality of teleportation.

An intuitive way to quantify nonclassicality (e.g., entanglement, EPR steering, nonlocality, etc.) is in terms of its robustness to noise. Such robustness measures are expressed as the maximal amount of noise which can be added to the given object before it becomes classical. Specifying the type of added noise allows for different types of robustness to be defined. Entanglement robustness [15] is a well known example: for a bipartite state ρ^{AB} the entanglement robustness is defined through the following optimization problem:

$$\begin{aligned} \epsilon(\rho^{AB}) &= \min_{r, \rho_s, \sigma_s} r \\ \text{s.t.} \quad & \frac{\rho^{AB} + r\rho_s}{1+r} = \sigma_s, \\ & \sigma_s \in \mathcal{S}, \end{aligned} \quad (16)$$

Depending on the properties of ρ_s , different types of entanglement robustness can be defined:

(i) *generalized entanglement robustness* [16] ϵ_{gen} , obtained when the only constraint is that ρ_s is a valid quantum state.

(ii) *separable entanglement robustness* ϵ_{sep} , obtained when the state ρ_s is separable.

(iii) *random entanglement robustness* ϵ_r , obtained when the state ρ_s is maximally mixed $\rho_s = \frac{\mathbb{1}}{d^2}$.

Based on the inclusion relations between the sets of states to which ρ_s belongs, it follows that

$$\epsilon_{\text{gen}} \leq \epsilon_{\text{sep}} \leq \epsilon_r.$$

Anticipating its role in estimating entanglement robustness let us define teleportation robustness. The central object in a teleportation experiment is the teleportation assemblage $\{\sigma_{a|\omega_x}\}_{a,x}$. Robustness of teleportation represents the maximal proportion of a “noise assemblage” with which $\{\sigma_{a|\omega_x}\}_{a,x}$ can be mixed before it becomes classical:

$$\begin{aligned} \tau(\{\sigma_{a|\omega_x}\}) &= \max_{r, \{\bar{\sigma}_{a|\omega_x}\}, \{M_a^*\}} r \\ \text{s.t.} \quad & \frac{\sigma_{a|\omega_x}^B + r\bar{\sigma}_{a|\omega_x}^B}{1+r} = \text{tr}_V[M_a^{*VB}(\omega_x^V \otimes \mathbb{1}^B)], \\ & \sum_a M_a^{*VB} = \mathbb{1}^V \otimes \frac{\sum_a \sigma_{a|\omega_x}^B + r \sum_a \bar{\sigma}_{a|\omega_x}^B}{1+r}, \\ & M_a^{*VB} \in \mathcal{S} \quad \forall a. \end{aligned} \quad (17)$$

The constraints on the noise assemblage $\{\bar{\sigma}_{a|\omega_x}\}_{a,x}$ determine different types of teleportation robustness:

(i) *generalized teleportation robustness* τ_{gen} , obtained when the only constraint on $\{\bar{\sigma}_{a|\omega_x}\}_{a,x}$ is that it is allowed by quantum theory.

(ii) *classical teleportation robustness* τ_{cl} , obtained when $\{\bar{\sigma}_{a|\omega_x}\}_{a,x}$ describes classical teleportation.

(iii) *random teleportation robustness* τ_r , obtained when each element of $\{\bar{\sigma}_{a|\omega_x}\}_{a,x}$ is proportional to the maximally mixed state.

Each type of teleportation robustness places a lower bound on the corresponding type of entanglement robustness of the shared state. This is to be expected, since it has previously been shown that it is possible to place lower bounds on different entanglement quantifiers of the shared state in a measurement-device-independent manner [17,18]. Since we have already established teleportation as corresponding to the one-sided measurement-device-independent scenario, it seems natural that teleportation quantifiers are also related to the entanglement quantifiers of the shared state ρ^{AB} [see (3)]. Moreover, we can show that if Alice applies a full Bell state measurement and the set of input states is tomographically complete, then each teleportation robustness equals the corresponding entanglement robustness of the shared state ρ^{AB} . Since the proofs are similar in spirit to those used for lower bounding entanglement negativity, we leave the detailed proofs for the Appendix A.

IV. TELEPORTATION WEIGHT AND BEST SEPARABLE APPROXIMATION

In the previous section we saw that the lower bound on entanglement robustness inferred from a teleportation experiment can be seen as a teleportation quantifier. In this section we start from the opposite direction, i.e., we first define a teleportation quantifier and then we show that it puts a lower bound on the corresponding entanglement measure. An operationally meaningful teleportation quantifier different to robustness-based quantifiers is the teleportation weight. Any teleportation assemblage can be written as a convex combination of an assemblage obtained via classical teleportation and a nonclassical one. The minimal proportion of the nonclassical teleportation assemblage defines the teleportation weight. It can be seen as an analog to the best separable approximation [19], EPR2 decomposition [20], and the steering weight [21]. Mathematically, we define the teleportation weight in the following way:

$$\begin{aligned} \text{TW}(\{\sigma_{a|\omega_x}^B\}) &= \min_{p, \{\bar{M}_a\}, \{\tilde{M}_a\}} p \\ \text{s.t.} \quad & \sigma_{a|\omega_x}^B = \text{tr}_V[(p\tilde{M}_a^{VB} + (1-p)\bar{M}_a^{VB})\omega_x^V \otimes \mathbb{1}^B] \\ & \sum_a \text{tr}_V[\tilde{M}_a^{VB}(\omega_x^V \otimes \mathbb{1}^B)] \\ & = \sum_a \text{tr}_V[\bar{M}_a^{VB}(\omega_{x'}^V \otimes \mathbb{1}^B)], \quad \forall x, x', \\ & \bar{M}_a^{VB} \geq 0, \quad \bar{M}_a^{VB} \in \mathcal{S}, \quad \forall a, \\ & (\tilde{M}_a^{VB})^{T_V} \geq 0, \quad \forall a. \end{aligned} \quad (18)$$

In this definition the channel operators \bar{M}_a^{VB} describe classical teleportation, which is why they have to be positive and

separable, while \tilde{M}_a^{VB} are channel operators corresponding to nonclassical teleportation, satisfying instead the constraint of the positivity of the partial transpose [see (A28)]. Nonzero teleportation weight witnesses that teleportation is nonclassical, which in turn means that the state Alice and Bob share is entangled. When the set of input states is tomographically complete and the state Alice and Bob share is maximally entangled the teleportation weight must be equal to 1. Moreover, any pure entangled shared state with tomographically complete set of input states has maximal teleportation weight.

Just as teleportation robustness quantifiers can be seen to provide bounds on the corresponding entanglement robustness quantifiers, so too does the teleportation weight of the teleportation assemblage $\{\sigma_{a|\omega_x}\}_{a,x}$ place a lower bound on the best separable approximation of the state ρ^{AB} . The best separable approximation of a bipartite state ρ^{AB} is a monotone which says how much of a separable state is contained in the state ρ^{AB} and is defined as

$$\begin{aligned} \epsilon_{\text{BSA}}(\rho^{\text{AB}}) &= \min_{p, \rho_s, \sigma_s} p \\ \text{s.t. } \rho^{\text{AB}} &= p\rho_s + (1-p)\sigma_s, \\ \sigma_s &\in \mathcal{S}, \end{aligned} \quad (19)$$

We leave the proof that $\text{TW}(\{\sigma_{a|\omega_x}^{\text{B}}\}) \leq \epsilon_{\text{BSA}}(\rho^{\text{AB}})$ for the Appendix B. We also show in the Appendix B that in case Alice has a tomographically complete set of input states and performs a Bell state measurement, the resulting teleportation assemblage has teleportation weight equal to the best separable approximation of the state Alice shares with Bob.

V. EXAMPLES

In this section we present several examples of the methods presented in this work. An online notebook reproducing all of the figures presented in this section can be found at [22].

Negativity estimation—In Fig. 1 we plot the teleportation-based lower bound on the negativity for the state $p|\Phi^+\rangle\langle\Phi^+| + (1-p)|01\rangle\langle 01|$, when Alice performs a full Bell state measurement and uses a tomographically complete set of input states (eigenstates of the Pauli operators). The graph was obtained by using the SDP (13). We see that, as expected, the lower bound is tight, and equal to the negativity of the state.

Robustness estimations—In Fig. 2 we show a graph showing different types of teleportation robustness, i.e., lower bounds to the corresponding types of entanglement robustness of teleportation assemblages produced by using state $\rho = p|\Phi^+\rangle\langle\Phi^+| + (1-p)|01\rangle\langle 01|$. The graphs are obtained by performing SDP optimizations (17) with different types of noise.

Teleportation weight—The teleportation weight of the state $p|\Phi^+\rangle\langle\Phi^+| + (1-p)\frac{\mathbb{1}}{4}$ corresponding to different scenarios (i.e., different sets of input states) is presented in Fig. 3. The graph is obtained by solving the SDP formulation of (18). We see that the teleportation weight for a tomographically complete set of input states is larger than zero whenever $p > \frac{1}{3}$, which is the separability bound for Werner states. This does not change even if Alice does not apply the full Bell state measurements, but projects only onto one of the Bell




FIG. 1. Teleportation-based lower bound on the negativity, and negativity of the state $p|\Phi^+\rangle\langle\Phi^+| + (1-p)|01\rangle\langle 01|$. Alice performs a full Bell state measurement and uses a tomographically complete set of input states. In this case, the lower bound on the negativity is tight, so the curves for the negativity and the negativity lower bound coincide.

states (i.e., a partial Bell state measurement). When the set of input states consists of eigenstates of two Pauli observables, nonclassical teleportation is detected only when $p > \frac{1}{2}$.

Bound entangled states—One of the most striking new insights resulting from using all the observable data in a teleportation experiment is that all entangled states can be used to certify nonclassical teleportation. Previously, all bound entangled states were considered to be useless for teleportation.




FIG. 2. Teleportation robustnesses for the state $\rho = p|\Phi^+\rangle\langle\Phi^+| + (1-p)|01\rangle\langle 01|$. The set of quantum inputs consists of all eigenstates of three Pauli operators. For this particular teleportation assemblage the generalized and classical teleportation robustness coincide. We can see that even when the average teleportation fidelity is smaller than $2/3$ the robustness quantifiers are larger than zero, demonstrating nonclassical teleportation.




FIG. 3. Teleportation weight for different scenarios involving the state $p|\Phi^+\rangle\langle\Phi^+| + (1-p)\frac{\mathbb{1}}{4}$: Alice either performs a full or partial Bell state measurement, and uses either a tomographically complete set of inputs (eigenstates of X , Y , and Z), or a tomographically incomplete set of measurements (eigenstates of X and Z). The teleportation weight is insensitive to the choice of measurements for both sets of inputs, indicating that it is only the conclusive events (corresponding to POVM elements that are entangled) that count.

One of the most famous examples of bound entangled states is the Horodecki state [23]:

$$\rho_H = \frac{1}{8a+1} \begin{pmatrix} a & 0 & 0 & 0 & a & 0 & 0 & 0 & a \\ 0 & a & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & a & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & a & 0 & 0 & 0 & 0 & 0 \\ a & 0 & 0 & 0 & a & 0 & 0 & 0 & a \\ 0 & 0 & 0 & 0 & 0 & a & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \frac{1+a}{2} & 0 & \frac{\sqrt{1-a^2}}{2} \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & a & 0 \\ a & 0 & 0 & 0 & a & 0 & \frac{\sqrt{1-a^2}}{2} & 0 & \frac{1+a}{2} \end{pmatrix}, \quad (20)$$

for values $a \in (0, 1)$. The dependence of the teleportation weight of the teleportation assemblage obtained by using the Horodecki state with parameter a is given in Fig. 4. The set of input states is chosen to be tomographically complete and a partial Bell state measurements is performed ($M_1^{VA} = |\Phi^+\rangle\langle\Phi^+|$, $M_2^{VA} = \mathbb{1} - M_1^{VA}$). The teleportation weight of the Horodecki state is small in value, but we observed that other bound entangled states give higher weights, even maintaining a partial Bell state measurement. For example, the ‘‘pyramid’’ UPB (unextendable product bases) state [24], with tomographically complete set of inputs, has teleportation weight equal to 0.2350.

VI. CONCLUSION

In this work we have discussed the estimation of entanglement from teleportation experiments and closely related problem of quantifying nonclassical teleportation. Taking into account the central role of quantum teleportation in various quantum information protocols, it is very important to improve our understanding of the role of entanglement in the protocol, both qualitatively and quantitatively. We showed




FIG. 4. Dependence of the teleportation quantifiers introduced here on parameter a the Horodecki state, using a tomographically complete set of input states (chosen randomly to produce this plot), and a partial Bell state measurement. For all values of $a \neq 0$ or 1, nonclassical teleportation is demonstrated. The curves for the classical and the generalized teleportation robustness coincide. Separability of the channel operators was relaxed to the requirement of having a 2-symmetric PPT extension [12].

that by using the full data available in an experiment we can put lower bounds on several entanglement measures, such as entanglement negativity, robustness of entanglement, and the best separable approximation. Importantly, in accordance with our previous work [9] we have shown that we can estimate nonzero entanglement even from teleportation experiments in which the average fidelity of teleportation falls below the classical limit. The inferred lower bound appears to be tight in case when Alice applies a full Bell state measurements and the set of input states is tomographically complete.

It is worth mentioning that lower bounds on different entanglement measures can actually be seen as teleportation quantifiers. We thus paid additional attention to operationally meaningful robustness based teleportation quantifiers, and the teleportation weight.

There are several possible directions for future research. One would be to identify other known entanglement measures which can be estimated from a teleportation experiment. A more demanding task would be to mirror the protocol of certifying and estimating nonclassicality of teleportation to some other protocols which use teleportation as a subroutine. Finally, while all our results regard finite-dimensional teleportation it would be interesting to understand what can be said about continuous variable teleportation.

ACKNOWLEDGMENTS

This work was supported by the Swiss National Science Foundation (Starting grant DIAQ), the Ramón y Cajal fellowship, Spanish MINECO (QIBEQI FIS2016-80773-P and Severo Ochoa SEV-2015-0522), the AXA Chair in Quantum Information Science, Generalitat de Catalunya (SGR875 and CERCA Programme), Fundació Privada Cellex and ERC CoG

QITBOX, COST project CA16218 NANOCOBYBRI, and a Royal Society University Research Fellowship (UHQT).

APPENDIX A: LOWER BOUNDS ON ENTANGLEMENT ROBUSTNESS

The entanglement present in the shared state ρ^{AB} is a necessary resource for nonclassicality of teleportation assemblage $\{\sigma_{a|\omega_x}\}_{a,x}$. Given robustness-based nonclassicality measures for both the state ρ^{AB} and the teleportation assemblage $\{\sigma_{a|\omega_x}\}_{a,x}$, it is instructive to compare their values.

In this paper we prove the following relations:

$$\tau_{\text{gen}}(\{\sigma_{a|\omega_x}\}) \leq \epsilon_{\text{gen}}(\rho^{AB}), \quad (\text{A1})$$

$$\tau_{\text{cl}}(\{\sigma_{a|\omega_x}\}) \leq \epsilon_{\text{sep}}(\rho^{AB}), \quad (\text{A2})$$

$$\tau_{\text{r}}(\{\sigma_{a|\omega_x}\}) \leq \epsilon_{\text{r}}(\rho^{AB}). \quad (\text{A3})$$

All inequalities are saturated when Alice teleports a tomographically complete set of inputs and performs a Bell state measurement.

1. Generalized teleportation robustness

Let us examine in detail the generalized teleportation robustness of a teleportation assemblage $\{\sigma_{a|\omega_x}\}_{a,x}$, obtained by using the measurement $\{M_a\}$ on a shared state ρ^{AB} , and a set of input states $\{\omega_x\}_x$. This quantity is given by (17) when the only constraint on $\{\bar{\sigma}_{a|\omega_x}\}_{a,x}$ is that it is admissible by quantum theory:

$$\begin{aligned} \tau_{\text{gen}}(\{\sigma_{a|\omega_x}\}) &= \min_{r, \{M_a^*\}, \{\bar{\sigma}_{a|\omega_x}\}} r \\ \text{s.t. } \frac{\sigma_{a|\omega_x}^B + r\bar{\sigma}_{a|\omega_x}^B}{1+r} &= \text{tr}_V [M_a^{*VB}(\omega_x^V \otimes \mathbb{1}^B)], \\ \bar{\sigma}_{a|\omega_x}^B &\in T_q, \\ \sum_a M_a^{*VB} &= \mathbb{1}^V \otimes \frac{\sum_a \sigma_{a|\omega_x} + r \sum_a \bar{\sigma}_{a|\omega_x}}{r+1}, \\ M_a^{*VB} &\geq 0, \quad M_a^{*VB} \in \mathcal{S}, \quad \forall a, \end{aligned} \quad (\text{A4})$$

where T_q is the set of teleportation assemblages admissible by quantum theory, i.e., those which can arise by performing a quantum measurement on a shared quantum state (this requirement will be made more explicit in the next section of the Appendix in Eq. (A28)). The set of constraints given above imposes that the mixture of the observed assemblage $\{\sigma_{a|\omega_x}\}_{a,x}$ with some other hypothetical assemblage $\{\bar{\sigma}_{a|\omega_x}\}_{a,x}$ can be simulated classically. We note that all constraints can be written in a linear form—with a suitable relaxation of the set of separable operators this problem is readily solved by using semidefinite programming optimization (SDP) [13]. The only nontrivial constraint regards the characterization of the set T_q , but in the next section we will show that membership in such a set also can be imposed as a semidefinite programming constraint.

From the definition of the generalized entanglement robustness we know that

$$\frac{\rho^{AB} + \epsilon_{\text{gen}}(\rho^{AB})}{1 + \epsilon_{\text{gen}}(\rho^{AB})} = \sigma_S, \quad (\text{A5})$$

for some state ρ_S and a separable state σ_S . By tensoring with ω_x^V and applying measurement $\{M_a^{VA}\}_a$, (A5) becomes

$$\begin{aligned} &\frac{1}{1 + \epsilon_{\text{gen}}(\rho^{AB})} \text{tr}_{VA} [(M_a^{VA} \otimes \mathbb{1}^B)(\omega_x^V \otimes \rho^{AB})] \\ &+ \frac{\epsilon_{\text{gen}}(\rho^{AB})}{1 + \epsilon_{\text{gen}}(\rho^{AB})} \text{tr}_{VA} [(M_a^{VA} \otimes \mathbb{1}^B)(\omega_x^V \otimes \rho_S^{AB})] \\ &= \text{tr}_{VA} [(M_a^{VA} \otimes \mathbb{1}^B)(\omega_x^V \otimes \sigma_S^{AB})], \end{aligned}$$

for every x and a . This is equivalent to

$$\frac{\sigma_{a|\omega_x} + \epsilon_{\text{gen}}(\rho^{AB})\bar{\sigma}_{a|\omega_x}}{1 + \epsilon_{\text{gen}}(\rho^{AB})} = \text{tr}_V [M_a^{*VB}(\omega_x^V \otimes \mathbb{1}^B)], \quad (\text{A6})$$

where

$$\bar{\sigma}_{a|\omega_x} = \text{tr}_{VA} [(M_a^{VA} \otimes \mathbb{1}^B)(\omega_x^V \otimes \rho_S^{AB})]$$

is an arbitrary teleportation assemblage (for an arbitrary state ρ_S). At the same time,

$$M_a^{*VB} = \text{tr}_A [(M_a^{VA} \otimes \mathbb{1}^B)(\mathbb{1}^V \otimes \sigma_S^{AB})] \quad (\text{A7})$$

is separable when σ_S is separable. Furthermore,

$$\begin{aligned} \sum_a M_a^{*VB} &= \text{tr}_A \left[\left(\sum_a M_a^{VA} \otimes \mathbb{1}^B \right) (\mathbb{1}^V \otimes \sigma_S^{AB}) \right] \\ &= \text{tr}_A (\mathbb{1}^V \otimes \sigma_S^{AB}) \\ &= \mathbb{1}^V \otimes \frac{\text{tr}_A \rho^{AB} + \epsilon_{\text{gen}}(\rho^{AB}) \text{tr}_A \rho_S^{AB}}{1 + \epsilon_{\text{gen}}(\rho^{AB})} \\ &= \mathbb{1}^V \otimes \frac{\sum_a \sigma_{a|\omega_x} + \epsilon_{\text{gen}}(\rho^{AB}) \sum_a \bar{\sigma}_{a|\omega_x}}{1 + \epsilon_{\text{gen}}(\rho^{AB})}, \end{aligned} \quad (\text{A8})$$

where the second line follows from the completeness relation $\sum_a M_a = \mathbb{1}$, the third line follows from (A5), and the last line is obtained by using the definitions of $\sigma_{a|\omega_x}$ and $\bar{\sigma}_{a|\omega_x}$.

From (A6), (A8), and the separability of M_a^{*VB} , it follows that $\epsilon_{\text{gen}}(\rho^{AB})$ satisfies all the constraints from (A4). Since the mixing teleportation assemblage $\bar{\sigma}_{a|\omega_x}$ did not have any special properties, apart from being realizable in quantum theory, the generalized teleportation robustness of $\{\sigma_{a|\omega_x}\}_{a,x}$ cannot be bigger than $\epsilon_{\text{gen}}(\rho^{AB})$:

$$\tau_{\text{gen}}(\{\sigma_{a|\omega_x}\}) \leq \epsilon_{\text{gen}}(\rho^{AB}). \quad (\text{A9})$$

2. Classical teleportation robustness

The classical teleportation robustness $\tau_{\text{cl}}(\cdot)$ is defined by (17) with the additional constraint that the mixing teleportation assemblage describes classical teleportation. Such a teleportation assemblage is characterized by positive and separable channel operators \bar{M}_a^{VB} as shown in (5) and (8). With these constraints, the classical teleportation robustness can be expressed as the following optimization problem:

$$\begin{aligned} \tau_{\text{cl}}(\{\sigma_{a|\omega_x}\}) &= \min_{\{r_a\}, \{\bar{M}_a\}, \{M_a^*\}} r \\ \text{s.t. } \frac{\sigma_{a|\omega_x}^B + r\bar{\sigma}_{a|\omega_x}^B}{1+r} &= \text{tr}_V [M_a^{*VB}(\omega_x^V \otimes \mathbb{1}^B)], \\ \bar{\sigma}_{a|\omega_x}^B &= \text{tr}_V [(\bar{M}_a^{VB})(\omega_x^V \otimes \mathbb{1}^B)] \quad \forall a, x, \end{aligned}$$

$$\begin{aligned}
 \bar{M}_a^{VB}, M_a^{*VB} &\geq 0, \quad \forall a, \\
 \bar{M}_a^{VB}, M_a^{*VB} &\in \mathcal{S}, \quad \forall a, \\
 \sum_a \bar{M}_a^{VB} &= \mathbb{1}^V \otimes \sum_a \bar{\sigma}_{a|\omega_x}, \\
 \sum_a M_a^{*VB} &= \mathbb{1}^V \otimes \frac{\sum_a \sigma_{a|\omega_x} + r \sum_a \bar{\sigma}_{a|\omega_x}}{1+r}. \quad (\text{A10})
 \end{aligned}$$

The classical teleportation robustness mirrors the classical entanglement robustness $\epsilon_{\text{sep}}(\rho^{AB})$ which satisfies the following equation:

$$\frac{\rho^{AB} + \epsilon_{\text{sep}}(\rho^{AB})\rho_S}{1 + \epsilon_{\text{cl}}(\rho^{AB})} = \sigma_S, \quad (\text{A11})$$

where now both states ρ_S and σ_S are separable. By tensoring with ω_x^V and applying the measurement $\{M_a^{VA}\}_a$ this becomes

$$\frac{\sigma_{a|\omega_x} + \epsilon_{\text{sep}}(\rho^{AB})\bar{\sigma}_{a|\omega_x}}{1 + \epsilon_{\text{sep}}(\rho^{AB})} = \text{tr}_V [M_a^{*VB}(\omega_x^V \otimes \mathbb{1}^B)], \quad (\text{A12})$$

where now the mixing assemblage can be expressed in terms of separable channel operators \bar{M}_a^{VB} ,

$$\begin{aligned}
 \bar{\sigma}_{a|\omega_x} &= \text{tr}_V [\bar{M}_a^{VB}(\omega_x^V \otimes \mathbb{1}^B)], \\
 \bar{M}_a^{VB} &= \text{tr}_A [(M_a^{VA} \otimes \mathbb{1}^B)(\mathbb{1}^V \otimes \rho_S^{AB})], \\
 \sum_a \bar{M}_a^{VB} &= \mathbb{1}^V \otimes \sum_a \bar{\sigma}_{a|\omega_x}
 \end{aligned}$$

The channel operators M_a^{*VB} remain separable and still satisfy relation (A8). Together with (A12) and the separability of \bar{M}_a^{VB} , this implies that $\epsilon_{\text{sep}}(\rho^{AB})$ satisfies all of the constraints of the classical teleportation robustness (of the teleportation assemblage $\{\sigma_{a|\omega_x}\}_{a,x}$), leading to

$$\tau_{\text{cl}}(\{\sigma_{a|\omega_x}\}) \leq \epsilon_{\text{sep}}(\rho^{AB}). \quad (\text{A13})$$

3. Random teleportation robustness

In [9] the random teleportation robustness was introduced as a special case of (17) with the constraint $\bar{\sigma}_{a|\omega_x} = \frac{1}{|o|} \frac{\mathbb{1}}{d}$, where $|o|$ is the number of outcomes of Alice’s measurement. Here we consider a more general definition, defined as the solution to the following optimization problem:

$$\begin{aligned}
 \tau_r(\{\sigma_{a|\omega_x}\}) &= \min_{r, \{p(a)\}, \{M_a^*\}} r \\
 \text{s.t.} \quad &\frac{\sigma_{a|\omega_x}^B + r p(a) \frac{\mathbb{1}^B}{d}}{1+r} = \text{tr}_V [M_a^{*VB}(\omega_x^V \otimes \mathbb{1}^B)], \\
 &\sum_a p(a) = 1, \quad M_a^{*VB} \geq 0, \quad M_a^{*VB} \in \mathcal{S}, \\
 &\sum_a M_a^{*VB} = \mathbb{1}^V \otimes \frac{\rho^B + r \frac{\mathbb{1}^B}{d}}{1+r}. \quad (\text{A14})
 \end{aligned}$$

Recall that the random entanglement robustness $\epsilon_r(\rho^{AB})$ satisfies

$$\frac{\rho^{AB} + \epsilon_r(\rho^{AB})\mathbb{1}/d^2}{1 + \epsilon_r(\rho^{AB})} = \sigma_S. \quad (\text{A15})$$

Analogously to the previous cases this equation implies

$$\frac{\sigma_{a|\omega_x}^B + \epsilon_r(\rho^{AB})\mathbb{1}/d}{1 + \epsilon_r(\rho^{AB})} = \text{tr}_V [M_a^{*VB}(\omega_x^V \otimes \mathbb{1}^B)],$$

where

$$p(a) = \text{tr} \left[M_a^{VA} \left(\omega_x^V \otimes \frac{\mathbb{1}^A}{d} \right) \right] \quad (\text{A16})$$

and M_a^{*VB} satisfies (A7) and (A8) when $\sum_a \bar{\sigma}_{a|\omega_x} = \frac{\mathbb{1}}{d}$. Note that (A16) confirms that $\sum_a p(a) = 1$. We have confirmed that $\epsilon_r(\rho^{AB})$ satisfies all the constraints from (A14), thus represents an upper bound to the random teleportation robustness

$$\tau_r(\{\sigma_{a|\omega_x}\}) \leq \epsilon_r(\rho^{AB}). \quad (\text{A17})$$

4. Tightness of the lower bound

In [9] using a form of random teleportation robustness, it was proven that every entangled state leads to nonclassical teleportation. This was done by proving that the random entanglement robustness of ρ^{AB} is proportional to the random teleportation robustness of the teleportation assemblage $\{\sigma_{a|\omega_x}\}_{a,x}$ that results from applying a (full or partial) Bell state measurement on ρ^{AB} and a tomographically complete set of input states $\{\omega_x\}_x$.

In this Appendix we prove that such an equivalence is true for each kind of teleportation/entanglement robustness and, moreover, whenever Alice applies the full Bell state measurement the inequalities (A1)–(A3) are saturated.

Before doing so, let us state a lemma which will be repeatedly used in these proofs.

Lemma 1. Every element of a teleportation assemblage $\{\sigma_{a|\omega_x}\}_{a,x}$ resulting from an arbitrary measurement M_a^{VA} and a shared state ρ^{AB} could have also been obtained, up to a multiplicative factor, by post-selecting on the measurement outcome $\Phi^{+VA} = |\Phi^+\rangle\langle\Phi^+|^{VA}$ applied to a suitable state $\rho_a'^{AB}$.

Proof. The identity (11) allows any member of a teleportation assemblage $\{\sigma_{a|\omega_x}\}_{a,x}$ to be written in the following way:

$$\begin{aligned}
 \sigma_{a|\omega_x}^B &= \text{tr}_{VA} [(M_a^{VA} \otimes \mathbb{1}^B)(\omega_x^V \otimes \rho^{AB})] \\
 &= d \text{tr}_{VV_1A} \{ [(\omega_x^V)^T \otimes M_a^{V_1A} \otimes \mathbb{1}^B] (\Phi^{+VV_1} \otimes \rho^{AB}) \} \\
 &= d \text{tr}_{VV_1A} \{ [(\omega_x^V)^T \otimes \mathbb{1}^{V_1AB}] (\mathbb{1}^V \otimes M_a^{V_1A} \otimes \mathbb{1}^B) \\
 &\quad \times (\Phi^{+VV_1} \otimes \rho^{AB}) \}. \quad (\text{A18})
 \end{aligned}$$

In case $M_a^{V_1A} = \Phi^{+V_1A}$ the previous equation reduces to

$$\sigma_{a|\omega_x}^B = \frac{1}{d} \text{tr}_V \{ [(\omega_x^V)^T \otimes \mathbb{1}^B] \rho^{VB} \}.$$

On the other hand, if $M_a^{V_1A}$ is not a Bell state measurement, (A18) reduces to

$$\begin{aligned}
 \sigma_{a|\omega_x}^B &= d \text{tr}_{VV_1} \{ [(\omega_x^V)^T \otimes \mathbb{1}^{V_1AB}] \rho_a'^{VV_1AB} \} \\
 &= d p(a) \text{tr}_V \{ [(\omega_x^V)^T \otimes \mathbb{1}^B] \rho_a'^{VB} \}, \quad (\text{A19})
 \end{aligned}$$

where

$$p(a) = \text{tr} [(\mathbb{1}^V \otimes M_a^{V_1A} \otimes \mathbb{1}^B)(\Phi^{+VV_1} \otimes \rho^{AB})] \quad (\text{A20})$$

and

$$\rho_a^{\prime\text{VB}} = \frac{1}{p(a)} \text{tr}_{\text{VA}}(\mathbb{1}^{\text{V}} \otimes M_a^{\text{VA}} \otimes \mathbb{1}^{\text{B}})(\Phi^{+\text{V}_1} \otimes \rho^{\text{AB}}).$$

The state $\rho_a^{\prime\text{VB}}$ can be obtained from the state ρ^{AB} through a stochastic local operation, which can be seen as a local version of entanglement swapping. To obtain ρ_a^{\prime} from ρ , Alice uses two auxiliary systems in the maximally entangled state, and applies the measurement M_a on one auxiliary system and her share of the state ρ^{AB} . After the measurement she discards the measured systems.

Finally, the expression given in (A19) can be written as

$$\begin{aligned} \sigma_{a|\omega_x}^{\text{B}} &= d^2 p(a) \text{tr}_{\text{VA}}[(\omega_x^{\text{V}} \otimes \mathbb{1}^{\text{AB}})(\Phi^{+\text{VA}} \otimes \mathbb{1}^{\text{B}})(\mathbb{1}^{\text{V}} \otimes \rho_a^{\prime\text{AB}})] \\ &= d^2 p(a) \text{tr}_{\text{VA}}[(\Phi^{+\text{VA}} \otimes \mathbb{1}^{\text{B}})(\omega_x^{\text{V}} \otimes \rho_a^{\prime\text{AB}})], \end{aligned} \quad (\text{A21})$$

which proves the lemma. The multiplicative factor, mentioned in the lemma, is equal to $d^2 p(a)$. ■

The assumption that Alice applies the Bell state measurement implicitly assumes that the dimension of Alice's reduced state $d_A = \text{tr}_{\text{B}}(\rho^{\text{AB}})$ is equal to the dimension of the input state d . In the general case, d_A can be different from d and in (A18)–(A21) there is no assumption about d_A .

5. Generalized teleportation robustness

Let us denote by $\tau_{\text{gen}}^*(\cdot)$ the generalized teleportation robustness of a teleportation assemblage obtained when Alice performs a Bell state measurement $\{M_a^{\text{VA}}\}$ and the set of input states $\{\omega_x\}_x$ is tomographically complete. In this case the first constraint from (A4) can be rewritten as

$$\begin{aligned} &\frac{\sigma_{a|\omega_x}^{\text{B}} + r \bar{\sigma}_{a|\omega_x}^{\text{B}}}{1+r} \\ &= \frac{1}{r+1} \text{tr}_{\text{VA}}[(U_a^{\text{A}} \Phi^{+\text{VA}} U_a^{\text{A}\dagger} \otimes \mathbb{1}^{\text{B}})(\omega_x^{\text{V}} \otimes \rho^{\text{AB}})] \\ &\quad + \frac{1}{r+1} \text{tr}_{\text{VA}}[(\Phi^{+\text{VA}} \otimes \mathbb{1}^{\text{B}})(\omega_x^{\text{V}} \otimes d^2 p(a) \rho_a^{\prime\text{AB}})] \\ &= \frac{1}{d} \text{tr}_{\text{V}} \left[\left(\frac{U_a^{\text{V}} \rho^{\text{VB}} U_a^{\text{V}\dagger} + r d^2 p(a) \rho_a^{\prime\text{VB}}}{r+1} \right)^{\text{T}_\text{V}} (\omega_x^{\text{V}} \otimes \mathbb{1}^{\text{B}}) \right] \\ &= \text{tr}_{\text{V}}[M_a^{*\text{VB}}(\omega_x^{\text{V}} \otimes \mathbb{1}^{\text{B}})]. \end{aligned} \quad (\text{A22})$$

The second line follows from the assumption that Alice applies a Bell state measurement and U_a are the local unitary transformations that shift between the different Bell states $M_a = U_a \Phi^{+\text{VA}} U_a^{\dagger}$. The third line uses Lemma 1 and (A21) to re-express the general teleportation assemblage $\{\bar{\sigma}_{a|\omega_x}\}_{a,x}$. The fourth line follows from the identity (11).

Given that the set of quantum inputs $\{\omega_x\}$ is tomographically complete the last equality implies

$$M_a^{*\text{VB}} = \frac{1}{d} \left(\frac{1}{r+1} U_a^{\text{V}} \rho^{\text{VB}} U_a^{\text{V}\dagger} + \frac{r}{1+r} d^2 p(a) \rho_a^{\prime\text{VB}} \right)^{\text{T}_\text{V}}. \quad (\text{A23})$$

With this in mind we can once more rewrite the optimization problem (A4):

$$\tau_{\text{gen}}^*(\{\sigma_{a|\omega_x}\}) = \min_{\{r, M_a^*, p(a), \rho_a^{\prime\text{VB}}\}} r$$

$$\text{s.t.} \quad \frac{U_a^{\text{V}} \rho^{\text{VB}} U_a^{\text{V}\dagger} + r d^2 p(a) \rho_a^{\prime\text{VB}}}{r+1} = d(M_a^{*\text{VB}})^{\text{T}_\text{V}}, \quad (\text{A24a})$$

$$M_a^{*\text{VB}} \geq 0, \quad M_a^{*\text{VB}} \in \mathcal{S} \quad \forall a, \quad (\text{A24b})$$

$$\sum_a p(a) \rho_a^{\prime\text{VB}} = \frac{\mathbb{1}^{\text{V}}}{d} \otimes \rho^{\text{B}}, \quad \sum_a p(a) = 1, \quad (\text{A24c})$$

$$\sum_a M_a^{*\text{VB}} = \mathbb{1}^{\text{V}} \otimes \frac{\rho^{\text{B}} + r \rho^{\text{B}}}{1+r}, \quad (\text{A24d})$$

which resembles the optimization problem defining the generalized entanglement robustness of the state $\rho^{\text{VB}} = \rho^{\text{AB}}$. Indeed the optimization problem (A24), for one specific value of a , say $a = 0$, for which $U_0 = \mathbb{1}$, is similar to (16), with the differences being that $d^2 p(a) \rho_a^{\prime}$ and $d M_a^{*\text{VB}}$ are not necessarily normalized, and the two additional constraints (A24c) and (A24d). The constraint (A24a), for $a = 0$, has a solution if $d \text{tr} M_0^{*\text{VB}} = [1 + r d^2 p(0)]/(1+r)$. Taking this into account the constraint can be rearranged in the following way:

$$\frac{\rho^{\text{VB}} + r d^2 p(0) \rho_0^{\prime\text{VB}}}{1 + r d^2 p(0)} = \frac{1}{\text{tr} M_0^{*\text{VB}}} (M_0^{*\text{VB}})^{\text{T}_\text{V}}, \quad (\text{A25})$$

which is now equivalent to the first constraint from (16). Thus, the minimal r satisfying this constraint for separable $M_0^{*\text{VB}}$ is equal to $\epsilon_{\text{gen}}(\rho^{\text{AB}})/[d^2 p(0)]$.

In a similar manner the minimal r satisfying (A24a) for $a \neq 0$ and separable $M_a^{*\text{VB}}$ is equal to $\epsilon_{\text{gen}}(\rho^{\text{AB}})/[d^2 p(a)]$, since the generalized entanglement robustness is the same for all states which are related by local unitary transformations. Let us, for a moment, suppose that there is at least one a such that $d^2 p(a) \geq 1$. Since there are d^2 different values of a , (A24c) implies that for some other value of a , say $a = a'$, it must be that $d^2 p(a') \leq 1$. But in this case the smallest r satisfying the constraints (A24a) and (A24b) for all a must be strictly bigger than $\epsilon_{\text{gen}}(\rho^{\text{AB}})$. On the other hand, if $d^2 p(a) = 1$ for all values of a , the smallest r satisfying (A24) is exactly equal to $\epsilon_{\text{gen}}(\rho^{\text{AB}})$. Since $p(a)$ are optimization variables, the minimal τ_{gen} will be achieved when all $p(a)$ s are mutually equal.

Finally, we have to make sure that the constraints (A24c) and (A24d) are satisfied by the solution $r = \epsilon_{\text{gen}}(\rho^{\text{VB}})$. If for $a = 0$ (A24a) is satisfied for some $\rho_0^{\prime\text{VB}}$ and $M_0^{*\text{VB}}$, for $a \neq 0$ it will be satisfied with the same r , $\rho_a^{\prime\text{VB}} = U_a^{\text{V}} \rho_0^{\prime\text{VB}} U_a^{\text{V}\dagger}$ and $M_a^{*\text{VB}} = U_a^{\text{V}} M_0^{*\text{VB}} U_a^{\text{V}\dagger}$ implying

$$\sum_a p(a) \rho_a^{\prime\text{VB}} = \frac{\mathbb{1}^{\text{V}}}{d} \otimes \rho^{\text{B}}$$

because $\sum_a U_a^{\text{V}} \rho_a^{\prime\text{VB}} U_a^{\text{V}\dagger} = d \mathbb{1}^{\text{V}} \otimes \rho^{\text{B}}$. The validity of (A24d) is verified by summing (A24a) over all different values

of a :

$$\begin{aligned} \sum_a M_a^{*VB} &= \frac{1}{d} \sum_a U_a^V \frac{(\rho^{VB})^{T_V} + r(\rho_0'^{VB})^{T_V}}{1+r} U_a^{\dagger V} \\ &= \mathbb{1}^V \otimes \frac{\rho^B + r\rho'^B}{1+r}. \end{aligned}$$

By establishing the equivalence between the optimization problem (A4) when Alice performs a full Bell state measurement and has access to a tomographically complete set of input states, and the optimization problem defining the generalized entanglement robustness, we can conclude that

$$\tau_{\text{gen}}^*(\{\sigma_{a|\omega_x}\}) = \epsilon_{\text{gen}}(\rho^{AB}). \quad (\text{A26})$$

6. Classical teleportation robustness

For easier comparison let us restate the definition of the separable entanglement robustness, which is obtained from (16) with the constraint that ρ_s is a separable state

$$\begin{aligned} \epsilon_{\text{sep}}(\rho^{AB}) &= \min_{r, \rho_s, \sigma_S} r \\ \text{s.t.} \quad &\frac{\rho^{AB} + r\rho_s}{1+r} = \sigma_S, \\ &\rho_s, \sigma_S \in \mathcal{S}. \end{aligned} \quad (\text{A27})$$

Let us further consider the classical teleportation robustness of a teleportation assemblage obtained when Alice applies a full Bell state measurement and uses a tomographically complete set of inputs, and denote it by $\tau_{\text{cl}}^*(\cdot)$. In order to reduce (A10) to (A27), it is useful to switch from the variables \bar{M}_a to $p(a)$ and ρ_a' which are related in the following way:

$$dp(a)(\rho_a'^{VB})^{T_V} = \bar{M}_a^{VB}. \quad (\text{A28})$$

With this in place, members of the teleportation assemblage $\bar{\sigma}_{a|\omega_x}^B$ can be written in the form given in (A21).

The simplification used in (A22) can again be used in exactly the same way, leading to

$$\begin{aligned} \text{tr}_V \left[\left(\frac{U_a^V \rho^{VB} U_a^{\dagger V} + rd^2 p(a) \rho_a'^{VB}}{r+1} \right)^{T_V} (\omega_x^V \otimes \mathbb{1}^B) \right] \\ = d \text{tr}_V [M_a^{*VB} (\omega_x^V \otimes \mathbb{1}^B)]. \end{aligned} \quad (\text{A29})$$

Since the set of input states is tomographically complete, (A29) implies

$$\frac{U_a^V \rho^{VB} U_a^{\dagger V} + rd^2 p(a) \rho_a'^{VB}}{r+1} = d(M_a^{*VB})^{T_V}.$$

However, in this case ρ_a' are separable [which is the consequence of (A28) and the separability of \bar{M}_a^{VB}]. It is thus the

case that the optimization (A10) reduces to

$$\tau_{\text{cl}}^*(\{\sigma_{a|\omega_x}\}) = \min_{r, \{\bar{M}_a, p(a)\rho_a'\}_a} r \quad (\text{A30a})$$

$$\text{s.t.} \quad \frac{U_a^V \rho^{VB} U_a^{\dagger V} + rd^2 p(a) \rho_a'^{VB}}{r+1} = d(M_a^{*VB})^{T_V}, \quad (\text{A30b})$$

$$M_a^{*VB} \geq 0, \quad M_a^{*VB} \in \mathcal{S} \quad \forall a, \quad (\text{A30c})$$

$$\rho_a'^{VB} \geq 0, \quad \rho_a'^{VB} \in \mathcal{S} \quad \forall a, \quad (\text{A30d})$$

$$\sum_a p(a) \rho_a'^{VB} = \frac{\mathbb{1}^V}{d} \otimes \bar{\rho}^B, \quad (\text{A30e})$$

$$\sum_a M_a^{*VB} = \mathbb{1}^V \otimes \frac{\rho^B + r\bar{\rho}^B}{1+r}. \quad (\text{A30f})$$

In order to emphasize the resemblance with (A27), let us rewrite (A30b) in the following way:

$$\frac{U_a^V \rho^{VB} U_a^{\dagger V} + rd^2 p(a) \rho_a'^{VB}}{1+rd^2 p_a} = \frac{1}{\text{tr} M_a^{*VB}} (M_a^{*VB})^{T_V}.$$

Since all states that are mutually related by local unitary transformations have the same value for the separable entanglement robustness, the smallest r satisfying the last equation for each a is equal to $\epsilon_{\text{cl}}/d^2 p(a)$. Analogously to the case of the generalized teleportation robustness, the optimal r is obtained when $p(a) = 1/d^2$ and $\rho_a' = U_a \rho_0' U_a^\dagger$ for all values of a , and is equal to $\epsilon_{\text{sep}}(\rho^{AB})$, which implies

$$\tau_{\text{cl}}^*(\{\sigma_{a|\omega_x}\}) = \epsilon_{\text{sep}}(\rho^{AB}). \quad (\text{A31})$$

7. Random teleportation robustness

Finally, we consider the random teleportation robustness of a teleportation assemblage obtained when Alice applies a full Bell state measurement and has access to a tomographically complete set of inputs. Let us denote it accordingly by $\tau_r^*(\cdot)$. We will compare it to the random entanglement robustness ϵ_r :

$$\begin{aligned} \epsilon_r(\rho^{AB}) &= \min_{r, \sigma_S} r \\ \text{s.t.} \quad &\frac{\rho^{AB} + \mathbb{1}/d^2}{1+r} = \sigma_S, \\ &\sigma_S \in \Sigma. \end{aligned} \quad (\text{A32})$$

Recall that the definition of random teleportation robustness of a teleportation assemblage $\{\sigma_{a|\omega_x}\}_{a,x}$ is given in (A14). The first constraint of (A14) in the case where Alice applies a Bell state measurement reads

$$\begin{aligned} \frac{\sigma_{a|\omega_x}^B + rp(a)\mathbb{1}^B/d}{1+r} \\ = \text{tr}_{VA} \left[(\Phi^{+VA} \otimes \mathbb{1}^B) \left(\omega_x^V \otimes \frac{U_a^A \rho^{AB} U_a^{\dagger A} + rp(a)\mathbb{1}^{AB}}{1+r} \right) \right] \\ = \frac{1}{d} \text{tr}_V \left[\left(\frac{U_a^V \rho^{VB} U_a^{\dagger V} + rp(a)\mathbb{1}^{VB}}{1+r} \right)^{T_V} (\omega_x^V \otimes \mathbb{1}^B) \right] \\ = \text{tr}_V [M_a^{*VB} (\omega_x^V \otimes \mathbb{1}^B)]. \end{aligned} \quad (\text{A33})$$

For a tomographically complete set of inputs this condition is satisfied if and only if

$$\frac{U_a^V \rho^{VB} U_a^{\dagger V} + r p(a) \mathbb{1}^{AB}}{1+r} = d(M_a^{*VB})^{T_V}. \quad (\text{A34})$$

Following this simplification, the optimization problem (A14) reduces to

$$\tau_r(\{\sigma_{a|\omega_x}\}) = \min_{r, \{M_a^*\}, \{p(a)\}} r$$

s.t. $\frac{U_a^V \rho^{VB} U_a^{\dagger V} + r p(a) \mathbb{1}^{VB}}{1+r} = d(M_a^{*VB})^{T_V},$ (A35a)

$$M_a^{*VB} \geq 0, \quad M_a^{*VB} \in \Sigma \quad \forall a, \quad (\text{A35b})$$

$$\sum_a M_a^{*VB} = \mathbb{1} \otimes \frac{\rho^B + r \frac{\mathbb{1}^B}{d}}{1+r}. \quad (\text{A35c})$$

For each value of a , (A35a) can be transformed in the following way:

$$\frac{U_a^V \rho^{VB} U_a^{\dagger V} + r d^2 p(a) \frac{\mathbb{1}^{AB}}{d^2}}{1+r d^2 p(a)} = \frac{1}{\text{tr} M_a^{*VB}} (M_a^{*VB})^{T_V}.$$

Thus, the smallest r satisfying (A35a) and (A35b) for each a separately is equal to $\epsilon_r(U_a^V \rho^{VB} U_a^{\dagger V})/d^2 p(a) = \epsilon_r(\rho^{VB})/d^2 p(a)$. Since there are d^2 different outcomes a , the smallest r which can simultaneously satisfy (A35a) for all values of a is equal to ϵ_r . By summing (A34) over a , we see that the last constraint from (A14) is satisfied, which finally implies

$$\tau_r^*(\sigma_{a|\omega_x}) = \epsilon_r(\rho^{AB}).$$

8. Teleportation using a partial Bell state measurement

Teleportation experiments where Alice performs a partial Bell state measurement using POVM $M_0^{VA} = \Phi^{+VA}$, $M_1^{VA} = \sum_{i=1}^{d^2-1} U_i^V \Phi^{+VA} U_i^{\dagger V}$ and has access to a tomographically complete set of input states are also of particular interest.

Let us denote the random teleportation robustness of a teleportation assemblage obtained by performing such a measurement as $\tau_r'(\cdot)$. Taking into account that the set of input states is tomographically complete, $\tau_r'(\cdot)$ can be expressed as the solution to the following optimization problem:

$$\tau_r'(\{\sigma_{a|\omega_x}\}) = \min_{r, \{M_a^*\}, \{p(a)\}} r$$

s.t. $\frac{\rho^{VB} + r p(0) \mathbb{1}^{VB}}{1+r} = d(M_0^{*VB})^{T_V},$ (A36a)

$$\frac{\sum_{i=1}^{d^2-1} U_i^V \rho^{VB} U_i^{\dagger V} + r p(1) \mathbb{1}^{VB}}{1+r} = d(M_1^{*VB})^{T_V}, \quad (\text{A36b})$$

$$M_a^{*VB} \geq 0, \quad M_a^{*VB} \in \mathcal{S} \quad \forall a, \quad (\text{A36c})$$

$$\sum_a M_a^{*VB} = \mathbb{1} \otimes \frac{\rho^B + r \frac{\mathbb{1}^B}{d}}{1+r}. \quad (\text{A36d})$$

Note that the constraint (A36b), based on (A36a) and satisfying (A36d) can be reduced to

$$M_1^{*VB} = \sum_{i=1}^{d^2-1} M_0^{*VB} + \frac{[p(1) - p(0)(d^2 - 1)] \mathbb{1}^{VB}}{d(1+r)},$$

which is separable whenever M_0^{*VB} is separable.¹ This means that every r satisfying (A36a) also satisfies (A36b), which in turn implies that $\tau_r'(\{\sigma_{a|\omega_x}\})$ is equal to the smallest r satisfying (A36a) and (A36c). Following the equivalence of (A36) and (A32), the smallest such r is equal to $\epsilon_r(\rho^{AB})/[d^2 p(0)]$. The optimal mixing assemblage is the trivial one $\{\mathbb{1}^B/d, 0\}$ leading to

$$\tau_r'(\{\sigma_{a|\omega_x}\}) = \frac{\epsilon_r(\rho^{AB})}{d^2}. \quad (\text{A37})$$

Note that in [9] a different bound was obtained, namely that $\tau_r'(\{\sigma_{a|\omega_x}\}) = \frac{2\epsilon_r(\rho^{AB})}{d^2}$. This is due to the different definition used for the random teleportation robustness. There, as noted earlier, the mixing teleportation assemblage had the form $\frac{\mathbb{1}}{|o|d}$, which automatically fixed $p(0)$ to be equal to $1/2$.

APPENDIX B: TELEPORTATION WEIGHT AND BEST SEPARABLE APPROXIMATION

In this Appendix we first show that the teleportation weight of the teleportation assemblage $\{\sigma_{a|\omega_x}^B\}_{a,x}$ puts a lower bound on the best separable approximation of the shared state ρ^{AB} . First, let us observe that for the state ρ^{AB} and its best separable approximation $\epsilon_{\text{BSA}}(\rho^{AB})$ there exist a corresponding quantum state $\tilde{\rho}^{AB}$ and separable state $\tilde{\rho}^{AB}$ such that

$$\rho^{AB} = \epsilon_{\text{BSA}}(\rho^{AB}) \tilde{\rho}^{AB} + [1 - \epsilon_{\text{BSA}}(\rho^{AB})] \tilde{\rho}^{AB}.$$

By tensoring ρ^{AB} with the state ω_x^V and applying a joint measurement M_a^{VA} , this implies

$$\begin{aligned} \text{tr}_{VA} [(M_a^{VA} \otimes \mathbb{1}^B)(\omega_x^V \otimes \rho^{AB})] \\ = \epsilon_{\text{BSA}}(\rho^{AB}) \text{tr}_{VA} [(M_a^{VA} \otimes \mathbb{1}^B)(\omega_x^V \otimes \tilde{\rho}^{AB})] \\ + [1 - \epsilon_{\text{BSA}}(\rho^{AB})] \text{tr}_{VA} [(M_a^{VA} \otimes \mathbb{1}^B)(\omega_x^V \otimes \tilde{\rho}^{AB})], \end{aligned}$$

i.e.,

$$\begin{aligned} \sigma_{a|\omega_x}^B = \text{tr}_V (\{ \epsilon_{\text{BSA}}(\rho^{AB}) \tilde{M}_a^{*VB} \\ + [1 - \epsilon_{\text{BSA}}(\rho^{AB})] \tilde{M}_a^{*VB} \} \omega_x^V \otimes \mathbb{1}^B), \end{aligned} \quad (\text{B1})$$

where

$$\begin{aligned} \tilde{M}_a^{*VB} = \text{tr}_A [(M_a^{VA} \otimes \mathbb{1}^B)(\mathbb{1}^V \otimes \tilde{\rho}^{AB})], \\ \tilde{M}_a^{*VB} = \text{tr}_A [(M_a^{VA} \otimes \mathbb{1}^B)(\mathbb{1}^V \otimes \tilde{\rho}^{AB})], \end{aligned} \quad (\text{B2})$$

for all a and x , (B1) is equivalent to the first constraint from the optimization problem (18). Moreover, the operators \tilde{M}_a^{*VB} and \tilde{M}_a^{*VB} defined in (B2) satisfy all the other constraints from (18). Thus, the teleportation weight of the teleportation assemblage

¹This is expected since constraint (A36b) corresponds to the member of teleportation assemblage which is obtained by using separable measurement M_1^{VA} .

$\{\sigma_{a|\omega_x}\}_{a,x}$ can only be smaller than the best separable approximation of the shared state ρ^{AB} , i.e.,

$$\text{TW}(\{\sigma_{a|\omega_x}^B\}_{a,x}) \leq \epsilon_{\text{BSA}}(\rho^{AB}).$$

Now we show that if the teleportation assemblage $\{\sigma_{a|\omega_x}^B\}_{a,x}$ is obtained by applying a Bell state measurement on Alice's share of the state ρ^{AB} and states from a tomographically complete set $\{\omega_x\}_x$, its teleportation weight is equal to the best separable approximation of the state ρ^{AB} . In such a scenario the first constraint from the optimization problem (18) can be rewritten in the following way:

$$\begin{aligned} \sigma_{a|\omega_x}^B &= \text{tr}_{\text{VA}} [(U_a^A \Phi^{+\text{VA}} U_a^{\dagger A} \otimes \mathbb{1}^B)(\omega_x^V \otimes \rho^{AB})] \\ &= p \text{tr}_{\text{VA}} \{(\Phi^{+\text{VA}} \otimes \mathbb{1}^B)[\omega_x^V \otimes d^2 \tilde{\rho}(a) \tilde{\rho}_a^{\text{AB}}]\} \\ &\quad + (1-p) \text{tr}_{\text{VA}} \{(\Phi^{+\text{VA}} \otimes \mathbb{1}^B)[\omega_x^V \otimes d^2 \bar{\rho}(a) \bar{\rho}_a^{\text{AB}}]\}. \end{aligned}$$

The constraints on \tilde{M}_a and \bar{M}_a impose that the states $\tilde{\rho}_a$ could be any quantum states, while the states $\bar{\rho}_a$ are separable. Furthermore, using identity (11) the last equation reduces to

$$\begin{aligned} \frac{1}{d} \text{tr}_V [(U_a^V \rho^{VB} U_a^{\dagger V})^{T_V} (\omega_x^V \otimes \mathbb{1}^B)] \\ = \frac{1}{d} \text{tr}_V \{[p d^2 \tilde{\rho}(a) \tilde{\rho}_a^{\text{VB}} + (1-p) d^2 \bar{\rho}(a) \bar{\rho}_a^{\text{VB}}]^{T_V} (\omega_x^V \otimes \mathbb{1}^B)\}. \end{aligned} \quad (\text{B3})$$

For tomographically complete set of inputs $\{\omega_x\}_x$ this equation is satisfied if and only if

$$U_a^V \rho^{VB} U_a^{\dagger V} = d^2 [p \tilde{\rho}(a) \tilde{\rho}_a^{\text{VB}} + (1-p) \bar{\rho}(a) \bar{\rho}_a^{\text{VB}}]. \quad (\text{B4})$$

Together with the constraints on the states $\tilde{\rho}_a$ and $\bar{\rho}_a$, we see that the optimization problem of finding the teleportation weight of a teleportation assemblage obtained by using a Bell state measurement and tomographically complete set of inputs can be reduced to

$$\begin{aligned} \text{TW}(\{\sigma_{a|\omega_x}^B\}) &= \min_{p, \tilde{\rho}(a), \tilde{\rho}_a, \bar{\rho}(a), \bar{\rho}_a} p \\ \text{s.t. } U_a^V \rho^{VB} U_a^{\dagger V} &= d^2 [p \tilde{\rho}(a) \tilde{\rho}_a^{\text{VB}} + (1-p) \bar{\rho}(a) \bar{\rho}_a^{\text{VB}}], \\ \sum_a \tilde{\rho}(a) &= 1, \quad \sum_a \bar{\rho}_a = 1, \\ \tilde{\rho}_a &\in \mathcal{S}. \end{aligned} \quad (\text{B5})$$

For every a the minimal p satisfying (B4) is similar to the constraint appearing in the expression for the best separable approximation of the state $U_a^V \rho^{VB} U_a^{\dagger V}$. The difference is that in (B4) the states $d^2 \tilde{\rho}(a) \tilde{\rho}_a^{\text{VB}}$ and $d^2 \bar{\rho}(a) \bar{\rho}_a^{\text{VB}}$ need not be normalized. If for some $a = a'$, $\text{tr}[d^2 \tilde{\rho}(a') \tilde{\rho}_{a'}^{\text{VB}}]$ is bigger than 1, the minimal p satisfying (B4) would be smaller than the best separable approximation of $U_{a'}^V \rho^{VB} U_{a'}^{\dagger V}$. But because $\sum_a d^2 \bar{\rho}(a) \bar{\rho}_a^{\text{VB}} = d^2$ and there are d^2 different values of a , it means that for some other $a = a''$ we will have $\text{tr}[d^2 \tilde{\rho}(a'') \tilde{\rho}_{a''}^{\text{VB}}] < 1$, which would make the smallest p satisfying (B4) strictly larger than the best separable approximation of $U_{a''}^V \rho^{VB} U_{a''}^{\dagger V}$. Since all the states related by local unitary transformations have the same best separable approximation, the optimal p satisfying all d^2 different constraints contained in (B4) must be equal to $\epsilon_{\text{BSA}}(\rho^{VB})$.

-
- [1] C. H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres, and W. K. Wootters, Teleporting an Unknown Quantum State via Dual Classical and Einstein-Podolsky-Rosen Channels, *Phys. Rev. Lett.* **70**, 1895 (1993).
- [2] N. Gisin, G. Ribordy, W. Titel, and H. Zbinden, Quantum cryptography, *Rev. Mod. Phys.* **74**, 145 (2002).
- [3] H.-J. Briegel, W. Dür, J. I. Cirac, and P. Zoller, Quantum Repeaters: The Role of Imperfect Local Operations in Quantum Communication, *Phys. Rev. Lett.* **81**, 5932 (1998).
- [4] D. Gottesman and I. L. Chuang, Demonstrating the viability of universal quantum computation using teleportation and single-qubit operations, *Nature (London)* **402**, 390 (1999).
- [5] R. Raussendorf and H.-J. Briegel, A One-way Quantum Computer, *Phys. Rev. Lett.* **86**, 5188 (2001).
- [6] S. Popescu, Bell's Inequalities Versus Teleportation: What is Nonlocality? *Phys. Rev. Lett.* **72**, 797 (1994).
- [7] M. Horodecki, P. Horodecki, and R. Horodecki, General teleportation channel, singlet fraction, and quasidistillation, *Phys. Rev. A* **60**, 1888 (1999).
- [8] M. Horodecki, P. Horodecki, and R. Horodecki, Mixed-State Entanglement and Distillation: Is there a "Bound" Entanglement in Nature? *Phys. Rev. Lett.* **80**, 5239 (1998).
- [9] D. Cavalcanti, P. Skrzypczyk, and I. Šupić, All Entangled States Can Demonstrate Non-Classical Teleportation, *Phys. Rev. Lett.* **119**, 110501 (2017).
- [10] G. Carvacho, F. Andreoli, L. Santodonato, M. Bentivegna, V. D'Ambrosio, P. Skrzypczyk, I. Šupić, D. Cavalcanti, and F. Sciarrino, Experimental Study of Nonclassical Teleportation Beyond Average Fidelity, *Phys. Rev. Lett.* **121**, 140501 (2018).
- [11] M. J. Hoban and A. B. Sainz, A channel-based framework for steering, non-locality and beyond, *New J. Phys.* **20**, 053048 (2018).
- [12] A. C. Doherty, P. A. Parrilo, and F. M. Spedalieri, Distinguishing Separable and Entangled States, *Phys. Rev. Lett.* **88**, 187904 (2002); Complete family of separability criteria, *Phys. Rev. A* **69**, 022308 (2004).
- [13] S. Boyd and L. Vandenberghe, *Convex Optimization* (Cambridge University Press, Cambridge, 2004).
- [14] G. Vidal and R. Werner, A computable measure of entanglement, *Phys. Rev. A* **65**, 032314 (2002).
- [15] G. Vidal and R. Tarrach, Robustness of entanglement, *Phys. Rev. A* **59**, 141 (1999).
- [16] M. Steiner, Generalized robustness of entanglement, *Phys. Rev. A* **67**, 054305 (2003).

- [17] I.Šupić, P. Skrzypczyk, and D. Cavalcanti, Measurement-device-independent entanglement and randomness estimation in quantum networks, *Phys. Rev. A* **95**, 042340 (2017).
- [18] D. Rosset, A. Martin, E. Verbanis, C. C. W. Lim, and R. Thew, Practical measurement-device-independent entanglement quantification, *Phys. Rev. A* **98**, 052332 (2018).
- [19] M. Lewenstein and A. Sanpera, Separability and Entanglement of Composite Quantum Systems, *Phys. Rev. Lett.* **80**, 2261 (1998).
- [20] A. C. Elitzur, S. Popescu, and D. Rohrlich, Quantum nonlocality for each pair in an ensemble, *Phys. Lett. A* **162**, 25 (1992).
- [21] P. Skrzypczyk, M. Navascués, and D. Cavalcanti, Quantifying Einstein-Podolsky-Rosen Steering, *Phys. Rev. Lett.* **112**, 180404 (2014).
- [22] Notebook available at <https://git.io/fN3Qh>
- [23] P. Horodecki, Separability criterion and inseparable mixed states with positive partial transposition, *Phys. Lett. A* **232**, 333 (1997).
- [24] C. H. Bennet, D. P. DiVincenzo, T. Mor, P. W. Shor, J. A. Smolin, and B. Terhal, Unextendible Product Bases and Bound Entanglement, *Phys. Rev. Lett.* **82**, 5385 (1999).