



Wild, S., Krützen, M., Rankin, R., Hoppitt, W., Gerber, L., & Allen, S. (2019). Long-term decline in survival and reproduction of dolphins following a marine heatwave. *Current Biology*, 29(7), R239-R240. <https://doi.org/10.1016/j.cub.2019.02.047>

Peer reviewed version

License (if available):
CC BY-NC-ND

Link to published version (if available):
[10.1016/j.cub.2019.02.047](https://doi.org/10.1016/j.cub.2019.02.047)

[Link to publication record on the Bristol Research Portal](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Cell Press at <https://www.sciencedirect.com/science/article/pii/S0960982219302179> . Please refer to any applicable terms of use of the publisher.

University of Bristol – Bristol Research Portal

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available: <http://www.bristol.ac.uk/red/research-policy/pure/user-guides/brp-terms/>

Long-term decline in reproduction and survival of dolphins following a marine heatwave

Sonja Wild*^{1,2}, Michael Krützen², Robert W. Rankin^{3,4}, William J.E. Hoppitt¹, Livia Gerber², Simon J. Allen^{2,5,6}

¹School of Biology, University of Leeds, Leeds, United Kingdom

²Evolutionary Genetics Group, Department of Anthropology, University of Zurich, Zurich, Switzerland

³Department of Biology, Georgetown University, Washington DC, USA

⁴National Socio-Environmental Synthesis Center (SESYNC), University of Maryland, Annapolis MD, USA

⁵School of Biological Sciences and Oceans Institute, University of Western Australia, Crawley, Western Australia, Australia

⁶School of Biological Sciences, University of Bristol, Bristol, United Kingdom

*corresponding author: bssw@leeds.ac.uk

One of many challenges in the conservation of biodiversity is the recent trend and predicted increase in the frequency and duration of extreme climatic events [1]. The Shark Bay World Heritage Area, Western Australia, endured an unprecedented marine heatwave in 2011. Catastrophic losses of habitat-forming seagrass meadows throughout the bay followed [2], along with mass mortalities of invertebrate and fish communities [3]. Our long-term demographic data on Shark Bay's iconic population of resident Indo-Pacific bottlenose dolphins (*Tursiops aduncus*) revealed a significant decline in female reproductive rates subsequent to the heatwave. Moreover, capture-recapture analyses indicated 5.9% and 12.2% post-heatwave declines in the survival of dolphins that use tools to forage and those that do not, respectively. This implies that the tool using dolphins may have been somewhat buffered against the cascading effects of habitat loss following the heatwave by having access to a less severely affected foraging niche [4]. Overall, however, lower survival levels have persisted post-heatwave, suggesting that habitat loss following extreme weather events may have prolonged negative impacts on even behaviourally flexible, higher-trophic level predators.

In the Austral summer/autumn of 2011, the coastal waters of Western Australia experienced an unprecedented marine heatwave, with water temperatures rising to 2-4° Celsius above long-term averages for more than two months [5]. In the shallow, subtropical embayment of Shark Bay, an estimated 36% of seagrass meadows were damaged [2]. Wide-ranging effects associated with the heatwave were documented across lower trophic levels, while those on long-lived or large vertebrates remained more elusive. We investigated the vital rates, *i.e.* survival and reproduction, of dolphins in the western gulf of Shark Bay, using long-term demographic and behavioural data collected between 2007 and 2017 (Fig. 1a; SI).

We assessed changes in apparent survival of dolphins over time using capture-recapture analyses (Fig. 1a; SI; Data S1). Using Akaike weights as approximate model probabilities, there was evidence of a considerable decrease in survival after the heatwave. We suggest that this was likely linked to

the well-documented and sizeable losses of prey during the heatwave [3]. We also found that, on average, survival remained lower in the seven years after the heatwave compared to before. Again, having most support from Akaike weights, we suggest that this result is best interpreted as evidence that dolphin mortality was impacted over a protracted period, rather than a single peak following the temperature anomaly, which had much lower support by AIC. As such, the catastrophic reduction in seagrass coverage [2], which shows little sign of recovery [6], appears to be responsible for preventing fish stock recovery, since well-established seagrass meadows represent important breeding grounds and refuge habitat for numerous species [7].

We also tested for potential differences in survival between dolphins occupying different habitats, and those that use marine sponges as foraging tools ('spongers') versus those that do not ('non-spongers') [4]. Interestingly, while there was an overall decline in dolphin survival post-heatwave, survival of spongers was not as adversely impacted as that of non-spongers (5.9% versus 12.2% decline in survival from pre- to post-heatwave, respectively; Fig. 1b). Sponge use is restricted to deeper channel habitats with no seagrass cover. While both spongers and non-spongers use these channels for foraging, sponge use allows access to a foraging niche which dolphins without the tool using know-how cannot access [4], implying that the spongers' foraging niche may have been less severely impacted by the heatwave and subsequent seagrass losses (Fig. 1b).

A significantly greater number of calves per female per year was detected pre-heatwave compared to post-heatwave (Poisson GLM; $z = -2.579$; effect size = 31.6%; 95% C.I. = [7.5%, 55.7%]; $p < 0.010$; Fig. 1c), while no significant differences were found when comparing reproductive success for females from different habitat types (Poisson GLM; $z = 1.468$; effect size = 18.17%; 95% C.I. = [-6.0%; 42.6%]; $p = 0.142$), or using different foraging techniques (spongers, non-spongers; Poisson GLM; $z = -0.673$; effect size = -9.6%; 95% C.I. = [-38.5%; 17.8%]; $p = 0.501$) (Data S1). This suggests that the spongers' access to a different foraging niche may have led to less severe effects on survival, but not

reproduction. Reduced reproductive success as a consequence of significant ecological changes is not surprising, since a decline in food availability is expected to affect the most vulnerable members of a population, *i.e.*, young individuals and those with high nutritional demands, such as pregnant or lactating females [8].

[Fig. 1 appr. here]

There are several plausible explanations for this reduced reproductive success: First, lower prey availability is likely to have forced female dolphins to spend more time foraging, leading to reduced vigilance and, ultimately, greater shark predation on calves. Second, reduced food availability can lead to either increased rates of abortion during pregnancy or increased neonate mortality, when both the mother's and the offspring's nutritional needs cannot be sustained [8]. Both abortions and increased mortality of young calves would effectively appear as lower reproductive success in the demographic data. Third, suppressed ovulation or delayed sexual maturity may have occurred when females did not reach a certain threshold of body weight [9]. Indeed, along with juvenile survival, fertility is the vital rate most sensitive to changes in resource availability in marine mammals [10].

Western Australia's 2011 marine heatwave negatively impacted habitat-forming and lower-trophic level organisms [2,3,5], and our results suggest cascading effects through to a behaviourally flexible, top-order predator. Long-lived taxa, such as some birds, great apes and cetaceans, are likely to experience changing ecological conditions throughout their lifetimes and, hence, may display behavioural plasticity in adapting to such changes. However, our findings suggest that the ecological consequences of extreme weather events may be too sudden or disruptive for even highly adaptable, cognitively advanced animals to respond, leading to long-term negative impacts on their population viability. Such impacts may persist if ecosystems fail to recover. Furthermore, our study adds to prior research illustrating that extreme events can drive biodiversity patterns and cause

shifts in community structure toward more depauperate states [5]. These findings raise concerns over the long-term viability of the dolphin population, given that marine heatwaves are occurring with greater frequency and duration in association with anthropogenic climate change [1].

Acknowledgements

This research was funded by grants from the Swiss Science Foundation (31003A_149956 to MK), National Geographic Society, Sea World Research and Rescue Foundation Inc., W.V. Scott Foundation and the A.H. Schultz Stiftung. We thank Mitsui, Shark Bay Resources Pty. Ltd., and the Useless Loop community for in-kind and logistical support for our research teams. We also thank the numerous team leaders and field assistants that have contributed to data collection for the Dolphin Innovation Project between 2007 and 2017, without whom this research would not have been possible. This research was carried out under scientific investigation permits from the Department of Biodiversity, Conservation and Attractions, as well as animal ethics approvals from the University of Zurich, University of Western Australia, Murdoch University and the University of New South Wales.

Author contributions

Conceptualization, SW, MK, SJA. Formal Analysis, SW, RR, WH. Investigation, SW, SJA, LG. Writing – Original Draft, SW, SJA, MK. Writing – Review & Editing, SW, SJA, MK, RR, WH, LG. Funding acquisition, MK.

Declaration of interest

The authors declare no competing interests.

References

1. Oliver, E.C.J., Donat, M.G., Burrows, M.T., Moore, P.J., Smale, D.A., Alexander, L. V, Benthuisen, J.A., Feng, M., Sen Gupta, A., Hobday, A.J., *et al.* (2018). Longer and more frequent marine heatwaves over the past century. *Nat. Commun.* 9, 1–12. Available at: <http://dx.doi.org/10.1038/s41467-018-03732-9>.
2. Arias-Ortiz, A., Serrano, O., Masqué, P., Lavery, P.S., Mueller, U., Kendrick, G.A., Rozaimi, M., Esteban, A., Fourqurean, J.W., Marbà, N., *et al.* (2018). A marine heatwave drives massive losses from the world's largest seagrass carbon stocks. *Nat. Clim. Chang.* 2018 8, 338–344. Available at: <https://www.nature.com/articles/s41558-018-0096-y>.
3. Pearce, A., Lenanton, R., Jackson, G., Moore, J., Feng, M., and Gaughan, D. (2011). The

- “marine heat wave” off Western Australia during the summer of 2010/11. Fish. Res. Rep. No. 222, 40pp. Available at: http://fish.gov.au/reports/Documents/Pearce_et_al_2011.pdf [Accessed April 22, 2014].
4. Krützen, M., Kreicker, S., Macleod, C.D., Learmonth, J., Kopps, A.M., Walsham, P., and Allen, S.J. (2014). Cultural transmission of tool use by Indo-Pacific bottlenose dolphins (*Tursiops* sp.) provides access to a novel foraging niche. *Proc. R. Soc. B* 281.
 5. Wernberg, T., Smale, D.A., Tuya, F., Thomsen, M.S., Langlois, T.J., de Bettignies, T., Bennett, S., and Rousseaux, C.S. (2012). An extreme climatic event alters marine ecosystem structure in a global biodiversity hotspot. *Nat. Clim. Chang.* 3, 78–82. Available at: <http://dx.doi.org/10.1038/nclimate1627>.
 6. Nowicki, R.J., Thomson, J.A., Burkholder, D.A., Fourqurean, J.W., and Heithaus, M.R. (2017). Predicting seagrass recovery times and their implications following an extreme climate event. *Mar. Ecol. Prog. Ser.* 567, 79–93.
 7. Heithaus, M.R. (2004). Fish Communities of Seagrass Meadows and Associated Habitats in Shark Bay, Western Australia. *Bull. Mar. Sci.* 75, 79–99.
 8. Trites, A.W., and Donnelly, C.P. (2003). The decline of Steller sea lions *Eumetopias jubatus* in Alaska: A review of the nutritional stress hypothesis. *Mamm. Rev.* 33, 3–28.
 9. Boyd, I.L., Lockyer, C., and Marsh, H.D. (1999). Reproduction in marine mammals. In *Biology of marine mammals*, J. E. Reynolds and S. A. Rommel, eds. (Washington, D.C.: Smithsonian Institution Press), pp. 218–286.
 10. King, S.L., Schick, R.S., Donovan, C., Booth, C.G., Burgman, M., Thomas, L., and Harwood, J. (2015). An interim framework for assessing the population consequences of disturbance. *Methods Ecol. Evol.* 6, 1150–1158.

Figure 1: Dolphin vital rates before and after the 2011 marine heatwave in western Shark Bay

a) The study area in the western gulf of Shark Bay, Western Australia, encompassing approximately 1,500 km². Over 5,000 dolphin group encounters have been documented between 2007 and 2017 (all points). To account for unequal survey effort in each field season, the study area was overlaid with a grid of 2 x 2 km cells. Only encounters within grid cells covered in all seasons (‘core’ study area) were considered for capture-recapture analyses (dark red points). **b)** While both spongers and non-spongers experienced a decline in survival from pre- to post-heatwave, spongers were less affected than non-spongers (5.9% decline versus 12.2% decline, respectively). **c)** The number of calves detected (controlled for number of known mothers observed each field season) was significantly higher pre- compared to post-heatwave, suggesting that female dolphin reproductive success was also negatively impacted by habitat degradation and prey species loss following the heatwave. (Note that the pre-heatwave data set encompassed only four years, while that post-heatwave was seven).

Data S1: Input data and R codes for assessment of dolphin survival and reproduction