



Weaving, H., Sands, B., & Wall, R. (2019). Reproductive sublethal effects of macrocyclic lactones and synthetic pyrethroids on dung beetle *Onthophagus similis*. *Bulletin of Entomological Research*. Advance online publication.  
<https://doi.org/10.1017/S0007485319000567>

Peer reviewed version

Link to published version (if available):  
[10.1017/S0007485319000567](https://doi.org/10.1017/S0007485319000567)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Cambridge University Press at <https://www.cambridge.org/core/journals/bulletin-of-entomological-research/article/reproductive-sublethal-effects-of-macrocyclic-lactones-and-synthetic-pyrethroids-on-the-dung-beetle-onthophagus-similis/86759921FD88C67D80EE9B3E006F4B1D>. Please refer to any applicable terms of use of the publisher.

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

# Reproductive sublethal effects of macrocyclic lactones and synthetic pyrethroids on dung beetle *Onthophagus similis*

H. Weaving, B. Sands and R. Wall

Veterinary Parasitology and Ecology Group, School of Biological Sciences, 24 Tyndall Avenue,  
University of Bristol, Bristol, BS8 1TQ, UK

## Abstract

Dung-colonising beetles provide a range of ecosystem services in farmland pasture systems. However, such beetles are declining in Northern temperate regions. This may, in part, be due to the widespread use of macrocyclic lactones (MLs) and synthetic pyrethroids (SPs) in livestock farming. These chemicals are used to control pests and parasites of cattle; the residues of which are excreted in dung at concentrations toxic to insects. While the lethal effects of such residues are well known, sublethal effects are less understood. Any effects, however, may have important consequences for beetle populations, particularly if they affect reproduction. To investigate, the impact of ML and SP exposure on the reproductive output of *Onthophagus similis* (Scriba), a Northern temperate dung beetle species, was examined. In laboratory trials, field-collected adult *O. similis* exposed to the ML ivermectin at 1 ppm (wet weight) over a period of 3 weeks had smaller oocytes ( $p=0.016$ ), smaller fat bodies and reduced motility compared to the control. In a farm-level investigation, cattle dung-baited pitfall trapping was undertaken on 23 beef cattle farms in SW England, which either used MLs ( $n=9$ ), SPs ( $n=7$ ) or neither chemical ( $n=7$ ). On farms that used no MLs or SPs, 24.2% of females caught were gravid. However, on farms that used MLs no gravid females were caught, and only 1% of the beetles caught on farms using SPs were gravid ( $p<0.001$ ). The association between ML and SP use and impaired reproductive output suggests that the use of such chemicals is likely to be ecologically damaging.

**Key words:** dung beetle, ivermectin, macrocyclic lactone, synthetic pyrethroid, reproductive output, pesticide, parasiticide

\*Author for correspondence  
Phone: 01173941205  
E-mail: hw14528@bristol.ac.uk

## Introduction

Insects provide essential ecosystem services as pollinators, detritivores and regulators of pest abundance (Losey & Vaughan, 2006). Dung-colonising beetles are no exception; they aerate dung and move it below ground, thereby reducing dung-dwelling parasites (Sands *et al.*, 2017), improving soil structure and quality, facilitating nitrogen and carbon recycling and preventing dung from building up on pasture (Bornemissza, 1970; Fincher, 1981; Yamada *et al.*, 2007; Nichols *et al.*, 2008). These services have estimated to contribute at least £400 million to the U.K. agricultural industry annually, calculated to the current day equivalent in 2019 (Beynon *et al.*, 2015; Eliassen, 2019). However, in Northern temperate pastureland systems many dung beetle species are in widespread decline, some having become locally extinct (Bistrom *et al.*, 1991; Lobo *et al.*, 2001; Roslin *et al.*, 2014; Natural England, 2016). Some of this decline can be attributed to an intensification of farming and habitat fragmentation (Hutton & Giller, 2003; Filgueiras *et al.*, 2011). However, chemicals used to treat cattle for pests and parasites, the residues of which are excreted in dung in toxic concentrations, also may have contributed to this decline via nontarget lethal and sublethal effects (Sommer *et al.*, 1992; Floate *et al.*, 2005; Jacobs & Scholtz, 2015).

The macrocyclic lactones (MLs) (including ivermectin, eprinomectin, doramectin and moxidectin) are used extensively in livestock farming (Stafford & Coles, 1999). They are broad spectrum endectocides, active against warbles, mites, lice and gastrointestinal worms (Sutherland & Campbell, 1990; Floate *et al.*, 2005). Administration of MLs to cattle can be via subcutaneous or intramuscular injection, topically (pour-on), or orally. Once metabolised in the body, up to 98% of the active ingredient is excreted in the faeces, although the amount and rate of elimination varies with route of administration (Campbell *et al.*, 1989; Sommer *et al.*, 1992; McKellar & Gokbulut, 2012). A second major class of chemicals used in livestock husbandry are the synthetic pyrethroids (SPs), for the control of nuisance flies and ectoparasites (Pickett, 2004). In cattle, the route of elimination for SPs is mainly via the faeces, and the amount and

rate of excretion also varies with the compound applied and route of administration (Venant *et al.*, 1990; Floate *et al.*, 2005).

At environmentally relevant concentrations, residues of both MLs and SPs in dung have been shown to have lethal effects on dung colonising insects (Wall & Strong, 1987; Kruger & Scholz, 1997; Vale *et al.*, 2004; Vale *et al.*, 2015) of all life-cycle stages (Ridsdill-Smith, 1988; Sommer *et al.*, 1992; Cruz Rosales *et al.*, 2012; Mann *et al.*, 2015; Perez-Cogollo *et al.*, 2015; Sands *et al.*, 2018). However, sublethal effects resulting from exposure to ML and SP residues have been less thoroughly investigated (Wardhaugh & Rodriguez-Menendez, 1988; Houlding *et al.*, 1991; Kruger & Scholtz, 1997; Dadour *et al.*, 2000; Bang *et al.*, 2007; Cruz Rosales *et al.*, 2012; Martinez *et al.*, 2017). In consequence, the ecosystem services dung beetles provide may be more severely compromised than indicated simply by an understanding of the lethal effects (Manning *et al.*, 2017).

The aim of the work described here was to investigate the effect of MLs and SPs used in cattle farming on the reproductive physiology and fecundity of *Onthophagus similis* (Scriba), a temperate dung beetle species. Specifically, the study asked two questions. First, does *in vitro* exposure to sublethal doses of the ML ivermectin in dung affect the reproductive output and physiology of *O. similis*? And secondly, on farms in South West England, does the long-term use of MLs and SPs in beef herds have an impact on the reproductive development of *O. similis* in the field?

## Methods

### *Laboratory investigation*

Twenty dung-baited pitfall traps were used to catch beetles at a registered organic farm in South West England from 9<sup>th</sup> – 13<sup>th</sup> May 2018. Pitfall traps comprised a pot (18 cm depth x 16 cm diameter), buried to rim level in soil. The soil was packed tightly around the pot rim to

ensure the capture of both flying and walking dung beetles. Traps were positioned in a field adjacent to grazing cattle between 09:00-11:00 am, and placed approximately 10 meters apart. Fresh cattle dung was collected as bait and placed on a chicken wire mesh on top of the buried pot, using a 20 cm diameter pat former, which held 1 kg dung. The bait covered the entire opening of the pot to prevent the exit of beetles. Traps were protected from rain by 15 cm diameter aluminium dishes, supported using 30 cm bamboo canes, dug in to the soil. The cattle from which dung was collected had not been treated with any MLs or SPs for approximately 6 months. Bait was replaced with fresh dung every other day, and on days where dung was not replaced, the crust was peeled off and discarded.

Dung for use in the experiment was collected from the same farm and was homogenised with an industrial paddle mixer (Silverline 850W; Lufton Trading Estate, Yeovil) for 15 min and stored at 4 °C before use. Dry weight analysis of a subsample, using sequential drying in an oven followed by reweighing, showed that the dung was 11.5% dry mass. Beetles were collected from traps daily and placed into terraria, made from plastic containers (8 cm, 11.5 cm, 17 cm) with holes in the lid to allow air exchange. The terraria contained 400 g of clean sand and 250 g of cattle dung, and were stored at 4 °C. *Onthophagus similis* was identified using the morphological keys of Jessop (1986) and Skidmore (1991).

Before the experiment commenced, beetles were starved for 24 h. Experimental terraria, as described above, were maintained at 20 °C, on a 12:12 h light:dark cycle in an illuminated cooled incubator (Sanyo Electric, Osaka City, 540-6226, Japan). Each terrarium was supplied with 250 g of dung spiked with ivermectin (Ivomec Super®, Boehringer Ingelheim Ltd., Berkshire) at concentrations of 0.01 ppm, 0.1 ppm and 1 ppm in wet dung. This product also contains 10% w/v clorsulon; a narrow-spectrum flukicide which is inactive against insects. Dilutions were made by mixing the required quantity of 1% ivermectin (Ivomec Super®) in 10 ml of ethanol per kg of dung. The solution was mixed into the dung for 10 minutes using a spoon

and glass rod to ensure it was thoroughly homogenised throughout the dung. The dung for the control group was spiked with ethanol only. To allow evaporation of the solvent, mixtures were left for 1 h before use.

Twelve terraria were each populated with 13 *O. similis* of approximately equal sex ratio. Dominant males (those with a horn) were also divided evenly between the groups. Once divided, terraria were randomly allocated to treatments (0.01 ppm, 0.1 ppm and 1 ppm ivermectin and ethanol control), with three repeats of each. The terraria were maintained for 3 weeks in total, with dung replaced weekly. After three weeks, all beetles were killed and stored in ethanol.

Dissections were carried out on all female beetles in Ringer's solution (EMD Millipore Corporation, Billerica MA 01821, USA) under a GX dissecting microscope (GT Vision Ltd, Standsfield, CO10 8LY). Females were designated a reproductive state: gravid or non-gravid. Individuals were defined as gravid if oocytes were differentiated, and non-gravid if oocytes were undifferentiated so that the coiled germarium was one mass. Oocyte counts and measurements were only taken for gravid individuals. The basal oocyte was measured; this was the most mature oocyte at the base of the coiled germarium. A count was also taken of the total number of maturing oocytes. Additionally, the left hind leg was removed for measurement of the tibia. This measurement was used as an indication of beetle size; it was assumed that tibia length and body size varied isometrically (e.g. Lease & Wolf, 2011; Kanpp & Knappova, 2013). The hind tibia of all females was measured, regardless of reproductive state. All egg and leg measurements were made using a Leica MZ12 dissecting microscope and Leica DFC295 digital camera and the Leica Application Suite V3.8, measurement tool (Leica Microsystems Ltd, Milton Keynes, MK14 6FG).

### *Field investigation*

Twenty-three beef cattle farms in the South West of England were selected. Some farms used macrocyclic lactones predominantly (n = 9), some used synthetic pyrethroids (n = 7), and the others used no MLs or SPs at all (n = 7). All farms had been under the same regime for the management of pests and parasites for 3 or more years.

Pitfall trapping was carried out from 13<sup>th</sup> June to 26<sup>th</sup> July 2016 on all 23 farms (as described by Sands & Wall, 2018). Traps were set up on pasture as previously described, between 09:00 and 12:00 noon, and left for 24 h. Ten traps were placed at each farm along a straight line transect, 5 m apart. The traps were separated from cattle by a fence to avoid damage, but were within 50 m of a field grazed by cattle during the trapping period. Traps were baited with homogenised dung collected from the nearest farm forgoing ML or SP use. After collection, *O. similis* were preserved in ethanol for later dissection of a random sample. Identification and dissections were carried out as described previously.

### *Data analysis*

Data were analysed using the statistical package RStudio (Version 3.4.4; R Core Team, 2018). A one-way ANOVA was used to assess the differences in oocyte length and number between ivermectin treatment groups in the laboratory investigation. For the field investigation, binary logistic regression was carried out with female reproductive state (defined as gravid or non-gravid) as the dependent variable, to determine whether farm type (MLs, SPs or none) could predict the reproductive state of a female. All means are stated  $\pm$  standard error. Where data are non-normal, medians ( $\pm$  interquartile range) are stated.

## Results

### *Laboratory investigation*

Observations indicated that at the lower concentrations of ivermectin the fat body was large and white to light yellow (Fig. 1). At higher concentrations of ivermectin, the fat body was smaller and darker yellow. At the highest concentrations, the fat body and oocytes were brown. At higher concentrations of ivermectin, beetles appeared to be slower in their movements, taking longer to become active when removed from the dung for counting. At the end of the three-week trial, 3.7% ( $\pm 1.8$ ) of beetles died in the control group and 0, 4.8% ( $\pm 2.0$ ) and 7.5% ( $\pm 2.5$ ) died when exposed to dung containing 0.01 ppm, 0.1 ppm and 1 ppm of ivermectin, respectively. All beetles, excluding one individual, died in the third week of the experiment.

In total, 90% of female *O. similis* were gravid and each gravid female matured an average of 7.5 ( $\pm 0.21$ ) oocytes. The number of mature oocytes in gravid females did not differ between the treatment groups ( $F_{3,77} = 0.78, p = 0.51$ ). However, there was a significant difference in oocyte length between the treatment groups ( $F_{3,77} = 3.66, p = 0.016$ ; Fig. 2). Oocyte length was significantly smaller in beetles that were exposed to dung containing 1 ppm ivermectin than the control group (Tukey HSD:  $p < 0.02$ ), and beetles exposed to dung containing 0.01 ppm ivermectin (Tukey HSD:  $p < 0.05$ ; Fig. 2). There was no relationship between the size of hind tibia and oocyte length ( $t_{79} = 1.42, p = 0.16$ ).

### *Field investigation*

Amongst the *O. similis* trapped on farms, of which a sample was selected ( $n = 145$ ), only 6% were gravid. The median number of oocytes found per gravid female was 4 ( $\pm 1.5$ ), and the median basal length of oocyte was 1.44 mm ( $\pm 0.13$ ) ( $n = 9$ ). Farm type was a significant predictor of the number of gravid *O. similis* ( $\chi^2_{2,63} = 15.42, p < 0.001$ ; Fig. 3). On farms where no MLs or SPs were used, 25% of females trapped were gravid. In contrast, 1% of the females



trapped were gravid on farms that used SPs, and no gravid females were trapped at farms that used MLs.

*Onthophagus similis* is assumed to be a spring-breeding species, but we are unaware of published reports to this effect. To confirm this assumption, we also dissected female *O. similis* (n = 81) collected in a related study on 16 of these same farms from 15<sup>th</sup> August – 8<sup>th</sup> September 2016 (Sands & Wall, 2018). No collected individuals were gravid.

## Discussion

In the present study, the length of the basal oocyte of *O. similis* was significantly reduced when beetles were fed on dung spiked with ivermectin at 1 ppm (ww), compared to 0.01 ppm ivermectin and the control group (Fig. 1; Fig. 2). The higher concentration corresponds to the peak residue concentration of ivermectin in dung 3 days after pour-on treatment to cattle with a standard dose of 200 µg/kg of ivermectin (Sommer *et al.*, 1992). There was also a significant association between the use of chemicals on farms and the percentage of *O. similis* that were gravid; farms using MLs and SPs had virtually no gravid females.

Other studies have demonstrated similar results. For example, a bioassay using *Euoniticellus intermedius* (Reiche), fed with ivermectin-spiked dung, showed a relationship between oocyte length and ivermectin concentration, with the smallest basal oocytes recorded at the highest ivermectin concentration of 0.3 ppm (ww) (Martinez *et al.*, 2017). Reduced oocyte length has also been demonstrated with exposure of beetles to the MLs doramectin and abamectin (Dadour *et al.*, 2000). Effects of MLs on reproductive output have also been demonstrated. Significantly fewer brood masses were made by *E. intermedius* in dung collected from cattle 1 to 14 days after a 200 µg/kg injection of ivermectin (Kruger and Scholz, 1997), and when *E. intermedius* was exposed to dung spiked with 0.1 ppm of ivermectin (Cruz Rosales *et al.*, 2012). Brood ball production of *Onthophagus binodis* (Thunberg) was reduced by 67% in dung

collected 1 week after cattle were treated by injection with avermectin B1 at 200 µg/kg (Ridsdill-Smith, 1988). Similar effects have been noted with SPs (Wardhaugh *et al.*, 1998; Vale *et al.*, 2004; Bang *et al.*, 2007), for example brood ball production by *Metacatharsius troglodytes* (Boheman) was reduced by 40% and 80% when exposed to dung spiked with 0.1 ppm (ww) and 1ppm (ww) of deltamethrin respectively, compared to controls (Sands *et al.*, 2018).

The reduced oocyte size in the present study may result from delays in development in response to ML residues. This may be due to suppressed feeding activity, which has been demonstrated in several species in the presence of ivermectin (Wardhaugh & Rodriguez-Menendez, 1988; Finnegan *et al.*, 1997). Suppressed feeding in the presence of ivermectin may result from paralysis of muscles used to process food or for locomotion. Additionally, detection of volatiles by *Scarabaeus cicatricosus* (Lucas) antennae has been shown to be significantly reduced when dung was spiked with 0.001 to 0.2 ppm ivermectin compared to controls (Verdú *et al.*, 2015). Dung patches are highly ephemeral so altered volatile sensitivity could reduce the ability of beetles to find food and mates. Suppressed feeding may result in a smaller fat body and therefore reduced energy reserves for oocyte production. Indeed, the size of the fat body appeared reduced in the current study, possibly resulting in the reduced oocyte size.

The combined lethal and sublethal effects of MLs and SPs at the egg, larval or adult stage are likely to reduce the long-term abundance of dung-beetles on farms using these chemical treatments. Models based on lethal effects, parametrized with laboratory data, have suggested that a single treatment of the ML eprinomectin could reduce the abundance of *Onthophagus taurus* (Schreber) by 35% when cattle were treated 14 days after the emergence of beetles (Wardhaugh *et al.*, 2001). The inclusion of sublethal effects to such models are likely to result in predictions of even greater reductions in abundance.

Previous work has highlighted an association between the long-term use of MLs and SPs and reduced beetle species richness (Sands & Wall, 2018). Effects were predominantly on paracoprids, such as *Onthophagus*, while paradoxically finding a higher abundance of endocoprids, such as *Aphodius*, on farms that used MLs or SPs. The authors speculated that competitive interactions between the beetle guilds may be responsible, along with differential sensitivity to chemicals. For example, Gittings and Giller (1999) suggested that the presence of the paracoprid *Geotrupes spiniger* (Marsham), in combination with weather conditions, may have prevented the development of *Aphodius* larvae by promoting the rapid removal of dung from pastures in Ireland in summer. Hence, when paracoprids such as *Onthophagus* species are absent, due to their sensitivity to MLs and SPs, the resource may become dominated by *Aphodius* and therefore, the diversity of species on pasture reduced.

There is evidence that more complex beetle communities achieve the greatest differences in ecosystem services. Over a period of 36 weeks, mixed species assemblages of *Aphodius* and *Onthophagus* dung beetles were shown to achieve the highest cattle dung decomposition rates in comparison to monoculture treatments, when the biomass of beetles was standardised (Beynon *et al.*, 2012). A similar conclusion was reached in a study by Manning *et al.* (2017) where the dung-removing capacity of *Aphodius ater* (De Geer), *Aphodius fossor* (L.) and *Onthophagus joannae* (Goljan) was assessed in polycultures or monocultures. The three species treatment and the monoculture containing *A. fossor* removed a larger quantity of dung than the other two monoculture treatments. A mixed community including the paracoprid beetle *Onthophagus nuchicornis* (L.) was also found to remove a greater quantity of dung than the endocoprids, *Aphodius erraticus* (L.) and *A. fossor* in monocultures (Manning & Cutler, 2018). Mixed dung beetle assemblages were shown to result in 32% lower CO<sub>2</sub> equivalent emissions compared to monoculture treatments (Piccini *et al.*, 2017).

However, some caution should be taken with the results from the long-term study; the sample size was low, only finding a total of 9 gravid females. To gain more confidence in the results, the study should be repeated, perhaps with a larger sample of beetles.

Overall, these studies suggest that dung beetle species are not functionally redundant; species rich assemblages, including the presence of paracoprid beetles such as *O. similis*, may contribute disproportionately towards ecosystem function. The sublethal effects on reproduction observed in the present study may therefore result in greater alterations to dung beetle communities over time than from lethal effects alone. Reductions in dung beetle species richness and/or abundance resulting from the combined lethal and sublethal effects of MLs and SPs are therefore likely to reduce ecosystem function on pastures.

### **Acknowledgements**

We would like to thank Mike and Paul Amos for allowing access to their farm for dung collection and the many farmers who allowed pitfall trapping in their pastures.

### **References**

- Bang, H.S., Lee, J.H., Na, Y.E. & Wall, R.** (2007) Reproduction of the dung beetle (*Copris tripartitus*) in the dung of cattle treated with cis-cypermethrin and chlorpyrifos. *Applied Soil Ecology* **35**, 546-552.
- Bang, H.S., Lee, J-H, Kwon, O.S., Na, Y. E., Jang, Y. S. & Kim, W.H.** (2005) Effects of paracoprid dung beetles (Coleoptera: Scarabaeidae) on the growth of pasture herbage and on the underlying soil. *Applied Soil Ecology* **29**, 165-171.

- Beynon, S.A., Mann, D.J., Slade, E.M. & Lewis, O.T.** (2012) Species-rich dung beetle communities buffer ecosystem services in perturbed agro-ecosystems. *Journal of Applied Ecology* **49**, 1365-1372.
- Beynon, S.A., Wainwright, W.A. & Christie, M.** (2015) The application of an ecosystem services framework to estimate the economic value of dung beetles to the UK cattle industry. *Ecological Entomology* **40**, 124-135.
- Biström, O., Silfverberg H. & Rutanen, I.** (1991) Abundance and distribution of coprophilous Histerini (Histeridae) and Onthophagus and Aphodius (Scarabaeidae) in Finland (Coleoptera). *Entomologica Fennica* **2**, 53-66.
- Bornemissza, G.F.** (1970) Insectary studies on the control of dung breeding flies by the activity of the dung beetle, *Onthophagus gazella* F. (Coleoptera: Scarabaeinae). *Austral Entomology* **9**, 31-41.
- Campbell, W.C.** (1989) Ivermectin and Abamectin. Springer, New York.
- Cruz Rosales, M., Martínez, M., López-Collado, J., Vargas-Mendoza, M., González-Hernández, H. & Fajersson, P.** (2012) Effect of ivermectin on the survival and fecundity of *Euoniticellus intermedius* (Coleoptera: Scarabaeidae). *Revista de Biología Tropical* **60**, 333-345.
- Dadour, I.R., Cook, D.F. & Hennessy, D.** (2000) Reproduction and survival of the dung beetle *Onthophagus binodis* (Coleoptera: Scarabaeidae) exposed to abamectin and doramectin residues in cattle dung. *Environmental Entomology* **29**, 1116-1122.
- Eliassen, A.** (2019) Available:  
<https://futureboy.us/fsp/dollar.fsp?quantity=367&cy=pounds&fromYear=2015>. Last accessed 26th May 2019.
- Filgueiras, B.K., Iannuzzi, L. & Leal, I.R.** (2011) Habitat fragmentation alters the structure of dung beetle communities in the Atlantic Forest. *Biological Conservation* **144**, 362-369.
- Fincher, G.T.** (1981) The potential value of dung beetles in pasture ecosystems. *Journal of the Georgia Entomological Society* **16**, 316-333.

- Finnegan, P.M., Flanagan, D.P. & Gormally, M.J.** (1997) Preliminary investigations of *Aphodius* species activity in cattle faeces treated with ivermectin. *Medical and Veterinary Entomology* **11**, 139-142.
- Floate, K.D., Wardhaugh, K.G., Boxall, A.B. & Sherratt, T.N.** (2005) Fecal residues of veterinary parasiticides: non-target effects in the pasture environment. *Annual Review of Entomology* **50**, 153-179.
- Hutton, S.A. & Giller, P.S.** (2003) The effects of the intensification of agriculture on northern temperate dung beetle communities. *Journal of Applied Ecology* **40**, 994-1007.
- Jacobs, C.T. & Scholtz, C.H.** (2015) A review on the effect of macrocyclic lactones on dung-dwelling insects: Toxicity of macrocyclic lactones to dung beetles. *Onderstepoort Journal of Veterinary Research* **82**, 1-8.
- Jessop, L.** (1986) Dung Beetles and Chafers Coleoptera: Scarabaeoidea. Handbooks for the Identification of British Insects Vol. 5 Part 11, Royal Entomological Society of London, London, UK.
- Knapp, M., Knappová, J. & Miller, T.** (2013) Measurement of body condition in a common carabid beetle, *Poecilus cupreus*: a comparison of fresh weight, dry weight, and fat content. *Journal of Insect Science* **13**, 1-10.
- Kruger, K. & Scholtz, C.H.** (1997) Lethal and sub-lethal effects of ivermectin on the dung-breeding beetles *Euoniticellus intermedius* (Reiche) and *Onitis alexis* Klug (Coleoptera: Scarabaeidae). *Agriculture, Ecosystems and Environment* **61**, 123-131.
- Lease, H.M. & Wolf, B.O.** (2011) Lipid content of terrestrial arthropods in relation to body size, phylogeny, ontogeny and sex. *Physiological Entomology* **36**, 29-38.
- Lobo, J.M.** (2001) Decline of roller dung beetle (Scarabaeinae) populations in the Iberian peninsula during the 20th century. *Biological Conservation* **97**, 43-50.
- Losey, J.E. & Vaughan, M.** (2006) The economic value of ecological services provided by insects. *AIBS Bulletin* **56**, 311-323.

- Mann, C.M., Barnes, S., Offer, B. & Wall, R.** (2015) Lethal and sub-lethal effects of faecal deltamethrin residues on dung-feeding insects. *Medical and Veterinary Entomology* **29**, 189-195.
- Manning, P. & Cutler, G.C.** (2018) Ecosystem functioning is more strongly impaired by reducing dung beetle abundance than by reducing species richness. *Agriculture, Ecosystems and Environment* **264**, 9-14.
- Manning, P., Beynon, S.A. & Lewis, O.T.** (2017) Quantifying immediate and delayed effects of anthelmintic exposure on ecosystem functioning supported by a common dung beetle species. *PloS one* **12**, e0182730.
- Martinez, I., Lumaret, J.P., Zayas, R.O. & Kadiri, N.** (2017) The effects of sublethal and lethal doses of ivermectin on the reproductive physiology and larval development of the dung beetle *Euoniticellus intermedius* (Coleoptera: Scarabaeidae). *The Canadian Entomologist* **149**, 461-72.
- McKellar, Q.A. & Gokbulut, C.** (2012) Pharmacokinetic features of the antiparasitic macrocyclic lactones. *Current Pharmaceutical Biotechnology* **13**, 888-911.
- Natural England** (2016) A review of the status of the beetles of Great Britain: The stag beetles, dor beetles, dung beetles, chafers and their allies - Lucanidae, Geotrupidae, Trogidae and Scarabaeidae. pp. 1-119.
- Nichols, E., Spector, S., Louzada, J., Larsen, T., Amezquita, S., Favila, M.E. and Network, T.R.** (2008) Ecological functions and ecosystem services provided by Scarabaeinae dung beetles. *Biological Conservation* **141**, 1461-1474.
- Nichols, E., Spector, S., Louzada, J., Larsen, T., Amezquita, S., Favila, M.E. & Network, T.S.R.** (2008) Ecological functions and ecosystem services provided by Scarabaeinae dung beetles. *Biological Conservation* **14**, 1461-1474.
- Pérez-Cogollo, L.C., Rodríguez-Vivas, R.I., Delfín-González, H., Reyes-Novelo, E. & Ojeda-Chi, M.M.** (2015) Lethal and sublethal effects of ivermectin on *Onthophagus landolti* (Coleoptera: Scarabaeidae). *Environmental Entomology* **44**, 1634-1640.

- Piccini, I., Arnieri, F., Caprio, E., Nervo, B., Pelissetti, S., Palestrini, C., Roslin, T. & Rolando, A.** (2017) Greenhouse gas emissions from dung pats vary with dung beetle species and with assemblage composition. *PLoS ONE* **12**, e0178077.
- Pickett, J.A.** (2004) New opportunities in neuroscience, but a great danger that some may be lost. *Neurotox* **3**, 1-10.
- R Core Team** (2018) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
- Ridsdill-Smith, T.J.** (1988) Survival and reproduction of *Musca vetustissima* Walker (Diptera: Muscidae) and a scarabaeine dung beetle in dung of cattle treated with avermectin B1. *Australian Journal of Entomology* **27**, 175-178.
- Roslin, T., Forshage, M., Ødegaard, F., Ekblad, C. & Liljeberg, G.** (2014). Nordens dyngbaggar (Dung Beetles of Northern Europe) Tibiale, Helsinki.
- Sands, B. & Wall, R.** (2017) Dung beetles reduce livestock gastrointestinal parasite availability on pasture. *Journal of Applied Ecology* **54**, 1180-1189.
- Sands, B. & Wall, R.** (2018) Sustained parasiticide use in cattle farming affects functionally important dung beetle assemblages. *Agriculture Ecosystems and Environment* **265**, 226-235.
- Skidmore, P.** (1991) Insects of the British Cow Dung Community. Field Studies Council, Monfort Bridge, UK.
- Sommer, C., Steffansen, B., Nielsen, B., Grønvold, J., Jensen, V.K., Jespersen, B.J., Springborg, J. & Nansen, P.** (1992) Ivermectin excreted in cattle dung after subcutaneous injection or pour-on treatment: Concentrations and impact on dung fauna. *Bulletin of Entomological Research* **82**, 257-264.
- Stafford, K. & Coles, G.C.** (1999) Nematode control practices and anthelmintic resistance in dairy calves in the south west of England. *The Veterinary Record* **144**, 659.
- Sutherland, I.H. & Campbell, W.C.** (1990) Development, pharmacokinetics and mode of action of ivermectin. *Acta Leidensia* **59**, 161-168.



- Vale, G.A., Grant, I.F., Dewhurst, C.F. & Aigreau, D.** (2004) Biological and chemical assays of pyrethroids in cattle dung. *Bulletin of Entomological Research* **94**, 273-282.
- Vale, G.A., Hargrove, J.W., Chamisa, A., Grant, I.F. & Torr, S.J.** (2015) Pyrethroid treatment of cattle for tsetse control: reducing its impact on dung fauna. *PLoS Neglected Tropical Diseases*, **9**, e0003560.
- Venant, A., Belli, P., Borrel, S. & Mallet, J.** (1990) Excretion of deltamethrin in lactating dairy cows. *Food Additives and Contaminants* **7**, 535-543.
- Verdú, J.R., Cortez, V., Ortiz, A.J., González-Rodríguez, E., Martínez-Pinna, J., Lumaret, J.P., Lobo, J.M., Numa, C. & Sánchez-Piñero, F.** (2015) Low doses of ivermectin cause sensory and locomotor disorders in dung beetles. *Scientific Reports* **5**, e13912.
- Wall, R. & Strong, L.** (1987) Environmental consequences of treating cattle with the antiparasitic drug ivermectin. *Nature* **327**, 418-421.
- Wardhaugh, K.G. & Rodriguez-Menendez, H.** (1988) The effects of the antiparasitic drug, ivermectin, on the development and survival of the dung-breeding fly, *Orthelia cornicina* (F.) and the scarabaeine dung beetles *Copris hispanus* L., *Bubas bubalus* (Oliver) and *Onitis belial* F. *Journal of Applied Entomology* **106**, 381-389.
- Wardhaugh, K.G., Holter, P. & Longstaff, B.** (2001) The development and survival of three species of coprophagous insect after feeding on the faeces of sheep treated with controlled-release formulations of ivermectin or albendazole. *Australian Veterinary Journal* **79**, 125-132.
- Wardhaugh, K.G., Longstaff, B.C. & Lacey, M.J.** (1998) Effects of residues of deltamethrin in cattle faeces on the development and survival of three species of dung breeding insect. *Australian Veterinary Journal* **76**, 273-280.
- Yamada, D., Imura, O., Shi, K. & Shibuya, T.** (2007) Effect of tunneler dung beetles on cattle dung decomposition, soil nutrients and herbage growth. *Grassland Science* **53**, 121-129.

## Figure legends

**Fig. 1.** Ovary of *Onthophagus similis* when exposed to dung treated with ethanol only (control - C) or ivermectin at 1 ppm (T). Dung beetles were exposed for 3 weeks, with dung replaced weekly.

**Fig. 2.** The mean oocyte length ( $\pm 95\%$  confidence intervals) of gravid *Onthophagus similis* after three weeks, when exposed to dung treated with ethanol only (control) or ivermectin at 0.01, 0.1, or 1 ppm. Letters above bars indicate significant differences between means as indicated by Tukey Multiple Range tests.

**Fig. 3** Percentage of gravid *Onthophagus similis* beetles collected by pitfall trapping during the summer of 2016 on farms with different pesticide/parasiticide history (none (n = 33), macrocyclic lactones (n = 26), synthetic pyrethroids (n = 86)) in South-West England. Percentages are calculated for each farm type. Bars display the standard error.

Fig 1




Fig 2




Fig 3.

