



Aris, I. M., Rifas-Shiman, S. L., Zhang, X., Yang, S., Switkowski, K., Fleisch, A. F., Hivert, M-F., Martin, R., Kramer, M. S., & Oken, E. (2019). Association of BMI with Linear Growth and Pubertal Development. *Obesity*, 27(10), 1661-1670.
<https://doi.org/10.1002/oby.22592>

Peer reviewed version

License (if available):
Unspecified

Link to published version (if available):
[10.1002/oby.22592](https://doi.org/10.1002/oby.22592)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the accepted author manuscript (AAM). The final published version (version of record) is available online via Wiley at <https://doi.org/10.1002/oby.22592>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Body Mass Index and its Association with Linear Growth and Pubertal Development

Izzuddin M Aris^{1,2,3}, Sheryl L. Rifas-Shiman¹, Xun Zhang⁴, Seungmi Yang⁵, Karen Switkowski¹, Abby F Fleisch^{6,7}, Marie-France Hivert^{1,8}, Richard M Martin^{9,10}, Michael S Kramer^{2,5,11}, Emily Oken^{1,12}

Author Affiliations

¹ Division of Chronic Disease Research Across the Lifecourse, Department of Population Medicine, Harvard Medical School and Harvard Pilgrim Health Care Institute, Boston, Massachusetts, USA

² Department of Obstetrics and Gynecology, Yong Loo Lin School of Medicine, National University of Singapore, Singapore, Singapore

³ Singapore Institute for Clinical Sciences, Agency for Science, Technology and Research, Singapore, Singapore

⁴ Centre for Outcomes Research and Evaluation, McGill University Health Centre Research Institute, Montreal, Canada

⁵ Department of Epidemiology, Biostatistics and Occupational Health, Faculty of Medicine McGill University, Montreal, Quebec, Canada

⁶ Pediatric Endocrinology and Diabetes, Maine Medical Center, Portland, Maine, USA

⁷ Center for Outcomes Research and Evaluation, Maine Medical Center Research Institute, Portland, Maine, USA

⁸ Diabetes Unit, Massachusetts General Hospital, Boston, Massachusetts, USA

⁹ Bristol Medical School, Department of Population Health Sciences, University of Bristol, Bristol, UK

¹⁰ National Institute for Health Research Bristol Biomedical Research Centre, University Hospitals NHS Trust and University of Bristol, Bristol, UK

¹¹ Department of Pediatrics, Faculty of Medicine McGill University, Montreal, Quebec, Canada

¹² Department of Nutrition, Harvard T.H. Chan School of Public Health, Boston, Massachusetts, USA


Address correspondence to:

Izzuddin M ARIS. Division of Chronic Disease Research Across the Lifecourse, Department of Population Medicine, Harvard Medical School and Harvard Pilgrim Health Care Institute. 401 Park Drive, Suite 401E, Boston, MA 02215. Email: Izzuddin_Aris@harvardpilgrim.org

SUPPLEMENTARY MATERIAL


Figure S1: Complete-case analyses of associations of BMI z-score with standardized length/height velocity in (A) PROBIT and (B) Project Viva.

A



A = Length velocity 0–3 months; B = Length velocity 3–12 months; C: Length/height velocity 12 months to 6.5 years; D = Height velocity 6.5–11.5 years; E = Height velocity 11.5–16 years
▲ = boys; ● = girls

B



A' = Length velocity 0–6.3 months; B' = Length/height velocity 6.3 months to 3.2 years; C: Height velocity 3.2–7.7 years; D' = Height velocity 7.7–12.9 years
▲ = boys; ● = girl

SUPPLEMENTARY MATERIAL

Table S1: Associations of BMI z-score with standing height at 16 years, mediated by pubertal development at 11.5 years in PROBIT.

BMI z-score	Standing height at 16y (cm)		
	Model 1 ^a	Model 2 ^b	Model 3 ^c
Boys (n=8,682)		β (95% CI)	
0m	0.56 (0.40,0.73)	0.57 (0.40,0.73)	0.56 (0.40,0.73)
3m	-0.01 (-0.16,0.13)	-0.03 (-0.17,0.12)	-0.03 (-0.18,0.11)
12m	0.14 (-0.02,0.30)	0.12 (-0.04,0.28)	0.12 (-0.04,0.28)
6.5y	0.27 (0.12,0.42)	0.22 (0.07,0.37)	0.21 (0.06,0.36)
11.5y	0.29 (0.12,0.46)	0.22 (0.05,0.39)	0.21 (0.04,0.39)
Girls (n=8,099)			
0m	0.53 (0.38,0.68)	0.53 (0.39,0.68)	0.53 (0.39,0.68)
3m	-0.07 (-0.21,0.08)	-0.08 (-0.22,0.07)	-0.08 (-0.23,0.07)
12m	0.03 (-0.13,0.19)	0.01 (-0.15,0.17)	0.01 (-0.15,0.16)
6.5y	0.10 (-0.05,0.25)	0.03 (-0.13,0.18)	0.04 (-0.12,0.19)
11.5y	-0.13 (-0.29,0.04)	-0.25 (-0.42,-0.08)	-0.22 (-0.39,-0.06)

^a Adjusted for maternal and paternal education level, height and BMI; maternal age and smoking during pregnancy; gestational age at delivery; geographic region and BMI z-score at all previous time points.

^b Model 1 + genital (for boys) or breast (for girls) development at 11.5 years.

^c Model 1 + pubic hair stage at 11.5 years.

SUPPLEMENTARY MATERIAL

Table S2: Associations of overweight and obesity (vs. normal or underweight) with standardized length/height velocity in PROBIT.

	Standardized height velocity (SD units/year)				
	0-3m	3 - 12m	12m - 6.5y	6.5 - 11.5y	11.5 - 16y
Boys (n=8,682)					
Overweight					
(vs. normal or underweight)					
	β (95% CI)^a				
0m	0.45 (0.12,0.79)^b	-0.08 (-0.45,0.30)	0.05 (-0.34,0.44)	-0.17 (-0.58,0.23)	0.15 (-0.25,0.55)
3m	–	0.47 (0.32,0.61)	-0.01 (-0.16,0.14)	0.06 (-0.10,0.22)	-0.04 (-0.20,0.11)
12m	–	–	0.26 (0.20,0.32)	-0.00 (-0.06,0.06)	-0.05 (-0.11,0.01)
6.5y	–	–	–	0.17 (0.11,0.24)	-0.26 (-0.33,-0.20)
11.5y	–	–	–	–	-0.15 (-0.20,-0.10)
Obese					
(vs. normal or underweight)					
0m	3.42 (2.45,4.38)	0.71 (-0.37,1.79)	-0.57 (-1.61,0.47)	-0.19 (-1.23,0.86)	-0.19 (-1.18,0.80)
3m	–	0.40 (-0.05,0.85)	-0.01 (-0.50,0.48)	0.21 (-0.32,0.73)	-0.13 (-0.60,0.35)
12m	–	–	0.45 (0.33,0.58)	0.02 (-0.12,0.16)	-0.17 (-0.31,-0.03)
6.5y	–	–	–	0.24 (0.14,0.35)	-0.42 (-0.52,-0.31)
11.5y	–	–	–	–	-0.21 (-0.29,-0.13)
Girls (n=8,099)					
Overweight					
(vs. normal or underweight)					
0m	0.44 (0.07,0.81)	-0.11 (-0.52,0.30)	-0.09 (-0.52,0.34)	0.05 (-0.37,0.47)	-0.13 (-0.55,0.30)
3m	–	0.37 (0.20,0.55)	0.04 (-0.16,0.24)	-0.10 (-0.31,0.11)	0.03 (-0.18,0.23)
12m	–	–	0.25 (0.18,0.32)	-0.05 (-0.13,0.02)	-0.07 (-0.14,-0.01)
6.5y	–	–	–	0.18 (0.10,0.26)	-0.34 (-0.42,-0.26)
11.5y	–	–	–	–	-0.14 (-0.20,-0.09)
Obese					
(vs. normal or underweight)					
0m	1.17 (0.40,1.95)	0.35 (-0.49,1.18)	-0.77 (-1.65,0.12)	-0.51 (-1.38,0.36)	0.84 (-0.05,1.72)
3m	–	0.92 (0.39,1.45)	0.03 (-0.52,0.58)	0.08 (-0.48,0.64)	0.10 (-0.45,0.65)
12m	–	–	0.49 (0.34,0.65)	-0.07 (-0.23,0.10)	-0.12 (-0.28,0.04)
6.5y	–	–	–	0.28 (0.14,0.42)	-0.51 (-0.64,-0.37)
11.5y	–	–	–	–	-0.16 (-0.25,-0.06)

^a Adjusted for maternal and paternal education level, height and BMI; maternal age and smoking during pregnancy; gestational age at delivery; geographic region and BMI z-score at all previous time points.

^b Bold values indicate statistically significant associations at p<0.05.

SUPPLEMENTARY MATERIAL

Table S3: Associations of overweight and obesity (vs. normal or underweight) with standardized length/height velocity in Project Viva.

	Standardized height velocity (SD units/year)			
	0-6.3m	6.3m - 3.2y	3.2 - 7.7y	7.7 - 12.9y
Boys (n=758)	β (95% CI)^a			
Overweight (vs. normal or underweight)				
0m	0.06 (-0.28,0.40)	-0.15 (-0.53,0.23)	-0.01 (-0.40,0.38)	-0.12 (-0.53,0.28)
6m	–	0.31 (0.03,0.58)^b	0.04 (-0.25,0.34)	-0.01 (-0.32,0.30)
3.2y	–	–	0.17 (-0.12,0.47)	-0.12 (-0.48,0.24)
7.7y	–	–	–	0.08 (-0.17,0.33)
Obese (vs. normal or underweight)				
0m	-0.07 (-1.18,1.05)	-0.36 (-1.51,0.78)	0.10 (-1.15,1.36)	0.35 (-0.99,1.69)
6m	–	0.41 (-0.10,0.91)	-0.36 (-0.88,0.16)	-0.00 (-0.54,0.54)
3.2y	–	–	0.32 (-0.21,0.84)	0.08 (-0.45,0.62)
7.7y	–	–	–	-0.01 (-0.34,0.32)
Girls (n=732)				
Overweight (vs. normal or underweight)				
0m	-0.02 (-0.45,0.41)	-0.13 (-0.54,0.29)	0.16 (-0.26,0.58)	0.00 (-0.44,0.45)
6m	–	0.13 (-0.17,0.43)	-0.22 (-0.55,0.12)	0.17 (-0.22,0.55)
3.2y	–	–	0.10 (-0.26,0.45)	-0.18 (-0.61,0.26)
7.7y	–	–	–	-0.10 (-0.33,0.14)
Obese (vs. normal or underweight)				
0m	1.11 (-0.02,2.24)	-0.25 (-1.17,0.67)	0.22 (-0.80,1.25)	-0.31 (-1.45,0.84)
6m	–	0.24 (-0.52,1.00)	-0.02 (-0.90,0.86)	-0.10 (-1.09,0.90)
3.2y	–	–	-0.06 (-0.76,0.63)	-0.36 (-1.08,0.37)
7.7y	–	–	–	-0.34 (-0.66,-0.01)

^a Adjusted for maternal and paternal education level, height and BMI; maternal age, smoking history and parity; gestational age at delivery; child ethnicity; breastfeeding initiation and BMI z-score at all previous time points.

^b Bold values indicate statistically significant associations at p<0.05.

SUPPLEMENTARY MATERIAL

Table S4: Associations of overweight and obesity (vs. normal or underweight) with standing height, trunk and leg length, and pubertal development at 11.5 and 16 years in PROBIT.

	β (95% CI) ^a				OR (95% CI) ^{a,b}		
	11.5y			16y	11.5y		16y
	Standing height (cm)	Trunk length (cm)	Leg length (cm)	Standing height (cm)	Genital development	Pubic hair stage	Pubic hair stage
Boys (n=8,682)							
Overweight (vs. normal or underweight)							
0m	0.24 (-2.52,3.00)	0.05 (-1.58,1.68)	0.18 (-1.75,2.11)	1.58 (-1.08,4.25)	1.06 (0.38,2.94)	0.52 (0.16,1.65)	1.91 (0.64,5.75)
3m	0.27 (-0.81,1.34)	0.74 (0.09,1.39)	-0.47 (-1.19,0.25)	-0.13 (-1.17,0.92)	1.18 (0.83,1.69)	1.26 (0.88,1.80)	1.20 (0.86,1.69)
12m	0.52 (0.11,0.93)^c	0.38 (0.13,0.63)	0.14 (-0.14,0.42)	0.32 (-0.08,0.71)	1.05 (0.92,1.21)	1.20 (1.05,1.38)	1.06 (0.93,1.21)
6.5y	2.28 (1.82,2.74)	1.63 (1.36,1.90)	0.65 (0.34,0.95)	0.39 (-0.04,0.83)	1.44 (1.24,1.67)	1.61 (1.39,1.86)	1.23 (1.06,1.43)
11.5y	2.38 (1.96,2.80)	1.60 (1.36,1.85)	0.78 (0.50,1.06)	0.44 (0.04,0.85)	1.55 (1.34,1.78)	1.68 (1.46,1.94)	1.23 (1.07,1.41)
16y	—	—	—	0.23 (-0.23,0.68)	—	—	1.14 (0.97,1.33)
Obese (vs. normal or underweight)							
0m	-1.49 (-8.76,5.79)	1.67 (-2.62,5.95)	-3.18 (-8.01,1.65)	-3.93 (-10.80,2.94)	0.85 (0.06,11.46)	1.84 (0.22,15.53)	2.38 (0.22,25.47)
3m	0.40 (-3.07,3.88)	0.35 (-1.69,2.40)	0.05 (-2.28,2.38)	0.45 (-2.73,3.63)	0.95 (0.32,2.81)	1.09 (0.34,3.48)	0.87 (0.32,2.34)
12m	0.51 (-0.41,1.42)	0.86 (0.32,1.40)	-0.35 (-0.95,0.24)	-0.23 (-1.12,0.66)	1.09 (0.82,1.46)	1.25 (0.93,1.68)	0.92 (0.68,1.25)
6.5y	4.63 (3.91,5.34)	3.28 (2.85,3.71)	1.35 (0.85,1.84)	1.18 (0.47,1.90)	1.27 (0.99,1.63)	1.54 (1.20,1.98)	1.24 (0.97,1.59)
11.5y	4.05 (3.42,4.67)	2.64 (2.29,3.00)	1.40 (0.97,1.84)	1.15 (0.52,1.79)	1.02 (0.83,1.26)	1.35 (1.09,1.67)	1.27 (1.03,1.57)
16y	—	—	—	0.17 (-0.59,0.92)	—	—	0.98 (0.76,1.26)
Girls (n=8,099)							
Overweight (vs. normal or underweight)							
0m	-0.08 (-3.09,2.93)	0.23 (-1.57,2.03)	-0.31 (-2.25,1.63)	-1.71 (-4.14,0.72)	1.46 (0.62,3.42)	1.28 (0.57,2.86)	1.39 (0.63,3.07)
3m	-0.34 (-1.77,1.09)	0.09 (-0.77,0.95)	-0.42 (-1.33,0.49)	0.11 (-1.08,1.29)	1.08 (0.71,1.65)	1.35 (0.90,2.03)	0.97 (0.65,1.42)
12m	0.14 (-0.36,0.64)	0.26 (-0.04,0.56)	-0.12 (-0.43,0.18)	-0.10 (-0.49,0.29)	1.12 (0.97,1.30)	1.07 (0.93,1.23)	0.99 (0.86,1.13)
6.5y	2.90 (2.36,3.43)	2.07 (1.74,2.39)	0.82 (0.48,1.16)	0.16 (-0.30,0.62)	2.38 (2.02,2.81)	2.19 (1.87,2.58)	1.18 (1.01,1.37)
11.5y	2.09 (1.61,2.57)	1.62 (1.34,1.91)	0.46 (0.16,0.77)	-0.05 (-0.45,0.35)	2.53 (2.18,2.94)	2.16 (1.87,2.49)	1.11 (0.96,1.27)
16y	—	—	—	-0.64 (-1.08,-0.20)	—	—	1.08 (0.94,1.25)
Obese (vs. normal or underweight)							
0m	-6.01 (-12.32,0.29)	-3.04 (-6.87,0.79)	-2.98 (-6.97,1.00)	-0.46 (-5.63,4.72)	0.36 (0.05,2.48)	0.22 (0.03,1.83)	1.03 (0.10,10.35)
3m	0.62 (-3.36,4.60)	0.38 (-2.03,2.80)	0.22 (-2.29,2.73)	-0.06 (-3.24,3.11)	1.33 (0.43,4.15)	1.96 (0.62,6.27)	1.01 (0.37,2.74)
12m	0.22 (-0.93,1.37)	0.08 (-0.62,0.78)	0.13 (-0.60,0.86)	-0.24 (-1.16,0.68)	1.20 (0.85,1.68)	1.33 (0.95,1.85)	1.03 (0.75,1.43)
6.5y	4.72 (3.76,5.68)	3.13 (2.56,3.71)	1.58 (0.95,2.21)	0.29 (-0.51,1.08)	3.76 (2.82,5.00)	2.87 (2.16,3.82)	1.09 (0.83,1.43)
11.5y	2.40 (1.56,3.23)	2.14 (1.64,2.64)	0.26 (-0.28,0.81)	-0.01 (-0.72,0.70)	3.03 (2.33,3.95)	2.13 (1.66,2.75)	1.04 (0.82,1.32)
16y	—	—	—	-0.42 (-1.32,0.47)	—	—	1.03 (0.76,1.39)

^a Adjusted for maternal and paternal education level, height and BMI; maternal age and smoking during pregnancy; gestational age at delivery; geographic region; BMI z-score at all previous time points and age at measurement.

^b OR (odds ratio) reflects relative odds of being in a higher Tanner stage category.

^c Bold values indicate statistically significant associations at p<0.05.

SUPPLEMENTARY MATERIAL

Table S5: Associations of overweight and obesity (vs. normal or underweight) with standing height, trunk and leg length, and pubertal development at 12.9 years in Project Viva.

	β (95% CI) ^a			OR (95% CI) ^{a,b}	
	Standing height (cm)	Trunk length (cm)	Leg length (cm)	Pubertal score (units)	Pubic hair stage
<i>Boys (n=758)</i>					
Overweight (vs. normal or underweight)					
0m	-0.67 (-3.59,2.25)	0.09 (-1.64,1.82)	-0.76 (-2.62,1.10)	-0.10 (-0.36,0.15)	1.00 (0.49,2.03)
6.3m	0.77 (-1.45,3.00)	0.72 (-0.58,2.02)	0.05 (-1.35,1.45)	0.06 (-0.13,0.26)	1.21 (0.68,2.17)
3.2y	1.09 (-1.26,3.45)	1.03 (-0.37,2.42)	0.07 (-1.43,1.57)	0.05 (-0.16,0.25)	0.78 (0.39,1.59)
7.7y	2.70 (1.01,4.40)^c	0.99 (-0.05,2.02)	1.72 (0.65,2.79)	0.21 (0.07,0.35)	1.71 (1.07,2.72)
12.9y	1.21 (-0.44,2.86)	1.46 (0.48,2.43)	-0.25 (-1.28,0.79)	0.10 (-0.04,0.25)	1.21 (0.75,1.95)
Obese (vs. normal or underweight)					
0m	2.03 (-8.16,12.21)	1.87 (-4.75,8.50)	0.15 (-6.09,6.40)	0.22 (-0.71,1.14)	1.51 (0.14,16.46)
6.3m	-1.05 (-5.24,3.14)	-0.14 (-2.66,2.38)	-0.90 (-3.48,1.67)	0.01 (-0.32,0.35)	0.72 (0.25,2.08)
3.2y	5.09 (1.13,9.06)	2.91 (0.52,5.30)	2.18 (-0.42,4.78)	0.15 (-0.19,0.49)	1.77 (0.62,5.06)
7.7y	4.17 (1.94,6.39)	1.58 (0.20,2.96)	2.58 (1.10,4.06)	0.25 (0.06,0.45)	2.37 (1.23,4.59)
12.9y	0.71 (-1.83,3.25)	1.56 (0.02,3.10)	-0.85 (-2.45,0.74)	0.13 (-0.08,0.35)	0.94 (0.47,1.87)
<i>Girls (n=732)</i>					
Overweight (vs. normal or underweight)					
0m	0.42 (-2.57,3.41)	0.31 (-1.43,2.04)	0.11 (-1.78,2.01)	0.02 (-0.26,0.30)	1.08 (0.42,2.81)
6.3m	0.45 (-1.90,2.80)	0.49 (-0.99,1.98)	-0.04 (-1.57,1.49)	0.10 (-0.12,0.32)	1.57 (0.71,3.46)
3.2y	0.64 (-1.98,3.27)	0.68 (-0.89,2.25)	-0.04 (-1.71,1.62)	0.06 (-0.18,0.29)	1.10 (0.46,2.66)
7.7y	1.94 (0.42,3.45)	1.25 (0.32,2.19)	0.68 (-0.28,1.64)	0.24 (0.10,0.38)	1.59 (0.99,2.54)
12.9y	0.91 (-0.66,2.48)	1.07 (0.11,2.03)	-0.16 (-1.14,0.82)	0.22 (0.08,0.37)	1.28 (0.79,2.09)
Obese (vs. normal or underweight)					
0m	-1.19 (-8.24,5.86)	-0.51 (-4.84,3.82)	-0.68 (-5.37,4.01)	-0.15 (-0.77,0.48)	0.27 (0.04,1.94)
6.3m	-0.65 (-6.36,5.05)	0.05 (-3.53,3.63)	-0.70 (-4.42,3.01)	0.06 (-0.48,0.60)	1.12 (0.19,6.74)
3.2y	-1.09 (-5.92,3.74)	-0.44 (-3.36,2.48)	-0.65 (-3.74,2.45)	0.04 (-0.40,0.47)	1.41 (0.32,6.21)
7.7y	1.32 (-0.76,3.39)	1.05 (-0.24,2.34)	0.27 (-1.03,1.57)	0.22 (0.03,0.42)	1.87 (0.96,3.64)
12.9y	0.92 (-1.39,3.23)	1.27 (-0.18,2.73)	-0.35 (-1.81,1.11)	0.18 (-0.04,0.40)	1.31 (0.61,2.81)

^a Adjusted for maternal and paternal education level, height and BMI; maternal age, smoking history and parity; gestational age at delivery; child ethnicity; breastfeeding initiation; BMI z-score at all previous time points and age at measurement.

^b OR (odds ratio) reflects relative odds of being in a higher Tanner stage category.

^c Bold values indicate statistically significant associations at p<0.05.

SUPPLEMENTARY MATERIAL

Table S6: Complete case analyses of associations of BMI z-score with standing height, trunk and leg length, and pubertal development at 11.5 and 16 years in PROBIT.

BMI z-score	β (95% CI) ^a				OR (95% CI) ^{a,b}		
	11.5y			16y	11.5y		16y
	Standing height (cm)	Trunk length (cm)	Leg length (cm)	Standing height (cm)	Genital development	Pubic hair stage	Pubic hair stage
Boys							
0m	0.39 (0.20,0.58)^c	0.23 (0.12,0.35)	0.16 (0.04,0.29)	0.57 (0.39,0.76)	1.06 (1.00,1.14)	1.05 (0.99,1.13)	1.00 (0.94,1.07)
3m	0.09 (-0.08,0.26)	0.12 (0.02,0.22)	-0.04 (-0.15,0.07)	-0.03 (-0.19,0.13)	1.01 (0.96,1.07)	1.06 (1.00,1.13)	1.08 (1.02,1.14)
12m	0.35 (0.17,0.54)	0.29 (0.19,0.40)	0.05 (-0.08,0.17)	0.23 (0.05,0.42)	1.06 (1.00,1.13)	1.09 (1.02,1.16)	1.02 (0.96,1.08)
6.5y	1.36 (1.19,1.53)	1.01 (0.91,1.10)	0.36 (0.25,0.48)	0.31 (0.14,0.48)	1.18 (1.11,1.25)	1.25 (1.17,1.32)	1.09 (1.03,1.15)
11.5y	1.67 (1.49,1.84)	1.16 (1.06,1.26)	0.52 (0.39,0.64)	0.41 (0.22,0.60)	1.31 (1.23,1.40)	1.43 (1.34,1.53)	1.28 (1.20,1.37)
16y	—	—	—	0.23 (-0.01,0.47)	—	—	1.28 (1.19,1.39)
Breast development							
Girls							
0m	0.33 (0.13,0.53)	0.15 (0.03,0.28)	0.19 (0.06,0.32)	0.54 (0.38,0.70)	1.00 (0.94,1.06)	0.97 (0.91,1.03)	0.97 (0.92,1.03)
3m	0.09 (-0.11,0.29)	0.20 (0.08,0.32)	-0.11 (-0.24,0.02)	-0.05 (-0.21,0.10)	1.11 (1.04,1.17)	1.12 (1.05,1.18)	1.03 (0.97,1.08)
12m	0.37 (0.15,0.59)	0.30 (0.17,0.43)	0.08 (-0.06,0.22)	0.04 (-0.14,0.21)	1.13 (1.06,1.21)	1.13 (1.06,1.20)	1.04 (0.98,1.10)
6.5y	1.64 (1.44,1.83)	1.19 (1.08,1.31)	0.45 (0.33,0.58)	0.01 (-0.15,0.17)	1.75 (1.65,1.86)	1.61 (1.51,1.71)	1.07 (1.01,1.13)
11.5y	1.65 (1.45,1.85)	1.32 (1.20,1.43)	0.33 (0.20,0.46)	-0.16 (-0.33,0.01)	2.50 (2.33,2.68)	2.00 (1.87,2.14)	1.08 (1.02,1.15)
16y	—	—	—	-0.66 (-0.87,-0.45)	—	—	1.07 (1.00,1.15)

^a Adjusted for maternal and paternal education level, height and BMI; maternal age and smoking during pregnancy; gestational age at delivery; geographic region; BMI z-score at all previous time points and age at measurement.

^b OR (odds ratio) reflects relative odds of being in a higher Tanner stage category.

^c Bold values indicate statistically significant associations at p<0.05.

SUPPLEMENTARY MATERIAL

Table S7: Complete case analyses of associations of BMI z-score with standing height, trunk and leg length, and pubertal development at 12.9 years in Project Viva.

BMI z-score	β (95% CI) ^a			OR (95% CI) ^{a,b}	
	Standing height (cm)	Trunk length (cm)	Leg length (cm)	Pubertal score (units)	Pubic hair stage
Boys					
Birth	-0.26 (-1.37,0.84)	0.11 (-0.49,0.71)	-0.40 (-1.12,0.33)	-0.01 (-0.09,0.07)	0.89 (0.67,1.19)
6.3m	-0.05 (-1.08,0.99)	-0.03 (-0.60,0.54)	-0.03 (-0.71,0.64)	0.00 (-0.07,0.08)	1.17 (0.90,1.52)
3.2y	0.93 (-0.44,2.30)	0.94 (0.17,1.70)	-0.01 (-0.91,0.89)	-0.01 (-0.11,0.10)	0.88 (0.62,1.23)
7.7y	1.99 (0.46,3.52)^c	0.48 (-0.38,1.34)	1.51 (0.52,2.50)	0.13 (0.02,0.25)	1.39 (0.93,2.08)
12.9y	3.14 (1.50,4.78)	2.07 (1.16,2.98)	1.08 (-0.01,2.18)	0.13 (0.00,0.26)	1.50 (0.96,2.35)
Girls					
Birth	0.34 (-0.43,1.11)	-0.04 (-0.54,0.46)	0.38 (-0.12,0.88)	-0.01 (-0.10,0.07)	1.10 (0.83,1.47)
6.3m	0.82 (-0.08,1.72)	0.65 (0.09,1.20)	0.17 (-0.42,0.76)	0.07 (-0.03,0.17)	1.35 (0.96,1.89)
3.2y	0.40 (-0.69,1.49)	0.31 (-0.36,0.98)	0.09 (-0.59,0.77)	0.03 (-0.09,0.16)	1.05 (0.69,1.61)
7.7y	1.75 (0.44,3.05)	1.22 (0.42,2.03)	0.52 (-0.27,1.32)	0.18 (0.03,0.32)	1.68 (1.02,2.78)
12.9y	1.86 (0.52,3.20)	1.43 (0.62,2.24)	0.42 (-0.42,1.26)	0.40 (0.26,0.53)	1.76 (1.02,3.04)

^a Adjusted for maternal and paternal education level, height and BMI; maternal age, smoking history and parity; gestational age at delivery; child ethnicity; breastfeeding initiation; BMI z-score at all previous time points and age at measurement.

^b OR (odds ratio) reflects relative odds of being in a higher Tanner stage category.

^c Bold values indicate statistically significant associations at $p < 0.05$.