



the PRACTICAL Consortium (2020). Appraising causal relationships of dietary, nutritional and physical-activity exposures with overall and aggressive prostate cancer: two-sample Mendelian randomization study based on 79,148 prostate cancer cases and 61,106 controls. *International Journal of Epidemiology*, 49(2), 587-596.
<https://doi.org/10.1093/ije/dyz235>

Peer reviewed version

Link to published version (if available):
[10.1093/ije/dyz235](https://doi.org/10.1093/ije/dyz235)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Oxford University Press at <https://academic.oup.com/ije/advance-article/doi/10.1093/ije/dyz235/5650986?searchresult=1>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Appraising causal relationships of dietary, nutritional and physical-activity exposures with overall and aggressive prostate cancer: two-sample Mendelian randomization study based on 79,148 prostate cancer cases and 61,106 controls

Dr. Nabila Kazmi (PhD)^{1,2}, Dr. Philip Haycock (PhD)^{1,2}, Dr Konstantinos Tsilidis (PhD)^{3,4}, Assoc Prof Brigid M. Lynch (PhD)⁵⁻⁷, Dr. Therese Truong (PhD)⁸, the PRACTICAL consortium, CRUK, BPC3, CAPS, PEGASUS*, Prof Richard M Martin (PhD)^{1,2,9†}, Dr. Sarah Lewis (PhD)^{1,2†*}

¹MRC Integrative Epidemiology Unit (IEU), Bristol Medical School, University of Bristol, Bristol, UK

²Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK

³Department of Hygiene and Epidemiology, University of Ioannina School of Medicine, Ioannina, Greece

⁴Department of Epidemiology and Biostatistics, School of Public Health, Imperial College London, London, UK

⁵Cancer Epidemiology Division, Cancer Council Victoria, Melbourne, Australia

⁶Centre for Epidemiology and Biostatistics, Melbourne School of Population and Global Health, University of Melbourne, Melbourne, Australia

⁷Physical-activity Laboratory, Baker Heart and Diabetes Institute, Melbourne, Australia

⁸Université Paris-Saclay, Université Paris-Sud, CESP (Center for Research in Epidemiology and Population Health), Inserm, Team Cancer and Environment, Villejuif, France.

⁹National Institute for Health Research (NIHR) Bristol Biomedical Research Centre, University Hospitals NHS Trust and University of Bristol, Bristol, UK.

*Members from the PRACTICAL Consortium, CRUK, BPC3, CAPS and PEGASUS are provided in the supplement.

† These authors contributed equally

Contact details of the corresponding authors:

Dr Sarah Lewis

Email: s.j.lewis@bristol.ac.uk

Telephone: +44 (0) 117 3313316

Address: Oakfield House, Oakfield Grove

University of Bristol

Bristol, BS8 2BN

UK

SUMMARY

Background: Prostate cancer is the second most common male cancer worldwide, but there is substantial geographical variation suggesting a potential role for modifiable risk factors in prostate carcinogenesis.

Methods: We identified previously reported prostate cancer risk factors from the World Cancer Research Fund's (WCRF) systematic appraisal of the global evidence (2018). We assessed whether each identified risk factor was causally associated with risk of overall (79,148 cases and 61,106 controls) or aggressive (15,167 cases and 58,308 controls) prostate cancer using Mendelian randomization (MR) based on genome wide association study (GWAS) summary statistics from the PRACTICAL and GAME-ON/ELLIPSE consortia. We assessed evidence for replication in UK Biobank (7,844 prostate cancer cases and 204,001 controls).

Results: WCRF identified 57 potential risk factors, of which 22 could be instrumented for MR analyses using single nucleotide polymorphisms (SNPs). For overall prostate cancer, we identified evidence compatible with causality for the following risk factors (odds ratio [OR] per standard deviation increase; 95% confidence interval): accelerometer-measured physical-activity, OR=0.49 (0.33-0.72; p=0.0003); serum iron, OR=0.92 (0.86-0.98; p=0.007); body mass index (BMI), OR=0.90 (0.84-0.97; p=0.003); and mono-unsaturated fat, OR=1.11 (1.02-1.20; p=0.02). Findings in our replication analyses in UK Biobank were compatible with our main analyses (albeit with wide confidence intervals). In MR analysis, height was positively associated with aggressive prostate cancer risk: OR=1.07 (1.01-1.15; p=0.03).

Conclusions: The results for physical-activity, serum iron, BMI, mono-unsaturated fat and height are compatible with causality for prostate cancer. The results suggest that interventions aimed at increasing physical activity may reduce prostate cancer risk.

Keywords: Mendelian randomization, prostate cancer, physical activity

KEY MESSAGES:

- Our MR analyses showed that physical activity, BMI, and serum iron levels were inversely associated with overall prostate cancer risk.
- Mono-unsaturated fat levels were positively associated with overall prostate cancer risk.
- These effects were likely to be causal.
- For these risk factors, the direction of association was consistent for aggressive prostate cancer risk.

- In addition, our MR analyses showed that height was positively associated with aggressive prostate cancer risk.

Introduction

In 2012, 1.1 million men were diagnosed with prostate cancer, making it the second most common male cancer worldwide(1, 2). There is wide global variation in prostate cancer incidence, with almost 70% of cases occurring in more developed regions of the world(2). This variation is thought in part to be related to the intensity of prostate-specific antigen (PSA) based screening practices(3, 4), although migration studies suggest an influence of environmental and lifestyle factors(5, 6). Established risk factors include advanced age, ethnicity and family history of prostate cancer(7, 8). In addition, several lifestyle and anthropometric factors have been hypothesised to play an aetiological role, and measures of adiposity a prognostic role(8). However, the epidemiological evidence to support a causal role for these potentially modifiable factors is weak. This is because inference from observational studies is limited by residual or unmeasured confounding, and other biases such as reverse causation and detection bias(9, 10).

Mendelian randomization (MR) is a methodological approach to addressing reverse causation and confounding within observational studies, based on long-established 'instrumental variable' (IV) principles(11). MR exploits the random assortment of alleles through meiotic cell division at conception, and can be thought of as a natural experiment that generates conditions equivalent to randomised controlled trials, where randomised treatment arms are analogous to randomly assigned genetic subgroups(12-15). At the population level, individuals defined by specific genotypes should on average only differ with respect to that genotype and its phenotypic consequences if certain IV assumptions are met. These assumptions are that the instrument is: i) robustly associated with the exposure it is acting as a proxy for; ii) independent of confounders; and iii) independent of the outcome conditional on the exposure (i.e. 'no pleiotropy' where a single locus influences the outcome through biological pathways that are independent of the exposure of interest)(16). If these assumptions can be shown to have been met, then the genetic polymorphism can be used in an IV framework (i.e. MR) to provide an unconfounded and unbiased estimate of the causal association between the potentially modifiable risk factor and outcome of interest(17). An extension of this methodology - two-sample MR - derives estimates for the required genotype-exposure and genotype-outcome associations from separate and non-overlapping samples of the same representative population(13, 18). Two-sample approaches exploit the rapidly growing availability of summary data from large consortia of genome-wide association studies (GWASs) and thus allow for greater sample sizes to improve statistical power(19).

The aim of this study was to systematically apply two-sample MR analyses to appraise the evidence of a causal link between previously reported lifestyle and anthropometric risk factors with overall and aggressive prostate cancer.

Methods

Selection of risk factors

The World Cancer Research Fund (WCRF) Continuous Update Project (CUP) is a rigorous and systematic synthesis of the global scientific literature on diet, weight and physical-activity in relation to prostate cancer risk, based largely on observational epidemiological studies. We selected all potential risk factors for prostate cancer identified by the WCRF CUP reported in 2018(8).

Defining genetic instruments

We searched for each of the risk factors included in the WCRF report, using exact wording and synonyms, in both the GWAS catalog (www.ebi.ac.uk/gwas) and the MR-Base repository of full GWAS association statistics (www.mrbase.org)(20). This was done to identify any studies that reported associations between single nucleotide polymorphisms (SNPs) and the specific risk factor of interest(20). Further details of how we defined the genetic instruments and their properties are provided in the supplemental material, including; selection criteria for the SNP(s) to proxy each risk factor, how we standardised the beta coefficient and standard error (SE) for each SNP-exposure association, and how we calculated the proportion of variance (R^2) in the risk factor explained by the SNP(s), the strength of the instrument represented by the F-statistic, and the power to detect an odds ratio (OR) of 1.2 (or conversely a protective OR of 0.80).

Outcome trait

GWAS results for prostate cancer were obtained from fixed-effects meta-analyses based on individuals of European ancestry in the PRACTICAL and GAME-ON/ELLIPSE consortia (PRACTICAL: Prostate Cancer Association Group to Investigate Cancer-Associated Alterations in the Genome; GAME-ON: Genetic Associations and Mechanisms in Oncology; ELLIPSE: Elucidating Loci Involved in Prostate Cancer Susceptibility). The summary data were derived from a GWAS of overall prostate cancer on 79,148 cases and 61,106 controls(21), and a GWAS of aggressive prostate cancer involving 15,167 cases and 58,308 controls(21). Aggressive prostate cancer was defined as Gleason score ≥ 8 , PSA >100 ng/mL, metastatic disease (M1), or death from prostate cancer.

Two-sample MR analysis

We used the inverse-variance weighted (IVW) method as our primary MR analytical approach. The IVW method estimates the effect of the exposure on the outcome from the slope of the relationship between β_{XG} (SNP-exposure association) and β_{YG} (SNP-outcome association). This approach performs an inverse variance weighted meta-analysis of each Wald ratio(22), effectively treating each SNP instrumenting a specific risk-factor as a valid natural experiment. We used a random effects IVW model by default, unless there was underdispersion in the causal estimates between SNPs, in which case a fixed effects model was used. The estimates from the random and fixed effects IVW models are the same but the variance for the random effects model is inflated to take into account heterogeneity between SNPs. For risk-factors that only had one SNP available as the instrument, we used the Wald ratio, which is equivalent to β_{YG}/β_{XG} (where Y= outcome, G= gene and X=exposure).

In sensitivity analyses, we applied weighted median(23), weighted mode(24) and MR-Egger regression(25) methods. The weighted median has the advantage that only half the SNPs need to be valid instruments (i.e. exhibiting no horizontal pleiotropy, no association with confounders, and a robust association with the exposure) for the causal effect estimate to be unbiased. The mode-based estimator clusters the SNPs into groups based on similarity of causal effects, and returns the causal effect estimate based on the cluster that has the largest number of SNPs. The weighted mode introduces an extra element similar to IVW and the weighted median, weighting each SNP's contribution to the clustering by the inverse variance of its outcome effect.

The MR-Egger method is similar to the IVW approach but relaxes the 'no horizontal pleiotropy' assumption. MR-Egger regression allows a non-zero intercept in the relationship between multiple SNP-outcome and SNP-exposure associations, where the intercept provides a formal statistical test for the presence of directional (bias inducing) pleiotropy. The slope of the MR-Egger regression between multiple SNP-outcome and SNP-exposure associations can be considered as an unbiased causal effect between the risk factors and prostate cancer, assuming any horizontal pleiotropic effects are not correlated with the SNP-exposure effects (strength of the instrument). The MR pleiotropy residual sum and outlier test (MR-PRESSO) was also implemented to identify outlying genetic variants and analyses were re-run after excluding these variants(26). Violations of the MR 'no horizontal pleiotropy' assumption were also assessed by visual inspection of funnel(27), forest, scatter and leave-one-out plots, and tests of heterogeneity(28) between SNPs making up a multi-allelic instrument(20).

All analyses were conducted using the TwoSampleMR and MRInstruments R packages, curated by MR-Base, www.mrbase.org.

Replication

The risk factors that showed suggestive evidence of association ($p < 0.05$) with overall prostate cancer were assessed for replication among men in the UK Biobank prospective cohort of 7,844 prostate cancer cases and 204,001 controls, using two sample MR. The information on prostate cancer diagnosis was obtained from National Cancer Registries, UK (<http://biobank.ndph.ox.ac.uk/showcase/field.cgi?id=40006>), based on ICD10 code for prostate cancer C61. The GWAS results of this study are available on www.mrbase.org.

Results

There were 57 potential risk factors for prostate cancer considered by the WCRF 2018 report(8)(Supplementary File 1; Table S1 and Table S2). The WCRF reported strong evidence that being obese (BMI, waist circumference and WHR) increases the risk of aggressive prostate cancer (Supplementary File 1; Table S2) and height increases the risk of prostate cancer (Supplementary File 1; Table S1 and S2). There was limited evidence that consumption of dairy products, diets high in calcium, and low plasma alpha-tocopherol and low plasma selenium concentrations increased prostate cancer risk. The evidence was too weak to draw any conclusions for the remaining risk factors. Of these 57 exposures, 22 could be analysed using MR because they had at least one SNP that was strongly associated with them (Table 1).

Mendelian randomization results

Of the 22 potential risk factors examined in our study, only four showed evidence of an association with overall prostate cancer risk (Figure 1, Supplementary File 1; Table S3). Physical-activity, assessed as ‘average acceleration’ (OR per SD change: 0.49; 95% CI: 0.33, 0.72; $P=0.0003$), serum iron levels (OR: 0.92; 95% CI: 0.86, 0.98; $P=0.007$), and BMI (OR: 0.90; 95% CI: 0.84, 0.97; $P=0.003$) were inversely associated with overall prostate cancer risk. Circulating mono-unsaturated fat (OR: 1.11; 95% CI: 1.02, 1.20; $P=0.02$) was positively associated with overall prostate cancer risk. Compared to results from observational studies(8) for overall prostate cancer risk (highest versus lowest total physical-activity; risk ratio (RR): 0.97; 95% CI: 0.9, 1.04), the estimate obtained by MR for the physical-activity measure (average acceleration) was much more strongly supportive of a protective effect (Figure 1). The WCRF reported strong evidence of association for increased body fatness (marked by BMI (RR: 1.08; 95% CI: 1.04, 1.12), waist circumference (RR: 1.12; 95% CI: 1.04, 1.21) and WHR (RR: 1.15; 95% CI: 1.03, 1.28)) with aggressive prostate cancer risk but these results were inconsistent with our MR results, which found evidence of protective association of BMI with overall prostate

cancer risk. In the WCRF report (2018), the observational analysis reported no association between intake of mono-unsaturated fat and overall prostate cancer risk (RR: 1.00; 95% CI: 0.99, 1.01) but our MR-analysis found a positive association between circulating mono-unsaturated fat and overall prostate cancer risk.

None of the risk factors we examined showed strong evidence of association with aggressive prostate cancer although height showed weak evidence of increasing risk; OR: 1.07; 95% CI: 1.01, 1.15; $P=0.03$ (Figure 2, Supplementary File 1; Table S4). The observational studies also reported positive association of height with overall prostate cancer (OR: 1.04.; 95% CI: 1.03, 1.05; $P=1.3 \times 10^{-15}$) and aggressive prostate cancer (8) (OR: 1.04; 95% CI: 1.02, 1.06; $P=6.4 \times 10^{-5}$). However, our MR analysis did not find evidence of association between height and overall prostate cancer risk. There was weak evidence for iron, and average acceleration, with effect-estimates being similar to those observed for overall prostate cancers. In the WCRF report, observational studies have reported positive effects of dairy products, calcium and low selenium concentration on overall prostate cancer, but we did not find strong evidence of associations with these in our MR analyses (Figure 1). However, the confidence intervals (CIs) for these risk factors were overlapping between observational and MR analyses. The power to detect the observationally reported effect size for these risk factors was $>74\%$). Low alpha-tocopherol concentration was positively associated with prostate cancer risk in WCRF report but due to lack of an instrument it was not possible to conduct MR analyses.

There were only two SNPs available for the MR analysis of physical-activity (average acceleration) so we could not perform extensive sensitivity analyses. The direction of association for both SNPs was consistent (Supplementary File 1; Table S5) and the p-value for heterogeneity test was 0.99. These SNPs were on different chromosomes so represent independent associations. After exploring MRBASE-PheWAS (<http://phewas.mrbase.org/>), we found that these two SNPs were associated with anthropometric traits other than physical activity (Supplementary File 1; Table S6). The results for the effect of serum iron, and increasing BMI on overall prostate cancer were consistent across the various sensitivity analyses (Supplementary File 1; Figure S1-S3). The test for directional horizontal pleiotropy by MR-Egger (serum iron: intercept: 0.0005, $P=0.97$ and BMI: intercept: -0.0002, $P=0.89$) didn't find evidence of pleiotropy. Mono-unsaturated fat showed results in the opposite direction using MR-Egger regression (OR: 0.85; 95% CI: 0.55, 1.31; $P=0.51$) compared to other MR-methods (Supplementary File 1; Figure S1 and S4). However, there was no strong evidence for directional pleiotropy for mono-unsaturated fat using MR-Egger (test for directional horizontal pleiotropy by MR-Egger: intercept: 0.04, $P=0.31$). The MR-Egger tests for both iron and mono-unsaturated fat had low power due to the small number of SNPs used, however all 5 SNPs for iron and 4/5 SNPs for mono-unsaturated fats showed

associations with prostate cancer in the same direction (Supplementary File 1; Table S7-S8). At MRBASE-PheWAS, five SNPs of iron were associated with haemoglobin concentration and blood cells count (Supplementary File 1; Table S9) and SNPs of mono-unsaturated fat were associated with lipids (Supplementary File 1; Table S10). The MR results for single SNP analyses of BMI (overall prostate cancer) and height (aggressive prostate cancer) are provided in Supplementary File 1; Table S11-S12.

Replication

The MR analyses were repeated using prostate cancer summary data generated from UK Biobank for physical-activity, iron, BMI, and mono-unsaturated fat (Figure 3). The point-estimates showed consistent directions of association for physical-activity (OR: 0.37; 95% CI: 0.13, 1.06; P=0.07), and BMI (OR: 0.83, 95% CI: 0.74, 0.94; P=0.002). The point-estimates for iron (OR: 1.07; 95% CI: 0.96, 1.20; P=0.24) and mono-unsaturated fat (OR: 0.89; 95% CI: 0.73, 1.07; P=0.20) were in the opposite direction, but the power to detect an effect with these risk factors in UK Biobank was low and confidence intervals for the replication analysis overlapped with our main analysis for all risk factors.

Discussion

We found consistent evidence that physical-activity (assessed as ‘average accelerations’, but not other measures of physical activity), and BMI have an inverse effect on overall prostate cancer risk. There was also evidence of an inverse effect of iron and a positive effect of circulating mono-unsaturated fat in our initial analyses, the effect sizes were in opposite direction in UK Biobank study however power for the replication analysis was low and their CIs overlapped with the PRACTICAL study. There was weak evidence for physical-activity (average accelerations), and iron had a similar effect on aggressive prostate cancer to that seen for overall prostate cancer. We found little evidence that any of the other risk factors studied have a causal role in overall or aggressive prostate cancer, but height showed a positive association with aggressive prostate cancer. The CIs of MR results overlapped with those seen in the observational analyses for all risk factors except for average acceleration and mono-unsaturated fat (overall prostate cancer). In fact, confidence intervals for our MR analysis of aggressive prostate cancer were wide and the power for these analyses was low for many risk factors.

The WCRF report meta-analysed self-reported physical-activity which was assessed in different studies by various methods (i.e. occupational, recreational and total physical-activity) as highest versus lowest level of total physical-activity, a relatively crude dichotomy that may have masked associations. Our MR analysis which proxied fraction accelerations was the most similar to the observational analyses and we did not find evidence of

an association of this measure with prostate cancer. However, we did find an association with average acceleration, which is a different measure and could be high if someone is consistently engaging in light-intensity activity across most of the waking day (vs lots of sitting and a 30 minutes bout of MVPA). Indeed, there is little genetic or phenotypic correlation between the two measures in the UKBiobank population(29). The mechanism for our association between average accelerations and prostate cancer is unclear, although this could be through improved insulin sensitivity or reduced insulin-like growth factor-1 (IGF-1)(30), reduced levels of testosterone and dihydrotestosterone(31), alterations in the antioxidant defence system(31), or improvements in the immune system through enhanced natural killer cell activity(31).

Whilst not consistent with observational analyses, our results for BMI are concordant with other lines of evidence. We have previously shown weak evidence that higher BMI is associated with a reduced prostate cancer risk, in a smaller sample from the PRACTICAL consortium(32). A study examined childhood and adult body size in relation to total incident prostate cancer in a prospective cohort of 47,491 US men(33). High BMI at age 21 was inversely associated with total prostate cancer, with fatal and advanced disease. The association for late adult BMI was more complex and differed by age this could represent confounding by other factors for example physical activity, and diet etc.

Despite showing a protective effect of BMI on prostate cancer risk, we did not find any strong evidence of waist circumference or WHR with risk. However, the results for all these risk factors were in the same direction. The point estimates for BMI, waist circumference and WHR were in the opposite direction to findings from observational results for aggressive prostate cancer. For aggressive prostate cancer, power calculations suggested that we would have good power to detect an effect of BMI (96%) but very low power (33% and 23%) to detect an OR of 1.20 (or, conversely a protective OR of at least 0.80) for waist circumference and WHR respectively. An increased estrogen production has been observed in obese men(34). The sensitivity of prostate cancer to sex hormones has been exploited for therapeutic purposes for many years. Androgen-deprivation therapy is a common treatment in prostate cancer(35), as is the therapeutic use of estrogen for patients with metastatic prostate cancer in both the US(36) and Europe(37), though not popular due to cardiovascular and other side effects. Hence the clinical prediction would be that obesity would be associated with a lower risk of prostate cancer. However, we cannot rule out detection bias(38) arising from delayed diagnosis and therefore a more advanced stage at diagnosis in obese men. This could arise due to lower accuracy of digital rectal examination in obese men or lower PSA values caused by obesity-related traits.

The analysis of iron as a risk factor for prostate cancer found that the evidence was too limited to draw conclusions in the WCRF Second Expert Report and this was not updated in the Third Expert Report due to a lack of new evidence(8, 39). Population studies that have examined the associations between serum iron and cancer outcomes are limited and have reported discordant findings. A prospective cohort study with 15-16 years follow up time reported higher serum iron concentrations increased non-skin cancer risk overall but conversely, in men, higher serum iron concentrations decreased the risk of non-skin cancer(40). A Swedish cohort reported increased serum iron concentrations were not associated with overall cancer risk except for a slightly higher risk of postmenopausal breast cancer(41). Previous findings have suggested that, although higher circulating iron concentrations may potentially increase the risk of cancer in women it may be protective against cancer in men(40) which is in accordance with our MR findings. Although again we were unable to replicate these findings in UK Biobank due to low power.

Our MR-analysis investigated the association of circulating levels of mono-unsaturated fat on prostate cancer, circulating levels of this nutrient have been shown to be poorly correlated with mono-unsaturated fat intake measured by questionnaire(42). The inclusion of an objective measure in our MR-analysis versus questionnaire data in observational studies could be the reason for the discordant results. Although discordance could also be due to negative confounding in the former studies resulting in the effect estimate being closer to the null (e.g., through other dietary, lifestyle, or molecular factors. Instruments for mono-unsaturated fat are correlated with instruments for saturated fat and other lipids. It could be possible that saturated fat or other lipids cause prostate cancer and this result might reflect that. More research is needed into the independence of mono-unsaturated fat of other lipid factors.

Our MR results showed positive association between height and aggressive prostate cancer and the results was consistent with observational studies. Adult height is associated with the rate of growth during fetal life and childhood(43, 44). Health and nutrition status in the neonatal period and childhood may affect the age of sexual maturity. These processes are mediated by changes in the hormonal microenvironment that may have both short and long term effects on circulating levels of growth factors, insulin and other endocrine or tissue specific mediators that may impact cancer risk(45).

The results from replication analyses were compatible with the main findings for BMI, and physical activity (albeit with wide confidence intervals for physical activity), which increases the likelihood that these findings are real. The UK Biobank sample size was however, smaller, and the effects were estimated with less precision

and whilst the results for iron and mono-unsaturated fats did not appear to replicate, due to low power in the replication study we cannot rule out causal effects of these nutrients.

This study's major strength is the use of MR, which is less susceptible to problems of measurement error, confounding and reverse causation in comparison to conventional observational studies. The use of two-sample MR enabled the use of the largest GWAS of prostate cancer(21) to date. We were also able to make use of the largest GWASs on the risk factors of interest, to increase the precision of the SNP-exposure estimates, which should reduce impact of weak instruments bias, which in turn increase statistical power assuming the SNP-exposure estimates are unbiased and risk factor/outcome samples come from the same population.

The study also has some limitations. We had only two SNPs for physical-activity assessed as average acceleration. If there were many independent SNPs available the causal inference could have been strengthened because a) each variant represents an independent natural experiment, and a more precise overall causal estimate (i.e. tighter CIs) can be obtained by meta-analysing the single estimates from each instrument; and b) potential bias arising from the violation of the assumptions can be detected or corrected by evaluating the consistency of effects across instruments(16, 24, 28, 46, 47). For many of the risk factors reported in the WCRF 2018 report, for example alpha-tocopherol, vitamin A, vitamin C etc, we did not find genetic instruments to conduct MR analyses. For the majority of the risk factors in overall prostate cancer, MR analyses were sufficiently powered to detect effect sizes of a modest magnitude (OR of 1.20 or 0.80) except for physical-activity traits (overall acceleration average, fraction of accelerations >425 milli-gravities, and sedentary behaviour), thus failure to detect strong evidence of effects for these risk factors could be due to low power to detect smaller effect sizes. Further identification of independent genetic variants that influence these risk factors will help to improve statistical power for future analyses.

In conclusion, we found evidence that physical-activity, serum iron, and BMI may be causally and inversely related to and circulating mono-unsaturated fat and height may be causally and positively related to, prostate cancer risk. Further studies should investigate the mechanisms by which these factors may lead to prostate cancer and investigate the potential to intervene to reduce risk.

Contributors

SL, RMM and NK conceived and designed the study. NK conducted the analyses. NK wrote the manuscript with input from all authors. Correspondence and material requests should be addressed to SL (s.j.lewis@bristol.ac.uk).


Declaration of interests


None

FUNDING

This work was supported by a grant awarded to SL for 3 years to identify modifiable risk factors for prostate cancer, by the World Cancer Research Fund International (Grant Reference Number: 2015/1421), a Cancer Research UK program grant (C18281/A19169) and the National Institute for Health Research (NIHR) Bristol Biomedical Research Centre. Dr Haycock is supported by CRUK Population Research Postdoctoral Fellowship (C52724/A20138). The views expressed are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care. Lynch was supported by a Mid-Career Research Fellowship from the Victorian Cancer Agency (MCRF18005).

Figure 1. Comparison between observational and MR estimates for the risk factors and overall prostate cancer risk







Only those risk factors are plotted whose observational estimates were reported in WCRF Second or Third Expert Report (8, 39).

The effect estimate for physical activity (Genetic studies) represents average acceleration.

Figure 2. Comparison between observational and MR estimates for the risk factors and aggressive prostate cancer risk




Risk factor



Only those risk factors are plotted whose observational estimates were reported in WCRF Second or Third Exper Report(8, 39).

Figure 3. Comparison of MR estimates (OR) from the main and replication analyses for the risk factors that showed evidence of association ($p < 0.05$) with overall prostate cancer in PRACTICAL



3

4

5

Risk factor	PubMed ID	N	# SNPs ¹	# SNPs ²	Units	SD	R ²	F	Power ¹	Power ²
<i>Anthropometrics and other measures</i>										
Birth weight	27680694	143677	46	45	g	1	1.69	53.54	>99	53
BMI	30124842	681275	535	525	kg/m ²	1	5.66	76.39	>99	96
Height	25282103	253288	433	426	m	1	12.01	79.71	>99	>99
Waist circumference	25673412	232101	45	44	cm	1	0.95	49.60	94	33
WHR	25673412	212244	31	28	ratio	1	0.65	44.95	83	23
<i>Circulating macro- and micro-nutrients</i>										
Sugar/Sucrose (glucose)	22885924	133010	15	15	mmol/l	0.73	0.52	46.65	74	21
Monounsaturated fat	27005778	13535	5	5	mmol/l	1	2.01	55.62	>99	61
Polyunsaturated fat	27005778	13549	19	19	mmol/l	1	13.79	113.93	>99	>99
Total fat	27005778	13505	12	12	mmol/l	1	4.62	54.51	>99	92
Alpha-carotene	28002826	433	3	3	μmol/l	0.23	22.08	40.53	>99	>99
Beta-carotene	19185284	3932	1	1	μmol/l	0.67	2.63	106.36	>99	73
Calcium	24068962	61079	5	5	mg/dl	1	0.65	79.85	83	25
Iron	25352340	72958	5	5	μmol/l	1	2.20	328.85	>99	66
Lycopene	26861389	441	1	1	N/A	N/A	8.34	39.93	>99	>99
Phosphorous	20558539	16264	4	4	mg/dl	0.49	1.52	62.82	99	50
Retinol	21878437	5006	2	2	μg/l	0.22	1.37	34.79	99	46
Selenium	25343990	9639	2	2	μg/L	0.18	1.44	70.33	>99	48
Vitamin D	23393431	42024	4	4	ng/ml	10	2.35	253.15	>99	68
<i>Consumption of foods and drinks</i>										
Alcohol	30643251	941280	77	76	drinks per week	1	0.56	68.35	77	22
Coffee	25288136	91462	4	4	cups/day	1.96	0.45	103.79	68	19
Dairy products (milk intake)	29071499	74241	1	1	glasses/week	1.7	0.52	388.87	74	21
<i>Physical-activity</i>										
Average acceleration	29899525	91084	2	2	milli gravities (mg)	8.14	0.10	44.68	20	7

Sedentary behaviour	30531941	91105	2	2	proportion units	1	0.08	34.91	17	6
Fraction accelerations	29899525	90667	1	1	inverse-normalized fraction of accelerations	1	0.04	39.06	11	5
Sleep duration	30531941	91105	7	7	proportion units	1	0.38	49.29	60	16

6 **Table 1. Details of the instruments used to proxy risk factors for prostate cancer risk**

7 BMI = body mass; WHR = waist-hip ratio; N is the sample size for the GWAS used to define the instruments; #
8 SNPs represents the number of SNPs used within the instrument for each risk factor after clumping,
9 harmonization and extraction of data from a GWAS of prostate cancer (# SNPs¹ for overall prostate cancer risk
10 and # SNPs² for aggressive prostate cancer risk); Units and SD represent the analysis scale and standard
11 deviation scale for betas and SE of SNPs for each risk factor(1 = the GWAS results were already on SD scale
12 otherwise the SD of population mean), respectively; R² represents the variance explained in the risk factor by
13 the instrument; F indicates strength of the instrument used for each risk factor (a strong instrument is sometimes
14 defined as an F-statistic >10); Power¹ represents the power to detect an odds ratio of 1.2 for an association of
15 the risk factor with overall prostate cancer at a significance level (P) of 0.05; Power² represents the power to
16 detect an odds ratio of 1.2 for an association of the risk factor with aggressive prostate cancer at a significance
17 level (P) of 0.05.

18

19 **References**

- 20 1. Stewart B, Wild CP. International Agency for Research on Cancer, World Cancer Report 2014,
21 ed. International Agency for Research on Cancer. 2014.
22 2. Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo M, et al. Cancer incidence and
23 mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. International journal
24 of cancer. 2015;136(5):E359-86.

- 25 3. Center MM, Jemal A, Lortet-Tieulent J, Ward E, Ferlay J, Brawley O, et al. International
26 variation in prostate cancer incidence and mortality rates. *European urology*. 2012;61(6):1079-92.
- 27 4. Wong MC, Goggins WB, Wang HH, Fung FD, Leung C, Wong SY, et al. Global Incidence and
28 Mortality for Prostate Cancer: Analysis of Temporal Patterns and Trends in 36 Countries. *European*
29 *urology*. 2016;70(5):862-74.
- 30 5. Haenszel W, Kurihara M. Studies of Japanese migrants. I. Mortality from cancer and other
31 diseases among Japanese in the United States. *Journal of the National Cancer Institute*.
32 1968;40(1):43-68.
- 33 6. McCredie M, Williams S, Coates M. Cancer mortality in East and Southeast Asian migrants to
34 New South Wales, Australia, 1975-1995. *British journal of cancer*. 1999;79(7-8):1277-82.
- 35 7. Gann PH. Risk factors for prostate cancer. *Reviews in urology*. 2002;4 Suppl 5:S3-s10.
- 36 8. **Diet, nutrition, physical activity and prostate cancer.** World Cancer Research
37 fund/American Institute For Cancer Research Continuous Update Project Expert Report 2018.
- 38 9. Lawlor DA, Davey Smith G, Kundu D, Bruckdorfer KR, Ebrahim S. Those confounded vitamins:
39 what can we learn from the differences between observational versus randomised trial evidence?
40 *Lancet (London, England)*. 2004;363(9422):1724-7.
- 41 10. Davey Smith G, Ebrahim S. Epidemiology--is it time to call it a day? *International journal of*
42 *epidemiology*. 2001;30(1):1-11.
- 43 11. Angrist JD, Imbens GW, Rubin DB. Identification of Causal Effects Using Instrumental
44 Variables. *Journal of the American Statistical Association*. 1996;91(434):444-55.
- 45 12. Davey Smith G ES. "Mendelian randomisation": can genetic epidemiology contribute to
46 understanding environmental determinants of disease? . *Int J Epidemiology* 2003;32:1-22.
- 47 13. Davey Smith G HG. Mendelian randomization: genetic anchors for causal inference in
48 epidemiology studies. *Hum Mol Genet*. 2014.
- 49 14. Lawlor DA HR, Sterne JAC, Timpson NJ, Davey Smith G. Mendelian randomization: using
50 genes as instruments for making causal inferences in epidemiology. *Stat Med*. 2008;27:1133-63.
- 51 15. Haycock PC, Burgess S, Wade KH, Bowden J, Relton C, Davey Smith G. Best (but oft-
52 forgotten) practices: the design, analysis, and interpretation of Mendelian randomization studies.
53 *The American Journal of Clinical Nutrition*. 2016.
- 54 16. Bowden J, Davey Smith G, Haycock PC, Burgess S. Consistent Estimation in Mendelian
55 Randomization with Some Invalid Instruments Using a Weighted Median Estimator. *Genetic*
56 *epidemiology*. 2016;40(4):304-14.
- 57 17. Timpson NJ, Wade KH, Smith GD. Mendelian randomization: application to cardiovascular
58 disease. *Current hypertension reports*. 2012;14(1):29-37.
- 59 18. Pierce BL, Burgess S. Efficient Design for Mendelian Randomization Studies: Subsample and
60 2-Sample Instrumental Variable Estimators. *American Journal of Epidemiology*. 2013;178(7):1177-84.
- 61 19. Yarmolinsky J, Wade KH, Richmond RC, Langdon RJ, Bull CJ, Tilling KM, et al. Causal Inference
62 in Cancer Epidemiology: What Is the Role of Mendelian Randomization? *Cancer epidemiology,*
63 *biomarkers & prevention* : a publication of the American Association for Cancer Research,
64 cosponsored by the American Society of Preventive Oncology. 2018;27(9):995-1010.
- 65 20. Hemani G, Zheng J, Elsworth B, Wade KH, Haberland V, Baird D, et al. The MR-Base platform
66 supports systematic causal inference across the human phenome. *eLife*. 2018;7.

- 67 21. Schumacher FR, Al Olama AA, Berndt SI, Benlloch S, Ahmed M, Saunders EJ, et al. Association
68 analyses of more than 140,000 men identify 63 new prostate cancer susceptibility loci. *Nature*
69 *genetics*. 2018;50(7):928-36.
- 70 22. Johnson T. Efficient calculation for Multi-SNP genetic risk scores. American Society of Human
71 Genetics Annual Meeting. 2012.
- 72 23. Bowden J, Davey Smith G, Haycock PC, Burgess S. Consistent Estimation in Mendelian
73 Randomization with Some Invalid Instruments Using a Weighted Median Estimator. *Genetic*
74 *Epidemiology*. 2016;40(4):304-14.
- 75 24. Hartwig FP, Davey Smith G, Bowden J. Robust inference in summary data Mendelian
76 randomization via the zero modal pleiotropy assumption. *International journal of epidemiology*.
77 2017;46(6):1985-98.
- 78 25. Bowden J, Davey Smith G, Burgess S. Mendelian randomization with invalid instruments:
79 effect estimation and bias detection through Egger regression. *International Journal of*
80 *Epidemiology*. 2015.
- 81 26. Verbanck M, Chen CY, Neale B, Do R. Detection of widespread horizontal pleiotropy in causal
82 relationships inferred from Mendelian randomization between complex traits and diseases. *Nature*
83 *genetics*. 2018;50(5):693-8.
- 84 27. Sterne JA, Sutton AJ, Ioannidis JP, Terrin N, Jones DR, Lau J, et al. Recommendations for
85 examining and interpreting funnel plot asymmetry in meta-analyses of randomised controlled trials.
86 *BMJ (Clinical research ed)*. 2011;343:d4002.
- 87 28. Bowden J, Del Greco MF, Minelli C, Davey Smith G, Sheehan N, Thompson J. A framework for
88 the investigation of pleiotropy in two-sample summary data Mendelian randomization. *Statistics in*
89 *medicine*. 2017;36(11):1783-802.
- 90 29. Klimentidis YC, Raichlen DA, Bea J, Garcia DO, Wineinger NE, Mandarino LJ, et al. Genome-
91 wide association study of habitual physical activity in over 377,000 UK Biobank participants identifies
92 multiple variants including CADM2 and APOE. *International journal of obesity (2005)*.
93 2018;42(6):1161-76.
- 94 30. Ngo TH, Barnard RJ, Tymchuk CN, Cohen P, Aronson WJ. Effect of diet and exercise on serum
95 insulin, IGF-I, and IGFBP-1 levels and growth of LNCaP cells in vitro (United States). *Cancer causes &*
96 *control : CCC*. 2002;13(10):929-35.
- 97 31. Friedenreich CM, Thune I. A review of physical activity and prostate cancer risk. *Cancer*
98 *causes & control : CCC*. 2001;12(5):461-75.
- 99 32. Davies NM, Gaunt TR, Lewis SJ, Holly J, Donovan JL, Hamdy FC, et al. The effects of height
100 and BMI on prostate cancer incidence and mortality: a Mendelian randomization study in 20,848
101 cases and 20,214 controls from the PRACTICAL consortium. *Cancer causes & control : CCC*.
102 2015;26(11):1603-16.
- 103 33. Moller E, Wilson KM, Batista JL, Mucci LA, Balter K, Giovannucci E. Body size across the life
104 course and prostate cancer in the Health Professionals Follow-up Study. *International journal of*
105 *cancer*. 2016;138(4):853-65.
- 106 34. Schneider G, Kirschner MA, Berkowitz R, Ertel NH. Increased estrogen production in obese
107 men. *The Journal of clinical endocrinology and metabolism*. 1979;48(4):633-8.
- 108 35. Wong YN, Ferraldeschi R, Attard G, de Bono J. Evolution of androgen receptor targeted
109 therapy for advanced prostate cancer. *Nature reviews Clinical oncology*. 2014;11(6):365-76.

- 110 36. Mohler JL, Kantoff PW, Armstrong AJ, Bahnson RR, Cohen M, D'Amico AV, et al. Prostate
111 cancer, version 2.2014. *Journal of the National Comprehensive Cancer Network* : JNCCN.
112 2014;12(5):686-718.
- 113 37. Heidenreich A, Bastian PJ, Bellmunt J, Bolla M, Joniau S, van der Kwast T, et al. EAU
114 guidelines on prostate cancer. Part II: Treatment of advanced, relapsing, and castration-resistant
115 prostate cancer. *European urology*. 2014;65(2):467-79.
- 116 38. Fowke JH, Motley S, Dai Q, Concepcion R, Barocas DA. Association between biomarkers of
117 obesity and risk of high-grade prostatic intraepithelial neoplasia and prostate cancer--evidence of
118 effect modification by prostate size. *Cancer letters*. 2013;328(2):345-52.
- 119 39. World Cancer Research Fund/ American Institute of Cancer Research. *Food, Nutrition,
120 Physical Activity, and the Prevention of Cancer: a Global Perspective*. Washington DC: AICR, 2007
121 (Second Expert Report). 2007.
- 122 40. Chua AC, Knuiman MW, Trinder D, Divitini ML, Olynyk JK. Higher concentrations of serum
123 iron and transferrin saturation but not serum ferritin are associated with cancer outcomes. *The
124 American journal of clinical nutrition*. 2016;104(3):736-42.
- 125 41. Gaur A, Collins H, Wulaningsih W, Holmberg L, Garmo H, Hammar N, et al. Iron metabolism
126 and risk of cancer in the Swedish AMORIS study. *Cancer causes & control : CCC*. 2013;24(7):1393-
127 402.
- 128 42. Marchioni DM, de Oliveira MF, Carioca AAF, Miranda AAM, Carvalho AM, Oki E, et al. Plasma
129 fatty acids: Biomarkers of dietary intake? *Nutrition (Burbank, Los Angeles County, Calif)*. 2019;59:77-
130 82.
- 131 43. Barker DJ, Thornburg KL. Placental programming of chronic diseases, cancer and lifespan: a
132 review. *Placenta*. 2013;34(10):841-5.
- 133 44. Rolland-Cachera MF. Rate of growth in early life: a predictor of later health? *Advances in
134 experimental medicine and biology*. 2005;569:35-9.
- 135 45. Le Roith D, Bondy C, Yakar S, Liu JL, Butler A. The somatomedin hypothesis: 2001. *Endocrine
136 reviews*. 2001;22(1):53-74.
- 137 46. Bowden J, Davey Smith G, Burgess S. Mendelian randomization with invalid instruments:
138 effect estimation and bias detection through Egger regression. *International journal of epidemiology*.
139 2015;44(2):512-25.
- 140 47. Hemani G, Tilling K, Davey Smith G. Orienting the causal relationship between imprecisely
141 measured traits using GWAS summary data. *PLoS genetics*. 2017;13(11):e1007081.

142

143