



Rachael, F., Apelqvist, J., Boyko, E. J., Fitridge, R., Hong, J. P., Katsanos, K., Mills, J. L., Nikol, S., Reekers, J., Venermo, M., Zierler, R. E., Hinchliffe, R. J., & Schaper, N. C. (2020). Effectiveness of revascularization of the ulcerated foot in patients with diabetes and peripheral artery disease: a systematic review. *Diabetes/Metabolism Research and Reviews*, 36(S1), Article e3279. Advance online publication. <https://doi.org/10.1002/dmrr.3279>

Peer reviewed version

Link to published version (if available):  
[10.1002/dmrr.3279](https://doi.org/10.1002/dmrr.3279)

[Link to publication record on the Bristol Research Portal](#)  
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Wiley at <https://doi.org/10.1002/dmrr.3279> . Please refer to any applicable terms of use of the publisher.

## University of Bristol – Bristol Research Portal

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/brp-terms/>

## **Effectiveness of revascularisation of the ulcerated foot in patients with diabetes and peripheral artery disease: a systematic review**

**Authors:** Rachael O. Forsythe<sup>1</sup>; Jan Apelqvist<sup>2</sup>; Ed J. Boyko<sup>3</sup>; Robert Fitzridge<sup>4</sup>; Joon Pio Hong<sup>5</sup>; Konstantinos Katsanos<sup>6</sup>; Joseph L. Mills<sup>7</sup>; Sigrid Nikol<sup>8</sup>; Jim Reekers<sup>9</sup>; Maarit Venermo<sup>10</sup>; R. Eugene Zierler<sup>11</sup>; Robert J. Hinchliffe<sup>12</sup>; Nicolaas C. Schaper<sup>13</sup>; on behalf of the International Working Group on the Diabetic Foot (IWGDF).

### **Institutions:**

<sup>1</sup>British Heart Foundation / University of Edinburgh Centre for Cardiovascular Science, University of Edinburgh, Edinburgh, Scotland, UK

<sup>2</sup>Department of Endocrinology, University Hospital of Malmö, Sweden

<sup>3</sup>Seattle Epidemiologic Research and Information Centre-Department of Veterans Affairs Puget Sound Health Care System and the University of Washington, Seattle, WA, USA.

<sup>4</sup>Vascular Surgery, The University of Adelaide, Adelaide, South Australia, Australia

<sup>5</sup>Asan Medical Center University of Ulsan, Seoul, Korea

<sup>6</sup>Patras University Hospital School of Medicine, Rion, Patras, Greece

<sup>7</sup>SALSA (Southern Arizona Limb Salvage Alliance), University of Arizona Health Sciences Center, Tucson, Arizona, USA

<sup>8</sup>Asklepios Klinik St. Georg, Hamburg, Germany

<sup>9</sup>Department of Vascular Radiology, Amsterdam Medical Centre, The Netherlands

<sup>10</sup>Helsinki University Hospital, University of Helsinki, Finland

<sup>11</sup>Department of Surgery, University of Washington, Seattle, Washington, USA

<sup>12</sup>Bristol Centre for Surgical Research, University of Bristol, Bristol, UK.

<sup>13</sup>Div. Endocrinology, MUMC+, CARIM and CAPHRI Institute, Maastricht, The Netherlands

Corresponding author: Rachael Forsythe; rachael.forsythe@ed.ac.uk

## **ABSTRACT**

In patients with diabetes, foot ulceration and peripheral artery disease (PAD), it is often difficult to determine whether, when and how to revascularise the affected lower extremity. The presence of PAD is a major risk factor for non-healing and yet clinical outcomes of revascularisation are not necessarily related to technical success. The International Working Group of the Diabetic Foot updated systematic review on the effectiveness of revascularisation of the ulcerated foot in patients with diabetes and PAD is comprised of 64 studies describing >13,000 patients. Amongst 60 case series and 4 non-randomised controlled studies, we summarised clinically relevant outcomes and found them to be broadly similar between patients treated with open versus endovascular therapy. Following endovascular revascularisation, the 1 year and 2 year limb salvage rates were 80% (IQR 78-82%) and 78% (IQR 75-83%), whereas open therapy was associated with rates of 85% (IQR 80-90%) at 1 year and 87% (IQR 85-88%) at 2 years, however these results were based on a varying combination of studies and cannot therefore be interpreted as cumulative. Overall, wound healing was achieved in a median of 60% of patients (IQR 50-69%) at 1 year in those treated by endovascular or surgical therapy, and the major amputation rate of endovascular versus open therapy was 2% vs 5% at 30 days, 10% vs 9% at 1 year and 13% vs 9% at 2 years. For both strategies, overall mortality was found to be high, with 2% (1-6%) peri-operative (or 30 day) mortality, rising sharply to 13% (9-23%) at 1 year, 29% (19-48%) at 2 years and 47% (39-71%) at 5 years. Both the angiosome concept (revascularisation directly to the area of tissue loss via its main feeding artery) or indirect revascularisation through collaterals, appear to be equally effective strategies for restoring perfusion.

Overall, the available data do not allow us to recommend one method of revascularisation over the other and more studies are required to determine the best revascularisation approach in diabetic foot ulceration.

**Keywords:** peripheral artery disease, diabetes, diabetic foot, foot ulcer, revascularisation, endovascular treatment, vascular surgery, mortality, amputation

## **INTRODUCTION**

Peripheral artery disease (PAD) affects up to 50% of patients with a diabetic foot ulcer (DFU) <sup>1 2 3</sup> and its presence is associated with poor outcomes. The natural history of patients with PAD and DFU is difficult to predict and is affected by more than just the severity of arterial disease. Concurrent infection increases the risk of poor outcomes even more, and other factors such as medical co-morbidities, microvascular dysfunction, poor glycaemic control and abnormal mechanical load may also contribute. Whilst it is important to recognise and address all of these clinical abnormalities, there is evidence that early revascularisation in patients with PAD is associated with improved outcomes <sup>4</sup>. However, the decision whether, when and how to revascularise is not straightforward. Up to 50% of patients for whom revascularisation is not thought technically possible or reasonable due to co-morbidities may heal their ulcer within a year without revascularisation <sup>5</sup>; conversely, amongst those who had a successful and patent revascularisation, >20% of patients underwent a major limb amputation within 12 months in one large series <sup>6</sup>.

The anatomical distribution of PAD is more challenging in patients with, as opposed to without, diabetes. There is a predilection for multi-level distal disease often involving multiple crural arteries, with long occlusions, poor propensity to form collaterals and a high prevalence of medial arterial calcification. All of these factors pose additional technical challenges when attempting revascularisation, either by open surgical or endovascular means. In addition, there are no major randomised trials addressing the most appropriate methods of revascularisation specifically for patients with DFU and PAD, nor are there informative subgroup analyses in general PAD trials.

The International Working Group on the Diabetic Foot (IWGDF) is a multi-disciplinary group of experts in the management of patients with DFU. The present systematic review of the effectiveness of revascularisation is an updated iteration of our previous review, launched in 2015 <sup>7</sup>, and informs the most recent IWGDF guidance addressing the diagnosis, prognosis and management of patients with PAD and DFU, which is published separately in this journal <sup>8</sup>.

## **METHODS**

### **Search methods**

We updated our previous systematic review <sup>7</sup>, guided by a recent consensus document on updating systematic reviews <sup>9</sup>. We searched the MEDLINE and EMBASE databases for studies relating to therapies to revascularise the ulcerated foot amongst patients with diabetes, updating the previous search and therefore capturing any new records published between June 2014 and September 2018. The search string is shown in Appendix A. Two reviewers independently screened the abstracts of retrieved articles to determine if they might meet the pre-set inclusion criteria, and a third reviewer adjudicated any conflicts. We accessed full-text articles of screened articles and assessed them for inclusion; members of the IWGDF PAD working group then extracted and verified data.

### **Inclusion / exclusion criteria**

To be eligible for inclusion, a study was required to meet the following criteria: 1) it reported on the outcomes of revascularisation (open or endovascular) for patients with DFU and PAD; 2) patients included had objective evidence of PAD (such as angiography); 3) >80% of patients included had tissue loss (defined as any lesion of the skin breaching the epithelium or ulceration or gangrene); 4) it included at least 40 patients with >80% of the population diagnosed with diabetes or where the results of at least 30 patients with diabetes were reported separately; and 5) it reported on primary outcomes including ulcer healing, limb salvage, major amputation or survival.

Studies reporting only on aorto-iliac disease were excluded, as the treatment of supra-inguinal disease is similar in patients with or without diabetes. Studies were excluded if they reported only on medical, pharmacological or topical therapies or if they compared different revascularisation technologies.

We only included studies published in the English language.

### **Primary outcome measures and definitions**

The primary outcome measures of interest included wound healing, limb salvage, major amputation and survival.

For the purpose of this systematic review, *PAD* was defined as any flow limiting atherosclerotic lesion of the arteries below the inguinal ligament. We accepted the diagnosis of *diabetes mellitus* as it was made according to the individual publication. *Tissue loss* was defined as any lesions of the skin breaching the epithelium, or the presence of ulceration or gangrene. *Early* mortality was considered within 30 days or within the period of the first hospital admission. A *major complication* was defined as any that resulted in a systemic disturbance of the patient or prolonged hospitalisation (or as defined by the individual study).

### **Data extraction and quality assessment**

This systematic review was performed according to Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) guidelines <sup>10</sup>. Two reviewers assessed studies for inclusion based on titles, then a review of the abstract, and finally upon review of the full text . The data for the evidence table were then extracted by members of the IWGDF PAD working group, confirmed by other members. Studies were assessed for methodological rigour using the Scottish Intercollegiate Guidelines Network (SIGN) guidelines <sup>11</sup>. Pooling of data and weighting of studies was not possible because of study heterogeneity and the generally low quality of evidence. Data was summarised for individual items as interquartile range (IQR) and median (not weighted).

Patient demographics were summarised, along with specific details of the foot lesion where available, and any reported concurrent infection. Objective assessments of perfusion were reported where available. We made no distinction amongst various endovascular techniques (e.g. angioplasty, stenting, subintimal angioplasty, atherectomy), which were all referred to as endovascular therapy. Neither did we distinguish among methods of open revascularisation (e.g. *in situ* versus reversed vein bypass graft).

## **RESULTS**

### **Search results**

Our updated search included articles published between June 2014 and June 2018 (Figure 1). A total of 8759 articles were screened, of which 295 articles were

assessed for inclusion based on the abstract and 80 assessed for inclusion based on the full text. Ultimately, 7 new studies deemed eligible were identified and, when added to those from our previous review <sup>7</sup>, a total of 64 studies, comprising 13,434 patients, are included in this updated systematic review.. There were no randomised controlled trials, all included studies were case series (SIGN 3) that reported on bypass surgery, endovascular intervention or both techniques used in combination. Few studies provided sufficient detail on severity of PAD or baseline foot lesion characteristics, however most studies provided adequate information on patient demographics and co-morbidities. In the event that more than one study reported on patients from the same institution, we highlighted this in the evidence table, but accepted that it is likely that some patients were reported more than once. A complete evidence table with all results from included studies can be found in Appendix B.

**PICO (patient, intervention, comparison, outcome):** What are the aims and methods of revascularisation and onward management in a person with diabetes, foot ulceration and PAD?

### **Summary of the literature**

#### **Patient demographics and comorbidities**

Among the 13,434 patients were included, of which 69% (IQRs 68-71%) were men, with a median age of 71 (69-72). As expected, there was a high prevalence of cardiovascular comorbidities, including 47% (43-55%) with coronary artery disease, 21% (18-23%) with cerebrovascular disease and 21% (20-26%) with end-stage renal disease (variably defined).

#### **Clinical outcomes**

It is important to note that the quantitative interpretation of outcomes cannot be considered cumulative, as there was a discrepancy of outcome reporting between studies and studies had different duration of follow-up. The results reported for each time point are based on a variable number of studies and should not be directly compared.

## **Wound healing**

Only 12 studies reported wound healing as an outcome measure, using variable follow-up periods. Wound healing was achieved in a median of 60% (50-69%) at 1 year. Of 3 studies reporting separately on endovascular outcomes, wound healing at 1 year was 75% (68-77%), while for two studies reporting on open therapy, the median wound healing at 1 year was lower (52%; 46-57%).

## **Limb salvage and major amputation**

Limb salvage was reported at a predefined time point by 39 studies. The overall limb salvage rates were 82% (IQR 79-89%) at 1 year (based on 28 studies), 86% (77-88%) at 2 years (based on 12 studies) and 76% (73-78%) at 5 years (based on 14 studies). Following endovascular treatment, 1 and 2 year limb salvage rates were 80% (78-82%; based on 9 studies) and 78% (75-83%; based on 5 studies), respectively. In those studies reporting limb salvage following open revascularisation (n=25), the rates were: 85% (80-90%) at 1 year (based on 19 studies) and 87% (86-89%) at 2 years (based on 7 studies).

The definition of major amputation was variable amongst the included studies but, where reported with a specified time point, the median rate of major amputations within 30 days was 4% (2-5%), which increased at 1 year to 9% (4-12%) and 11% (7-18%) at 2 years. The early (30 day) major amputation rate appeared somewhat lower for endovascular therapy when compared to open therapy (2% vs 5%) but higher at 1 year (10% vs 9%) and 2 years (13% vs 9%). However, this finding must be interpreted with great caution, due to the heterogeneity of studies and patients included and the inconsistency in reporting time points between studies.

The rates of minor amputation varied widely, with a median of 38% (IQR 23-59%). There were similar rates of minor amputation between patients undergoing open (36%, 23-57%) and endovascular (38%, 23-57%) therapies.

## **Mortality and survival**



Peri-operative or 30 day mortality was reported in 30 studies, and was 2% (IQR 1-5%) overall. Peri-operative mortality the same following endovascular versus open revascularisation (2% vs 2%). Overall 1 year mortality was 13% (9-23%), rising to 29% (19-48%) at 2 years and 47% (39-71%) at 5 years. At 2 years, the highest mortality rates were amongst patients with ESRD (72% in one study<sup>15</sup>). At 5 years, amongst those in seven studies who underwent open therapy, the mortality rate was 43% (39-60%). In the single study that reported 5 year mortality following endovascular therapy it was 74%<sup>16</sup>, however very small numbers remained in follow up at 5 years and the validity of these findings is therefore questionable.

The five studies reporting survival as compared to mortality yielded poorer results, with a median survival of 83% (75-87%) at 1 year (four studies) and 39% (26-46%) at 5 years (four studies), which translates into mortalities of 17% and 62% at 1 and 5 years, respectively.

### **Specific revascularisation strategies**

#### **Pedal bypass**

Among the twelve studies that reported on pedal bypass surgery, the median 1, 2 and 5 year limb salvage rates were 86% (80-92%), 85% (68-86) and 87% (78-95%). Peri-operative mortality was similar to that in the overall population at 2% (1-4%), with 1 year, 2 year and 5 year mortality rates of 14% (11-32%), 48% (32-60%) and 42% (39-45%), respectively. These outcomes are again based on a variable number of studies, with different duration of follow-up, reporting at each time point.

#### **Infra-popliteal angioplasty**

Eight studies reported exclusively on infra-popliteal angioplasty; at one year the median rate of wound healing was 71% (65-75%) and limb salvage was 77% (71-85%).

In a review of 448 patients (83% with diabetes) who underwent infrapopliteal angioplasty for tissue loss (86%) or rest pain (14%), there was no significant difference in major amputation rates at one year between single-vessel angioplasty and multiple-vessel intervention (16% vs 10%,  $p=0.24$ )<sup>17</sup>. Another

smaller study (n=92) <sup>18</sup>, also comparing single versus multiple endovascular revascularisation attempts, found no difference in outcomes between the two groups. In a third study of 92 patients undergoing endovascular therapy, the presence of a complete pedal arch following the intervention was associated with increased rates of wound healing, limb salvage and survival at 1 year compared to absence of a complete pedal arch <sup>19</sup>.

### **Angiosome-directed revascularisation**

Angiosome-directed revascularisation is a method of improving perfusion of an area of tissue loss directly, via its feeding artery (also called direct revascularisation, DR). This is based on the theory that the foot can be divided into 3 dimensional units of tissue (angiosomes), each of which is supplied by a specific feeding artery. By targeting a feeding artery that directly supplies an area of tissue loss, the concept is that this is the best way to improve perfusion to that specific area. The more traditional 'best vessel approach' can be considered non-angiosome directed or indirect revascularisation (IR), whereby the most suitable target vessel is chosen for revascularisation, regardless of its anatomical location. The majority of literature relating to outcomes based on the angiosome concept in patients with diabetes and foot ulcers is retrospective. We identified eight such studies, all of which were retrospective case series or cohort studies and had a high risk of bias. Six studies concluded that direct revascularisation according to the angiosome concept was associated with higher rates of wound healing when compared with indirect revascularisation <sup>20 18 21 22 23 24</sup>, however only four of these studies reported higher rates of limb salvage.

In one study <sup>18</sup>, patients who underwent indirect revascularisation (ie non-angiosome directed) were stratified further by the role of collaterals – those in whom pulsatile flow was restored to the affected area indirectly (ie not targeted to the feeding artery) but by collaterals (indirect revascularisation, via collaterals; IRc) and those in whom there were no collaterals between the target artery and the affected angiosome (indirect revascularisation, no collaterals). In this series, 92 patients underwent endovascular infra-popliteal intervention (including those with concurrent supra-popliteal angioplasty), and the outcomes were similar for DR and IRc – limb salvage rates were 89% and 85% at 2 years and wound healing

was 66% and 68% at 12 months. However, in those patients who underwent indirect revascularisation with no collaterals, limb salvage and wound healing rates were poor (59% and 7%, respectively).

A more recent study compared angiosome-based intervention during endovascular and open techniques in a series of 545 patients with diabetes and foot ulceration undergoing a first-time infra-popliteal revascularisation <sup>23</sup>. The highest rates of wound healing were in patients who underwent open revascularisation classified as DR (77% healing at 1 year) and the worst rates were in those who underwent endovascular treatment classified as IR (52% healing at 1 year). Amputation rates were also highest in the IR endovascular group, but if IR was achieved via collaterals then this improved limb salvage.

### **Significant clinical comorbidities**

#### **End-stage renal disease**

In the nine studies that analysed patients with end-stage renal disease (ESRD, defined variably) separately, its presence increased peri-operative mortality (4.6%, 2.6-8.8%) and high mortality over 5 years (48% and 72% at 2 years <sup>25 26</sup>, 56% at 3 years <sup>27</sup> and 91% at 5 years <sup>12</sup>), albeit based on small study numbers. For those who did survive to 1 year, salvage rates were 70% (65-70%).

#### **Infection**

Whilst it is a well recognised risk factor for poor outcomes in patients with diabetes and foot ulceration, only two studies specifically reported on outcomes in patients with diabetes, PAD and foot infection <sup>28 29</sup>. Both studies were retrospective case series of patients with an acutely infected ulcer. In a study of 53 patients who underwent pedal bypass <sup>28</sup>, 20% required minor amputation prior to bypass; however, the resulting limb salvage was excellent, at 98% at 1 year, 98% at 2 years and 95% at 3 years. In a study of 114 patients with foot infection, 38% underwent open revascularisation and the overall limb salvage for the case series was 87% at 2 years and 73% at 4 years <sup>29</sup>

**Evidence statement:** Evidence is inadequate to establish whether an endovascular, open or hybrid revascularisation technique is superior in restoring blood flow and improving prognosis in patients with diabetic foot ulceration and PAD

**Quality of the evidence:** Low, based on cohort studies and case series

**Evidence statement:** Restoration of direct blood flow to at least one of the foot arteries, preferably the artery that supplies the anatomical region of the ulcer is associated with the best outcome. Both revascularisation directly to the area of tissue loss via its main feeding artery or indirect revascularisation through collaterals, appear to be equally effective strategies for restoring perfusion

**Quality of evidence:** Low, based on cohort studies and case series

## **Discussion**

This systematic review is an update of the version launched in 2015 <sup>30</sup> and, in addition to the 56 studies included previously, includes a further seven studies describing outcomes of revascularisation in patients with diabetes, PAD and foot ulceration. All of the new studies were retrospective reviews of medical care encounter databases in either single or multiple institutions across the world and had a high risk of bias, as well as significant heterogeneity in included participants as well as outcome reporting. Of particular note, none of these new studies included a control group. There was also inconsistency in reporting of some key variables such as the severity of arterial perfusion deficit, indication for revascularisation and ulcer characteristics. For these reasons, it was not possible to pool the data or conduct a meta-analysis.

One of the main obstacles in revascularisation of patients with diabetes is the often complex, distal pattern of disease, which poses a challenge for performing both endovascular and open therapy. Crural vessel angioplasty and distal bypasses are often more time-consuming and technically demanding than revascularisation

with a more proximal outflow target vessel, and thus require additional expertise. The presence of multi-level disease, highly calcified lesions and paucity of collaterals each further contributes to this challenge. Overall, however, the current rates of reported limb salvage presented here (82% at 1 year and up to 78% at 5 years) are greatly improved on those reported by two studies documenting the natural history of patients with diabetes and foot ulcers for whom revascularisation was deemed unsuitable (50-54% at 1 year) <sup>31 5</sup>. Even amongst patients included in this review who underwent pedal bypass or crural angioplasty, limb salvage rates were reported as 86% and 77% at 1 year.

Because there are no randomised trials comparing methods of revascularisation specific to patients with diabetic foot disease, it is not possible to determine which method of revascularisation is more effective. Results from the Bypass versus Angioplasty in Severe Ischaemia of the Leg (BASIL) trial, the only completed RCT comparing open or endovascular treatment of severe limb ischaemia, published in 2005 <sup>32</sup>, cannot be directly extrapolated to patients with diabetes and foot ulceration, as only around 40% of the patients in BASIL had diabetes and the main pattern of disease was femoropopliteal. Moreover, there has been substantial advancement of endovascular technologies that were not captured by the BASIL trial. A more recent sub-group analysis of the BASIL trial that included only those enrolled patients with infra-popliteal disease demonstrated no difference in outcomes in open versus endovascular intervention <sup>33</sup>; however since <50% of these patients had diabetes, this information is again not directly transferable to the problem of the best revascularisation method for the diabetic foot. At the time of writing, BASIL-2 is recruiting patients with severe limb ischaemia due to infra-popliteal disease (with or without the presence of femoropopliteal disease) and will compare vein bypass-first versus best endovascular-first revascularisation. Given that a higher proportion of patients in BASIL-2 are likely to have diabetes, this study may facilitate further sub-group analyses that may be more informative for patients with diabetic foot ulcers and PAD.

Nevertheless, the outcomes presented in this review are broadly similar for open and endovascular therapy, albeit in very heterogeneous populations. Wound

healing was infrequently reported as an outcome (and variably defined) and whilst it appeared that endovascular compared to open revascularisation was associated with improved ulcer healing at 1 year (75% vs 52%), one of the two studies reporting open therapy included patients in whom almost 75% had ultra distal bypass, which may skew the overall outcomes due to its technical complexity. Otherwise, the rates of limb salvage and major amputation were broadly similar between patients undergoing open versus endovascular treatment. The results of both of these methods, however, will of course depend on the expertise and resources at a given institution. Our review does not summarise the data pertaining to technical success or feasibility of revascularisation in patients with diabetes, the most important outcomes to report are in our opinion clinical. Whilst the aim should be to use a durable revascularisation approach, the overall goal of healing the foot may be met even if a revascularisation site does not remain patent in the long term.

The angiosome concept is an area of much debate. Traditionally, the best vessel is chosen as the target for revascularisation, with a goal of restoring inline pulsatile blood flow to the foot through any means. More recently described, angiosome-directed revascularisation is based on the principle that the foot can be divided into six 3-dimensional blocks of tissue, each supplied by a feeding artery. The theory is that by identifying the specific feeding artery to an area of tissue loss and targeting that artery for revascularisation (ie direct revascularisation; DR), restoration of pulsatile blood flow directly to an area of tissue ischaemia renders it more likely to heal. Alternatively, non-angiosome directed therapy (indirect revascularisation; IR) adopts the 'best vessel' approach, whereby the most suitable target artery is chosen, regardless of whether it relates to the area of tissue loss and blood flow is therefore restored to the area by collaterals. Given that patients with diabetes typically have poor collaterals, it seems intuitively that angiosome-directed revascularisation might be more effective.

The results of six out of eight retrospective studies in this review suggest that direct revascularisation (ie angiosome-based therapy) is associated with improved wound healing and in four studies, this translated to improved limb

salvage. Those of one study<sup>18</sup> suggest that indirect endovascular revascularisation through collaterals has similar outcomes to direct revascularisation and that both offer significantly improved rates of limb salvage and wound healing when compared to indirect revascularisation without collaterals. These findings are reflected in two recent meta-analyses that combined include >4000 limbs with foot ulcers (>80% of which had diabetes), which both concluded that endovascular IR significantly improves wound healing and major amputation rates, but that, in the presence of collaterals, IR and DR have similar outcomes<sup>34 35</sup>. However, the populations included were highly variable, the definitions poorly defined and it is not possible to draw any firm conclusions from these data. Despite this, we think it would be sensible to pursue an angiosome-based revascularisation strategy if possible, particularly in patients with diabetes who have poor collaterals and may therefore benefit from the restoration of blood flow directly to the area of tissue loss.

That said, since several patients, >20% of patients in one study, who undergo a successful revascularisation procedure will still require a major amputation within 12 months<sup>6</sup>, there is clearly more to the story than optimising perfusion. While other clinical variables that affect outcomes (such as infection, wound characteristics, neuropathy, and comorbidities), it is difficult to know what proportion each of these factors contributes to an individual patient's chance of clinical success. The evidence indicates that patients with diabetes, ulceration and severe perfusion deficit should be considered for early revascularisation, however the role for intervention in those with mild to moderate ischaemia is less well defined. The studies included in our review provided few data about the perfusion deficit, the duration of attempted conservative management or the indication for vascular investigation and treatment. In addition, there is no known threshold value of perfusion towards which we should aim when attempting revascularisation. So as part of a goal to optimise all approaches to heal a diabetic foot ulcer, these patients mild to moderate ischaemia should be investigated further for the presence of a perfusion defect amenable to revascularisation if there is no significant improvement in healing within 4-6 weeks of optimal care.

Unfortunately, there are some clinical characteristics that contribute significantly to poor outcomes but cannot be optimised. ESRD is a known risk factor for foot ulceration and major amputation in patients with diabetes <sup>36</sup> and outcomes of revascularisation in this group are typically poor. Our review found that patients with ESRD undergoing revascularisation had lower 1 year limb salvage rates (70% vs 82%). Moreover, there was increased peri-operative mortality, and >50% of patients were dead at 3 years and up to 90% at 5 years. This appears much higher than the overall median mortality rates of almost 30% at 2 years and 46% at 5 years.

The poor survival rates in patients with diabetes, PAD and foot ulceration is in a large part attributable to the systemic nature of arterial disease. Many patients with PAD will also have ischaemic heart disease and cerebrovascular disease on presentation. It is imperative, therefore, to ensure that all patients with PAD receive comprehensive medical management of concomitant cardiovascular disease and control of cardiovascular risk factors, including at a minimum, smoking cessation, anti-platelet and statin therapy.

This systematic review once again highlights the paucity of robust evidence to guide the treatment of patients with diabetes, PAD and foot ulcers. To date, there are no randomised trials of endovascular versus open revascularisation in these patients and the vast heterogeneity of the available data precludes meaningful data pooling or meta-analysis. There remains an urgent need for improved research in this field. Better study design, use of pre-defined and standardised outcomes reporting, including wound healing, will go some way towards producing better quality evidence. However, the fate of a patient with diabetes, foot ulceration and PAD remains difficult to predict, and is affected by a multitude of clinical characteristics that cannot always be identified and ameliorated. The high mortality rates observed amongst these patients should alert the clinician to the significant contribution of concomitant cardiovascular diseases to outcomes, and the overall frailty of the patient cohort. The goals of revascularisation should be reflected in the decision-making process between both the patient and the treating clinician, given the many areas of uncertainty that exist.



## **Conclusions**

This updated systematic review of studies including more than 13,000 patients with diabetes and PAD demonstrates adverse high event rates even when revascularisation is undertaken. This highlights the importance of optimising care of concomitant cardiovascular disease, including medical management and lifestyle modification, particularly in light of the mortality rates of almost 50% at 5 years, worse than many common cancers. There is no appreciable difference in clinical outcomes when comparing endovascular and open therapy and both remain reasonable strategies, depending on the local expertise. Planning a revascularisation approach based on the angiosome concept appears to be a sensible approach in patients with diabetes (especially in those undergoing angioplasty), who typically have poor collateralisation and would likely benefit from revascularisation directly to the feeding artery at the area of tissue loss. However, the data to support this concept are almost entirely retrospective, lacking in standardisation of techniques, definitions and outcome measures. More robust evidence is therefore required in order to understand the best strategy for revascularisation in patients with diabetes, foot ulceration and PAD.

## **Acknowledgements:**

We would like to thank the following external experts for their review of our PICOs for clinical relevance: Stephan Morbach (Germany), Heidi Corcoran (Hongkong), Vilma Urbančič (Slovenia), Rica Tanaka (Japan), Florian Dick (Switzerland), Taha Wassila (Egypt), Abdul Basit (Pakistan), Yamile Jubiz (Colombia), Sriram Narayanan (Singapore), Eduardo Alvarez (Cuba).

We would like to thank Benjamin A. Lipsky (on behalf of the IWGDF editorial board) and Toby Richards (independent external expert) for their peer review of the manuscript.

In addition, we thank Jack Brownrigg for his input into the previous version of this systematic review.

## **Conflict of interest statements**

Production of the 2019 IWGDF Guidelines was supported by unrestricted grants from: Molnlycke Healthcare, Acelity, ConvaTec, Uro Medical, Edixomed, Klaveness, Reaplix, Podartis, Aurealis, SoftOx, Woundcare Circle, and Essity. These sponsors did not have any communication related to the systematic reviews of the literature or related to the guidelines with working group members during the writing of the guidelines, and have not seen any guideline or guideline-related document before publication.

All individual conflict of interest statement of authors of this guideline can be found at: <https://iwgdfguidelines.org/about-iwgdg-guidelines/biographies/>

**Author contributions:** RachaelF performed the updated literature search, screened the titles, abstracts and full papers, assessed the literature, extracted data, drew conclusions for the PICOs, completed the evidence tables, and wrote the manuscript. JA checked the evidence table and reviewed the manuscript. EB assessed the literature, extracted data, checked and revised the evidence tables, reviewed and critically revised the manuscript. RobertF screened the abstracts, assessed the literature, extracted data, checked and revised the evidence tables, and reviewed the manuscript. JPH checked the evidence tables and reviewed the manuscript. KK checked the evidence tables and reviewed the manuscript. JLM extracted data, checked the evidence tables and reviewed the manuscript. SN checked the evidence tables and reviewed the manuscript. JR checked the evidence tables and reviewed the manuscript. MV checked the evidence tables and reviewed the manuscript. REZ extracted data, checked the evidence tables and reviewed the manuscript. NCS assessed the literature, drew conclusions for the PICOs, checked and revised the evidence tables, reviewed and critically revised the manuscript. RJH reviewed and provided final consensus for the data extraction, drew conclusions for the PICOs, reviewed and critically revised the manuscript. RachaelF acted as secretary of the working group, RJH as chair of the working group. The authors take full responsibility for the content of the publication.

## References

1. Prompers L, Schaper N, Apelqvist J, Edmonds M, Jude E, Mauricio D, Uccioli L, Urbancic V, Bakker K, Holstein P, Jirkovska A, Piaggese A, Ragnarson-Tennvall G, Reike H, Spraul M, Van Acker K, Van Baal J, Van Merode F, Ferreira I, Huijberts M. Prediction of outcome in individuals with diabetic foot ulcers: focus on the differences between individuals with and without peripheral arterial disease. The EURODIALE Study. *Diabetologia*. 2008;51(5):747-755. doi:10.1007/s00125-008-0940-0.
2. Jeffcoate WJ, Chipchase SY, Ince P, Game FL. Assessing the outcome of the

- management of diabetic foot ulcers using ulcer-related and person-related measures. *Diab Care*. 2006;29(8):1784-1787. doi:10.2337/dc06-0306.
3. Beckert S, Witte M, Wicke C, Königsrainer A, Coerper S. A new wound-based severity score for diabetic foot ulcers: A prospective analysis of 1,000 patients. *Diab Care*. 2006;29(5):988-992. doi:10.2337/diacare.295988.
  4. Elgzyri T, Larsson J, Nyberg P, Thörne J, Eriksson K-F, Apelqvist J. Early Revascularization after Admittance to a Diabetic Foot Center Affects the Healing Probability of Ischemic Foot Ulcer in Patients with Diabetes. *Eur J Vasc Endovasc Surg*. 2014;48(4):440-446. doi:10.1016/j.ejvs.2014.06.041.
  5. Elgzyri T, Larsson J, Thörne J, Eriksson K-F, Apelqvist J. Outcome of ischemic foot ulcer in diabetic patients who had no invasive vascular intervention. *Eur J Vasc Endovasc Surg*. 2013;46(1):110-117. doi:10.1016/j.ejvs.2013.04.013.
  6. Faglia E, Clerici G, Clerissi J, Gabrielli L, Losa S, Mantero M, Caminiti M, Curci V, Quarantiello A, Lupattelli T, Luppattelli T, Morabito A. Long-term prognosis of diabetic patients with critical limb ischemia: a population-based cohort study. *Diab Care*. 2009;32(5):822-827. doi:10.2337/dc08-1223.
  7. Hinchliffe RJ, Andros G, Apelqvist J, Bakker K, Friederichs S, Fiedrichs S, Lammer J, Lepántalo M, Mills JL, Reekers J, Shearman CP, Valk G, Zierler RE, Schaper NC. A systematic review of the effectiveness of revascularization of the ulcerated foot in patients with diabetes and peripheral arterial disease. Schaper N, Houtum W, Boulton A, eds. *Diabetes Metab Res Rev*. 2012;28 Suppl 1(S1):179-217. doi:10.1002/dmrr.2249.
  8. Hinchliffe RJ, Forsythe RO, Apelqvist JAP, Boyko EJ, Fitridge RA, Hong JP, Katsanos K, Mills JLS, Nikol S, Reekers JA, Venermo M, Zierler RE, Schaper NC, Foot OBOTIWGOTD. IWGDF Guideline on diagnosis, prognosis and management of peripheral artery disease in patients with a foot ulcer and diabetes. *Diabetes Metab Res Rev*, in press.
  9. Garner P, Hopewell S, Chandler J, MacLehose H, Schunemann HJ, Akl EA, Beyene J, Chang S, Churchill R, Dearnness K, Guyatt G, Lefebvre C, Liles B, Marshall R, Martinez Garcia L, Mavergames C, Nasser M, Qaseem A, Sampson M, Soares-Weiser K, Takwoingi Y, Thabane L, Trivella M, Tugwell P, Welsh E, Wilson EC. When and how to update systematic reviews: consensus and checklist. *BMJ*. 2016;354. doi:10.1136/bmj.i3507.
  10. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Journal of Clinical Epidemiology*. 2009;62(10):1006-1012. doi:10.1016/j.jclinepi.2009.06.005.
  11. *SIGN 50: a Guideline Developers' Handbook*. 2001.

<http://www.sign.ac.uk/pdf/sign116.pdf>.

12. Owens CD, Ho KJ, Kim S, Schanzer A, Lin J, Matros E, Belkin M, Conte MS. Refinement of survival prediction in patients undergoing lower extremity bypass surgery: stratification by chronic kidney disease classification. *J Vasc Surg.* 2007;45(5):944-952. doi:10.1016/j.jvs.2007.01.025.
13. Schneider PA, Caps MT, Ogawa DY, Hayman ES. Intraoperative superficial femoral artery balloon angioplasty and popliteal to distal bypass graft: an option for combined open and endovascular treatment of diabetic gangrene. *J Vasc Surg.* 2001;33(5):955-962. doi:10.1067/mva.2001.114210.
14. Rigatelli G, Cardaioli P, Dell'avvocata F, Giordan M, Lisato G, Mollo F. Endovascular management of patients with coronary artery disease and diabetic foot syndrome: A long-term follow-up. *J Geriatr Cardiol.* 2011;8(2):78-81. doi:10.3724/SP.J.1263.2011.00078.
15. Johnson BL, Glickman MH, Bandyk DF, Esses GE. Failure of foot salvage in patients with end-stage renal disease after surgical revascularization. *J Vasc Surg.* 1995;22(3):280-5-discussion285-6.
16. Jämsén T, Manninen H, Tulla H, Matsi P. The final outcome of primary infrainguinal percutaneous transluminal angioplasty in 100 consecutive patients with chronic critical limb ischemia. *J Vasc Interv Radiol.* 2002;13(5):455-463.
17. Darling JD, O'Donnell TFX, Deery SE, Norman AV, Vu GH, Guzman RJ, Wyers MC, Hamdan AD, Schermerhorn ML. Outcomes after first-time lower extremity revascularization for chronic limb-threatening ischemia in insulin-dependent diabetic patients. *J Vasc Surg.* 2018;68(5):1455-1464.e1. doi:10.1016/j.jvs.2018.01.055.
18. Acín F, Varela C, López de Maturana I, de Haro J, Bleda S, Rodriguez-Padilla J. Results of Infrapopliteal Endovascular Procedures Performed in Diabetic Patients with Critical Limb Ischemia and Tissue Loss from the Perspective of an Angiosome-Oriented Revascularization Strategy. *Int J Vasc Med.* 2014;2014(1, part 2):1-13.
19. Troisi N, Turini F, Chisci E, Ercolini L, Frosini P, Lombardi R, Falciani F, Baggio C, Anichini R, Michelagnoli S. Pedal arch patency and not direct-angiosome revascularization predicts outcomes of endovascular interventions in diabetic patients with critical limb ischemia. *Int Angiol.* 2017;36(5):438-444. doi:10.23736/S0392-9590.17.03809-3.
20. Lejay A, Georg Y, Tartaglia E, Gaertner S, Geny B, Thaveau F, Chakfe N. Long-Term Outcomes of Direct and Indirect Below-The-Knee Open Revascularization Based on the Angiosome Concept in Diabetic Patients with Critical Limb Ischemia. *Ann Vasc Surg.* 2014;28(4):983-989. doi:10.1016/j.avsg.2013.08.026.

21. Söderström M, Albäck A, Biancari F, Lappalainen K, Lepäntalo M, Venermo M. Angiosome-targeted infrapopliteal endovascular revascularization for treatment of diabetic foot ulcers. *J Vasc Surg.* 2013;57(2):427-435. doi:10.1016/j.jvs.2012.07.057.
22. Kabra A, Suresh KR, Vivekanand V, Vishnu M, Sumanth R, Nekkanti M. Outcomes of angiosome and non-angiosome targeted revascularization in critical lower limb ischemia. *J Vasc Surg.* 2013;57(1):44-49. doi:10.1016/j.jvs.2012.07.042.
23. Špillerová K, Settembre N, Biancari F, Alback A, Venermo M. Angiosome Targeted PTA is More Important in Endovascular Revascularisation than in Surgical Revascularisation: Analysis of 545 Patients with Ischaemic Tissue Lesions. *Eur J Vasc Endovasc Surg.* 2017;53(4):567-575. doi:10.1016/j.ejvs.2017.01.008.
24. Jeon E-Y, Cho YK, Yoon DY, Kim DJ, Woo JJ. Clinical outcome of angiosome-oriented infrapopliteal percutaneous transluminal angioplasty for isolated infrapopliteal lesions in patients with critical limb ischemia. *Diagn Interv Radiol.* 2016;22(1):52-58. doi:10.5152/dir.2015.15129.
25. Leers SA, Reifsnyder T, Delmonte R, Caron M. Realistic expectations for pedal bypass grafts in patients with end-stage renal disease. *J Vasc Surg.* 1998;28(6):976-980. doi:10.1016/s0741-5214(98)70023-0.
26. Johnson BL, Glickman MH, Bandyk DF, Esses GE. Failure of Foot Salvage in Patients with End-Stage Renal-Disease After Surgical Revascularization. *J Vasc Surg.* 1995;22(3):280-286. doi:10.1016/s0741-5214(95)70142-7.
27. Sigala F, Georgopoulos S, Langer S, Baunach C, Papalambros E, Sigalas K, Bramis J, Bakoyiannis C, Bastounis E, Hepp W. Outcome of infrainguinal revascularization for critical limb ischemia in diabetics with end stage renal disease. *VASA.* 2006;35(1):15-20. doi:10.1024/0301-1526.35.1.15.
28. TANNENBAUM GA, Pomposelli FB, MARCACCIO EJ, Gibbons GW, Campbell DR, Freeman DV, Miller A, Logerfo FW, MEIER G. Safety of Vein Bypass-Grafting to the Dorsal Pedal Artery in Diabetic-Patients with Foot Infections. *J Vasc Surg.* 1992;15(6):982-990.
29. Taylor LM, Porter JM. The Clinical Course of Diabetics Who Require Emergent Foot Surgery Because of Infection or Ischemia. *J Vasc Surg.* 1987;6(5):454-459.
30. Hinchliffe RJ, Brownrigg JRW, Andros G, Apelqvist J, Boyko EJ, Fitridge R, Mills JL, Reekers J, Shearman CP, Zierler RE, Schaper NC, International Working Group on the Diabetic Foot (IWGDF). Effectiveness of revascularisation of the ulcerated foot in patients with diabetes and peripheral artery disease: a systematic review. *Diabetes Metab Res Rev.* 2015;32 Suppl 1:n/a-n/a. doi:10.1002/dmrr.2705.
31. Lepäntalo M, Mätzke S. Outcome of unreconstructed chronic critical leg

- ischaemia. *Eur J Vasc Endovasc Surg*. 1996;11(2):153-157.  
doi:10.1016/S1078-5884(96)80044-X.
32. Bradbury AW, Ruckley CV, Fowkes F, Forbes JF, Gillespie I, Adam DJ, Beard JD, Cleveland T, Bell J, Raab G, Storkey H, Participants BT. Bypass versus angioplasty in severe ischaemia of the leg (BASIL): multicentre, randomised controlled trial. *The Lancet*. 2005;366(9501):1925-1934.  
doi:10.1016/S0140-6736(05)67707-5.
  33. Popplewell MA, Davies HOB, Narayanswami J, Renton M, Sharp A, Bate G, Patel S, Deeks J, Bradbury AW. A Comparison of Outcomes in Patients with Infrapopliteal Disease Randomised to Vein Bypass or Plain Balloon Angioplasty in the Bypass vs. Angioplasty in Severe Ischaemia of the Leg (BASIL) Trial. *Eur J Vasc Endovasc Surg*. 2017;54(2):195-201.  
doi:10.1016/j.ejvs.2017.04.020.
  34. Jongsma H, Bekken JA, Akkersdijk GP, Hoeks SE, Verhagen HJ, Fioole B. Angiosome-directed revascularization in patients with critical limb ischemia. *J Vasc Surg*. 2017;65(4):1208-1219.e1.  
doi:10.1016/j.jvs.2016.10.100.
  35. Chae KJ, Shin JY. Is Angiosome-Targeted Angioplasty Effective for Limb Salvage and Wound Healing in Diabetic Foot? : A Meta-Analysis. Passi AG, ed. *PLoS ONE*. 2016;11(7):e0159523. doi:10.1371/journal.pone.0159523.
  36. Ndip A, Lavery LA, Boulton AJM. Diabetic Foot Disease in People with Advanced Nephropathy and Those on Renal Dialysis. *Curr Diab Rep*. 2010;10(4):283-290. doi:10.1007/s11892-010-0128-0.
  37. AhChong AM, Chiu KM, Wong M, Hui HK, Yip A. Diabetes and the outcome of infrainguinal bypass for critical limb ischaemia. *ANZ J Surg*. 2004;74(3):129-133.
  38. Alexandrescu V, Hubermont G, Philips Y, Guillaumie B, Ngongang C, Coessens V, Vandenbossche P, Coulon M, Ledent G, Donnay JC. Combined Primary Subintimal and Endoluminal Angioplasty for Ischaemic Inferior-limb Ulcers in Diabetic Patients: 5-year Practice in a Multidisciplinary "Diabetic-Foot" Service. *Eur J Vasc Endovasc Surg*. 2009;37(4):448-456.  
doi:10.1016/j.ejvs.2008.12.005.
  39. Bargellini I, Petruzzi P, Scatena A, Cioni R, Cicorelli A, Vignali C, Rizzo L, Piaggese A, Bartolozzi C. Primary infrainguinal subintimal angioplasty in diabetic patients. *Cardiovasc Intervent Radiol*. 2008;31(4):713-722.  
doi:10.1007/s00270-008-9366-9.
  40. DAVIDSON JT, CALLIS JT. Arterial Reconstruction of Vessels in the Foot and Ankle. *Ann Surg*. 1993;217(6):699-710. doi:10.1097/0000658-199306000-00012.
  41. Dosluoglu HH, Cherr GS, Lall P, Harris LM, Dryjski ML. Peroneal artery-only runoff following endovascular revascularizations is effective for limb

- salvage in patients with tissue loss. *J Vasc Surg.* 2008;48(1):137-143.  
doi:10.1016/j.jvs.2008.02.070.
42. Dorweiler B, Neufang A, Schmiedt W, Oelert H. Pedal arterial bypass for limb salvage in patients with diabetes mellitus. *Eur J Vasc Endovasc Surg.* 2002;24(4):309-313.
  43. Faglia E, Mantero M, Caminiti M, Caravaggi C, De Giglio R, Pritelli C, Clerici G, Fratino P, De Cata P, Dalla Paola L, Mariani G, Poli M, Settembrini PG, Sciangula L, Morabito A, Graziani L. Extensive use of peripheral angioplasty, particularly infrapopliteal, in the treatment of ischaemic diabetic foot ulcers: clinical results of a multicentric study of 221 consecutive diabetic subjects. *J Intern Med.* 2002;252(3):225-232.  
doi:10.1046/j.1365-2796.2002.01015.x.
  44. Faglia E, Dalla Paola L, Clerici G, Clerissi J, Graziani L, Fusaro M, Gabrielli L, Losa S, Stella A, Gargiulo M, Mantero M, Caminiti M, Ninkovic S, Curci V, Morabito A. Peripheral angioplasty as the first-choice revascularization procedure in diabetic patients with critical limb ischemia: prospective study of 993 consecutive patients hospitalized and followed between 1999 and 2003. *Eur J Vasc Endovasc Surg.* 2005;29(6):620-627.  
doi:10.1016/j.ejvs.2005.02.035.
  45. Ferraresi R, Centola M, Ferlini M, Da Ros R, Caravaggi C, Assaloni R, Sganzaroli A, Pomidossi G, Bonanomi C, Danzi GB. Long-term Outcomes after Angioplasty of Isolated, Below-the-knee Arteries in Diabetic Patients with Critical Limb Ischaemia. *Eur J Vasc Endovasc Surg* 2009;37(3):336-342. doi:10.1016/j.ejvs.2008.12.001.
  46. Gargiulo M, Maioli F, Ceccacci T, Morselli-Labate AM, Faggioli G, Freyrie A, Giovanetti F, Testi G, Muccini N, Stella A. What's next after optimal infrapopliteal angioplasty? Clinical and ultrasonographic results of a prospective single-center study. *J Endovasc Ther.* 2008;15(3):363-369.  
doi:10.1583/08-2423.1.
  47. Gibbons GW, Burgess AM, Guadagnoli E, Pomposelli FB Jr, Freeman DV, Campbell DR, Miller A, Marcaccio EJ Jr., Nordberg P, Logerfo FW. Return to well-being and function after infrainguinal revascularization. *J Vasc Surg.* 1995;21(1):35-45. doi:10.1016/S0741-5214(95)70242-3.
  48. Hering J, Angelkort B, Keck N, Wilde J, Amann B. Long-term outcome of successful percutaneous transluminal angioplasty of the fibular artery in diabetic foot syndrome and single-vessel calf perfusion depends on doppler wave pattern at the forefoot. *VASA.* 2010;39(1):67-75.  
doi:10.1024/0301-1526/a000007.
  49. Hertzner NR, Bena JF, Karafa MT. A personal experience with the influence of diabetes and other factors on the outcome of infrainguinal bypass grafts for occlusive disease. *J Vasc Surg.* 2007;46(2):271-279.  
doi:10.1016/j.jvs.2007.03.050.

50. Bypass to plantar and tarsal arteries: An acceptable approach to limb salvage. *J Vasc Surg.* 2004;40(6):1149-1157. doi:10.1016/j.jvs.2004.08.037.
51. Isaksson L, Lundgren F. Prognostic factors for failure of primary patency within a year of bypass to the foot in patients with diabetes and critical ischaemia. *Eur J Surg.* 2000;166(2):123-128. doi:10.1080/110241500750009465.
52. Kalra M, Gloviczki P, Bower TC, Panneton JM, Harmsen WS, Jenkins GD, Stanson AW, Toomey BJ, Canton LG. Limb salvage after successful pedal bypass grafting is associated with improved long-term survival. *J Vasc Surg.* 2001;33(1):6-16. doi:10.1067/mva.2001.112300.
53. Kandzari DE, Kiesz RS, Allie D, Walker C, Fail P, Ramaiah VG, Cardenas J, Vale J, Chopra A, Gammon RS. Procedural and clinical outcomes with catheter-based plaque excision in critical limb ischemia. *J Endovasc Ther.* 2006;13(1):12-22. doi:10.1583/05-1634.1.
54. Leers SA, Reifsnyder T, Delmonte R, Caron M. Realistic expectations for pedal bypass grafts in patients with end-stage renal disease. *J Vasc Surg.* 1998;28(6):976-983. doi:10.1016/S0741-5214(98)70023-0.
55. Malmstedt J, Leander K, Wahlberg E, Karlström L, Alfredsson L, Swedenborg J. Outcome after leg bypass surgery for critical limb ischemia is poor in patients with diabetes: a population-based cohort study. *Diab Care.* 2008;31(5):887-892. doi:10.2337/dc07-2424.
56. Mills JL, Gahtan V, Fujitani RM, Taylor SM, Bandyk DF. The utility and durability of vein bypass grafts originating from the popliteal artery for limb salvage. *Am J Surg.* 1994;168(6):646-651. doi:10.1016/S0002-9610(05)80138-2.
57. Mohan CR, Hoballah JJ, Martinasevic M, Chalmers RT, Sharp WJ, Kresowik TF, Corson JD. Revascularization of the ischemic diabetic foot using popliteal artery inflow. *Int Angiol.* 1996;15(2):138-143.
58. Owens CD, Ho KJ, Kim S, Schanzer A, Lin J, Matros E, Belkin M, Conte MS. Refinement of survival prediction in patients undergoing lower extremity bypass surgery: stratification by chronic kidney disease classification. *J Vasc Surg.* 2007;45(5):944-952. doi:10.1016/j.jvs.2007.01.025.
59. Panneton JM, Gloviczki P, Bower TC, Rhodes JM, Canton LG, Toomey BJ. Pedal Bypass for Limb Salvage: Impact of Diabetes on Long-Term Outcome. *Ann Vasc Surg.* 2000;14(6):640-647. doi:10.1007/s100169910114.
60. Pomposelli FB Jr, Marcaccio EJ, Gibbons GW, Campbell DR, Freeman DV, Burgess AM, Miller A, Logerfo FW. Dorsalis pedis arterial bypass: Durable limb salvage for foot ischemia in patients with diabetes mellitus. *J Vasc Surg.* 1995;21(3):375-384. doi:10.1016/S0741-5214(95)70279-2.



61. Pomposelli FB, Kansal N, Hamdan AD, Belfield A, Sheahan M, Campbell DR, Skillman JJ, Logerfo FW. A decade of experience with dorsalis pedis artery bypass: Analysis of outcome in more than 1000 cases. *J Vasc Surg.* 2003;37(2):307-315. doi:10.1067/mva.2003.125.
62. Pua U, Wong DES. Angioplasty in critical limb ischaemia: one-year limb salvage results. *Ann Acad Med Singap.* 2008;37(3):224-229.
63. Ramdev P, Rayan SS, Sheahan M, Hamdan AD, Logerfo FW, Akbari CM, Campbell DR, Pomposelli FB Jr. A decade experience with infrainguinal revascularization in a dialysis-dependent patient population. *J Vasc Surg.* 2002;36(5):969-974. doi:10.1067/mva.2002.128297.
64. Reed AB, Conte MS, Belkin M, Mannick JA, Whittemore AD, Donaldson MC. Usefulness of autogenous bypass grafts originating distal to the groin. *J Vasc Surg.* 2002;35(1):48-54-discussion-54-5. doi:10.1067/mva.2002.120380.
65. Rosenblum BI, Pomposelli FB, Giurini JM, Gibbons GW, Freeman DV, Chrzan JS, Campbell DR, Habershaw GM, Logerfo FW. Maximizing foot salvage by a combined approach to foot ischemia and neuropathic ulceration in patients with diabetes. A 5-year experience. *Diab Care.* 1994;17(9):983-987. doi:10.2337/diacare.17.9.983.
66. Saltzberg SS, Pomposelli FB Jr, Belfield AK, Sheahan MG, Campbell DR, Skillman JJ, Logerfo FW, Hamdan AD. Outcome of lower-extremity revascularization in patients younger than 40 years in a predominately diabetic population. *J Vasc Surg.* 2003;38(5):1056-1059. doi:10.1016/S0741-5214(03)00938-8.
67. Schneider JR, Walsh DB, McDaniel MD, Zwolak RM, Besso SR, Cronenwett JL. Pedal bypass versus tibial bypass with autogenous vein: a comparison of outcome and hemodynamic results. *J Vasc Surg.* 1993;17(6):1029-38-discussion1038-40.
68. Schneider PA, Caps MT, Ogawa DY, Hayman ES. Intraoperative superficial femoral artery balloon angioplasty and popliteal to distal bypass graft: An option for combined open and endovascular treatment of diabetic gangrene. *J Vasc Surg.* 2001;33(5):955-962. doi:10.1067/mva.2001.114210.
69. Söderström M, Arvela E, Alback A, Aho P-S, Lepäntalo M. Healing of ischaemic tissue lesions after infrainguinal bypass surgery for critical leg ischaemia. *Eur J Vasc Endovasc Surg.* 2008;36(1):90-95. doi:10.1016/j.ejvs.2008.01.027.
70. Stonebridge PA, Tsoukas AI, Pomposelli FB Jr, Gibbons GW, Campbell DR, Freeman DV, Miller A, Logerfo FW. Popliteal-to-distal bypass grafts for limb salvage in diabetics. *Eur J Vasc Endovasc Surg.* 1991;5(3):265-269. doi:10.1016/S0950-821X(05)80508-0.

71. Toursarkissian B, Stefanidis D, Hagino RT, D'Ayala M, Schoolfield J, Shireman PK, Sykes MT. Early duplex-derived hemodynamic parameters after lower extremity bypass in diabetics: implications for mid-term outcomes. *Ann Vasc Surg.* 2002;16(5):601-607. doi:10.1007/s10016-001-0272-8.
72. Uccioli L, Gandini R, Giurato L, Fabiano S, Pampana E, Spallone V, Vainieri E, Simonetti G. Long-Term Outcomes of Diabetic Patients With Critical Limb Ischemia Followed in a Tertiary Referral Diabetic Foot Clinic. *Diab Care.* 2010;33(5):977-982. doi:10.2337/dc09-0831.
73. Verhelst R, Bruneau M, Nicolas AL, Frangi R, Khoury El G, Noirhomme P, Dion R. Popliteal-to-distal bypass grafts for limb salvage. *Ann Vasc Surg.* 1997;11(5):505-509.
74. Werneck CCF, Lindsay TF. Tibial angioplasty for limb salvage in high-risk patients and cost analysis. *Ann Vasc Surg.* 2009;23(5):554-559. doi:10.1016/j.avsg.2009.05.007.
75. Woelfle KD, Lange G, Mayer H, Bruijnen H, Loeprecht H. Distal vein graft reconstruction for isolated tibioperoneal vessel occlusive disease in diabetics with critical foot ischaemia--does it work? *Eur J Vasc Surg.* 1993;7(4):409-413.
76. Wölfle KD, Bruijnen H, Reeps C, Reutemann S, Wack C, Campbell P, Loeprecht H, Häuser H, Bohndorf K. Tibioperoneal arterial lesions and critical foot ischaemia: successful management by the use of short vein grafts and percutaneous transluminal angioplasty. *VASA.* 2000;29(3):207-214. doi:10.1024/0301-1526.29.3.207.
77. Woelfle, Bruijnen, Loeprecht. Infrapopliteal arterial occlusive disease in diabetics with critical foot ischaemia: The role of distal-origin bypass grafts. *VASA.* 2013;30(Supplement 58):40-43. doi:10.1024/0301-1526.30.S58.40.
78. Zayed H, Halawa M, Maillardet L, Sidhu PS, Edmonds M, Rashid H. Improving limb salvage rate in diabetic patients with critical leg ischaemia using a multidisciplinary approach. *Int J Clin Pract.* 2009;63(6):855-858. doi:10.1111/j.1742-1241.2007.01608.x.
79. Park SW, Kim JS, Yun IJ, Hwang JJ, Lee SA, Chee HK, Chang S-H, Shin DH, Jung HG, Soo Chang II. Clinical outcomes of endovascular treatments for critical limb ischemia with chronic total occlusive lesions limited to below-the-knee arteries. *Acta Radiol.* 2013;54(7):785-789. doi:10.1258/ar.2012.120217.
80. Zhan LX, Bharara M, White M, Bhatnagar S, Lepow B, Armstrong DG, Mills JL. Comparison of initial hemodynamic response after endovascular therapy and open surgical bypass in patients with diabetes mellitus and critical limb ischemia. *J Vasc Surg.* 2012;56(2):380-6-discussion386.

doi:10.1016/j.jvs.2012.01.050.

81. Alexandrescu V, Vincent G, Azdad K, Hubermont G, Ledent G, Ngongang C, Filimon A-M. A Reliable Approach to Diabetic Neuroischemic Foot Wounds: Below-the-Knee Angiosome-Oriented Angioplasty. *J Endovasc Ther.* 2011;18(3):376-387. doi:10.1583/10-3260.1.
82. Darling JD, Bodewes TCF, Deery SE, Guzman RJ, Wyers MC, Hamdan AD, Verhagen HJ, Schermerhorn ML. Outcomes after first-time lower extremity revascularization for chronic limb-threatening ischemia between patients with and without diabetes. *J Vasc Surg.* 2018;67(4):1159-1169. doi:10.1016/j.jvs.2017.06.119.
83. Mohapatra A, Henry JC, Avgerinos ED, Boitet A, Chaer RA, Makaroun MS, Leers SA, Hager ES. Bypass versus endovascular intervention for healing ischemic foot wounds secondary to tibial arterial disease. *J Vasc Surg.* 2018;68(1):168-175. doi:10.1016/j.jvs.2017.10.076.
84. Lo ZJ, Lin Z, Pua U, Quek LHH, Tan BP, Punamiya S, Tan GWL, Narayanan S, Chandrasekar S. Diabetic Foot Limb Salvage-A Series of 809 Attempts and Predictors for Endovascular Limb Salvage Failure. *Ann Vasc Surg.* 2018;49:9-16. doi:10.1016/j.avsg.2018.01.061.
85. Hicks CW, Najafian A, Farber A, Menard MT, Malas MB, Black JH, Abularrage CJ. Diabetes does not worsen outcomes following infrageniculate bypass or endovascular intervention for patients with critical limb ischemia. *J Vasc Surg.* 2016;64(6):1667-1674.e1. doi:10.1016/j.jvs.2016.07.107.

**Figure 1: PRISMA Flow Diagram**



