



Drake, M. J., Lewis, A. L., Young, G., Abrams, P. H., Blair, P. S., Chapple, C., Glazener, C., Horwood, J. P., McGrath, J., Noble, S. M., Taylor, G., & Lane, J. A. (2020). Diagnostic Assessment of Lower Urinary Tract Symptoms in Men Considering Prostate Surgery: A Noninferiority Randomised Controlled Trial of Urodynamics in 26 Hospitals. *European Urology*, 78(5), 701-710.
<https://doi.org/10.1016/j.eururo.2020.06.004>

Peer reviewed version

Link to published version (if available):
[10.1016/j.eururo.2020.06.004](https://doi.org/10.1016/j.eururo.2020.06.004)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Elsevier at <https://doi.org/10.1016/j.eururo.2020.06.004> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Including UDS in the diagnostic tests for male LUTS resulted in a non-inferior symptom outcome (IPSS) to routine care, 18-months post-randomisation. However, there was no reduction in rates of surgery for treating BOO. Accordingly, the findings show that routine use of UDS in evaluation of uncomplicated LUTS has a limited role and should be used selectively.