



Hudson, P., Emmett, P. M., & Taylor, C. M. (2021). Pre-pregnancy maternal BMI classification is associated with preschool childhood diet quality and childhood obesity in the Avon Longitudinal Study of Parents and Children. *Public Health Nutrition*, 24(18), 6137-6144. <https://doi.org/10.1017/S1368980021001476>

Peer reviewed version

Link to published version (if available):  
[10.1017/S1368980021001476](https://doi.org/10.1017/S1368980021001476)

[Link to publication record on the Bristol Research Portal](#)  
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Cambridge University Press at [10.1017/S1368980021001476](https://doi.org/10.1017/S1368980021001476). Please refer to any applicable terms of use of the publisher.

## University of Bristol – Bristol Research Portal

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available: <http://www.bristol.ac.uk/red/research-policy/pure/user-guides/brp-terms/>

# **Pre-pregnancy maternal BMI classification is associated with preschool childhood diet quality and childhood obesity in the Avon Longitudinal Study of Parents and Children**

Paul Hudson<sup>1</sup>, Pauline M Emmett, Caroline M Taylor

Centre for Academic Child Health, Bristol Medical School, University of Bristol, Bristol, UK

<sup>1</sup>Present address: 48A Osborne Road, Portsmouth P05 3LT, Hants, UK

**Corresponding author:** Dr Caroline M Taylor, Centre for Academic Child Health, Bristol Medical School, University of Bristol, 1–5 Whiteladies Rd, Bristol BS8 1NU, UK

Caroline.m.taylor@bristol.ac.uk

**Short title:** Maternal BMI and childhood diet quality

**Acknowledgements:** We are extremely grateful to all the families who took part in this study, the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists and nurses.

**Financial Support:** The UK Medical Research Council and the Wellcome Trust (Grant ref: 102215/2/13/2) and the University of Bristol currently provide core support for ALSPAC. A comprehensive list of grants funding is available on the ALSPAC website (<http://www.bristol.ac.uk/alspac/external/documents/grant-acknowledgements.pdf>). CMT was supported by a Wellcome Career Re-entry Fellowship (Grant ref: 104077/Z/14/Z), by the Elizabeth Blackwell Institute for Health Research, University of Bristol, and the Wellcome Trust Institutional Strategic Support Fund, and by an MRC Career Development Award (Grant ref: MR/T010010/1). This publication is the work of the authors who serve as guarantors for the contents of the paper.

**Conflict of interest:** PH has no conflicts of interest. CMT and PME have received research funding from Nestlé Nutrition for unrelated research.

**Authorship:** PME and CMT conceived the study. PH undertook all analyses with consultation with PME and CMT. PH wrote the manuscript with critical inputs from PME and CMT. All authors approved the final version of the paper.

**Ethical standards disclosure:** This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving research study participants were approved by the ALSPAC Ethics and Law Committee and the Local Research Ethics Committees. **Informed consent for the use of data collected via questionnaires and clinics was obtained from participants following the recommendations of the ALSPAC Ethics and Law Committee at the time.**

## Abstract

**Objective** To explore the effect of maternal BMI class pre-pregnancy (overweight/obese versus healthy weight/underweight) on childhood diet quality and on childhood overweight/obesity risk.

**Design** Dietary data were collected using 3-day parental-completed food records for their children at ages 18 and 43 months. ~~Items in the food records were linked to weights and energy contents.~~ An index of diet quality was derived by classification of food items into core and non-core foods. Adjusted multiple linear regression analyses were used to explore the effect of maternal BMI class (overweight/obese (BMI  $\geq 25$  kg/m<sup>2</sup>) compared to healthy weight/underweight) on diet quality in their children.

**Setting** Avon, UK.

**Participants** A 10% subsample of the Avon Longitudinal Study of Parents and Children. 908 children provided complete dietary data at 18 months and 769 at 43 months.

**Results** Children with overweight/obese mothers consumed greater amounts of energy from non-core foods than children with healthy weight/underweight mothers (0.20 MJ [48 kcal]/day more at 18 months (p<0.001); 0.19 MJ [45 kcal]/day more at 43 months (p=0.008)) in adjusted models. ~~This included 0.06 MJ [14 kcal]/day more free sugar at 18 months (p=0.010).~~ Diet quality deteriorated between 18 and 43 months (children reduced their dietary energy intake from core foods (p<0.001) and increased intake from non-core foods (p<0.001)). However, this change was not associated with maternal BMI class in adjusted models. ~~(core foods p=0.271, non-core foods p=0.332).~~ Having an overweight/obese mother was associated with an increased odds of the child being overweight/obese at 43 months (OR 1.74 (1.17, 2.58)).

**Conclusion** Children aged 18 and 43 months with overweight/obese mothers are likely to have a poorer quality diet than those with healthy-/underweight mothers. Parents should be supported in discouraging the consumption of non-core foods in children at these ages.

[239-243 words]

**Keywords** ALSPAC, Obesity, Overweight, Diet, Childhood, Maternal BMI

## Introduction

Childhood obesity is a major and increasing public health concern in the UK and worldwide. In England, 28% of children **age 2-15 years old** are overweight, including 15% who are obese<sup>(1)</sup>. Obese children are more likely to suffer from medical conditions such as type 2 diabetes, sleep apnoea and cardiovascular disease<sup>(2)</sup>. They are also more likely to face socio-emotional difficulties such as discrimination and social marginalisation<sup>(3)</sup>.

In the UK, the general quality of diet in children is poor, including during early childhood. UK children between the ages 1.5 to 3 years, on average, consume more than double the maximum 5% energy intake from free sugars that is recommended by the Scientific Advisory Committee on Nutrition<sup>(4)</sup>. They also consume nearly one-third more than the recommended percentage energy intake from saturated fat<sup>(4)</sup>. In addition, only 18% of children between the ages 5 and 15 years eat the recommended five or more portions of fruit and vegetables a day<sup>(5)</sup>. When obesity develops in childhood, it significantly increases the odds of obesity in adulthood<sup>(6)</sup>. Therefore, addressing the causes of poor diet and obesity early in life is vitally important in reducing the prevalence and impact of obesity.

Maternal overweight/obesity is a significant risk factor for childhood obesity<sup>(7)</sup>. In England, the odds of childhood obesity more than doubles if a child has two overweight parents compared with a child with two healthy weight parents<sup>(7)</sup>. A number of transgenerational risk factors likely contribute to this relationship, such as parental influences over their child's diet, physical activity, behaviour and genetic vulnerability<sup>(8)</sup>. Obese mothers are also less likely to breastfeed, and do so for a shorter duration, than non-obese mothers<sup>(9)</sup>. Despite these findings, few studies have explored how the solid food diets of children with overweight/obese parents compare with those of children with healthy weight parents.

A previous study in the Avon Longitudinal Study of Parents and Children (ALSPAC) showed that childhood diet quality significantly deteriorates from ages 18 months to 43 months, and that this lower quality diet is maintained throughout childhood<sup>(10)</sup>. The primary aim of this study was to determine the association of maternal BMI classification on the quality of early childhood diet and risk of childhood obesity by exploring the effect on the consumption of core and non-core foods and soft drinks in children at ages 18 and 43 months, and on childhood BMI classification at 43 months of age. As a secondary aim, the predictive value of maternal BMI classification for a change in diet quality from ages 18 to 43 months was also explored.

## **Methods**

### ***Participants***

ALSPAC is an observational prospective cohort study investigating influences on health and development across the life course<sup>(11, 12)</sup>. The cohort has been described in greater detail elsewhere<sup>(11, 12)</sup>. In brief, the study initially recruited 14,541 pregnant women in the former Avon Health Authority in south-west England, with an expected delivery date between April 1991 and December 1992. From these initial pregnancies, 14,676 fetuses were produced, resulting in 14,062 live births and a total of 13,988 children who were alive at 1 year of age (Supplementary Figure 1). The cohort has since collected health, lifestyle and dietary information, through questionnaires and clinical visits from these women and their resulting offspring. The cohort is broadly representative of the UK population, although the participants are slightly more prosperous and the cohort is under-representative of ethnic minority participants<sup>(11)</sup>. The study website contains details of all the data that are available through a fully searchable data dictionary and variable search tool <http://www.bris.ac.uk/alspac/researchers/data-access/data-dictionary>.

Participants in ALSPAC's Children in Focus (CiF) subsample, equivalent to 10% of the entire ALSPAC cohort, were selected from the last 6 months of ALSPAC births. The selected children along with their parents were invited to attend research clinics. These clinics took place in intervals from 4 months to 61 months of age, with 1432 children attending at least one clinic. Children from this subsample are biased towards having mothers with a higher educational attainment compared with the entire ALSPAC cohort (39.6% of children in CiF have mothers whose educational attainment was A levels (Advanced level) or above compared with 35.4% in the entire ALSPAC cohort<sup>(13)</sup>).

### ***Exposures***

Pre-pregnancy maternal BMI was derived from self-reported pre-pregnancy weight and height, from which BMI was calculated ( $\text{weight}/\text{height}^2$ ). A pre-pregnancy maternal BMI classification variable was derived from the World Health Organization<sup>(14)</sup> categories: 'overweight/obese' ( $\text{BMI} \geq 25 \text{ kg/m}^2$ ) (ow/ob) or 'healthy weight/underweight' ( $\text{BMI} < 25 \text{ kg/m}^2$ ) (hw/uw).

### ***Outcomes***

Dietary information was collected when the child was aged 18 months and again aged 43 months. Detailed methods have been described elsewhere<sup>(10)</sup>. In summary, parents were invited by post to complete structured food records, recording all food and drink their child consumed over three individual days (preferable including two weekdays and one weekend day). The food records were checked by a member of the nutrition team for completeness with the parent and to clarify any uncertainties.

Food and drink items from the food record were linked to weights and energy contents and the data were grouped according to the concepts of 'core' or 'non-core' foods or soft drinks (Supplementary Table 1) based on the Australian Guide to Healthy Eating<sup>(15)</sup>. Core foods are described as those which provide the important nutrients the body needs; non-core food or 'extra foods' are described as those which are not necessary to provide the nutrients the body needs and often contain added fat, sugar or salt<sup>(15)</sup> and soft drinks are non-alcoholic, non-dairy drinks with low nutrient content. The core food groups created were: (1) Fruit; (2) Potatoes, vegetables and legumes; (3) Unprocessed meat, fish, poultry and eggs; (4) Cereal foods. The non-core food groups created were: (1) Processed meat, fish and poultry; (2) Potatoes cooked in fat; (3) Spreads, soups and sauces; (4) Sweetened processed foods. **Soft drinks were defined as a 'cold sweet drink that does not contain alcohol' and subdivided into three subgroups:**(1) Unsweetened fruit juice; (2) Sugar-sweetened soft drinks; (3) Artificially sweetened soft drinks (Supplementary Table 1).

The BMI of the children was measured at CiF clinics at 43 months of age. These were derived from weight and height measurements made by trained clinical staff using standardised procedures.

### ***Confounders***

Confounders were selected based on whether the characteristic occurred at about the same time as the exposure, were scientifically plausible as confounders and had a p-value <0.2 in the baseline characteristic analyses.

Data collected on maternal characteristics from maternal completed postal questionnaires included age (years), maternal education, parity and whether the mother had ever breastfed this child. Birthweight (g) was obtained from medical records. Family Adversity Index (FAI) was derived from responses to questions relating to measures of financial, social, emotional and physical adversity, which have been detailed elsewhere<sup>(16)</sup>. Scores range from 0 to 8. For maternal education, mothers were either categorised as having achieved Certificate of School



Education (CSE), vocational or O-level (Ordinary level) qualifications only, or as having achieved A-level, degree or higher qualifications. For maternal parity, mothers were either categorised as 0 (no previous births) or  $\geq 1$  (1 or more previous births).

### **Data analysis**

All statistical tests were performed on Stata, version 15 (StataCorp, College Station, TX, USA). BMI classifications were calculated using the Zantho package in STATA. Maternal age, maternal education, parity and FAI were used as covariates in all adjusted analyses.

Unpaired t-tests and chi-squared tests were used to test for baseline characteristic differences between ow/ob mothers and hw/uw mothers.

**Effect of maternal pre-pregnancy BMI class on diet at 18 and 43 months** The mean daily intakes of all core- and non-core food and soft drink groups (weight and energy) were calculated using the dietary data available at 18 and at 43 months of age, with the children categorised by maternal BMI class.

Multiple linear regression analyses were used to determine the association of maternal BMI classification with the daily mean energy intakes from total food, total core and non-core food and soft drinks at 18 months and 43 months.

Multiple linear regression analyses were repeated to determine the association of maternal BMI class with mean daily energy intakes from individual core and non-core food and soft drinks groups, and from individual macronutrients at 18 months and 43 months.

**Odds of childhood ow/ob at 43 months** Logistic regression was used to determine the odds of a child being overweight or obese at 43 months if they had an ow/ob mother (maternal pre-pregnancy BMI  $\geq 25$  kg/m<sup>2</sup>). Child BMI classifications are based on the British 1990 growth reference centile curves<sup>(17)</sup>: overweight or obese defined as BMI  $\leq$  85th or 95th centile respectively of BMIs within their age group and sex<sup>(18)</sup>.

**Changes in diet from 18 to 43 months** Paired t-tests were used to determine the percentage change, from 18 to 43 months, of total energy intake from core and non-core food and soft drink groups. Linear regression analyses were undertaken to explore the effect of maternal ow/ob status on this percentage change from 18 to 43 months.

## **Results**

### ***Demographic characteristics of study group***

The demographic characteristics of the study groups are shown in Table 1. In summary, ow/ob mothers had lower educational attainment ( $p < 0.001$ ) than hw/uw mothers, they were more likely to not have ever breastfed ( $p < 0.001$ ), and their babies had a greater birthweight ( $p < 0.001$ ). They also had a higher FAI, but the difference was weak ( $p = 0.061$ ). There were no differences between the groups for smoking status, parity and age.

### ***Changes in diet from 18 to 43 months***

Energy intakes in children at 18 and 43 months categorised by maternal BMI class are shown in Supplementary Table 2. As expected, total energy intake increased between 18 and 43 months by 23% for children with hw/uw mothers and by 21% for children with ow/ob mothers. However, energy intake from total non-core foods increased by a greater proportion in children with ow/ob mothers (non-core food energy intake increased by 68% amongst children with ow/ob mothers and by 58% amongst children with hw/uw mothers). There were many more consumers of soft drinks at 43 months than at 18 months. The percentage of consumers of sugar sweetened drinks rose from 34% to 60% in both groups of children while low calorie/diet drinks went from 64% to 73% in children of hw/uw mothers and from 72% to 82% in those of ow/ob mothers.

### ***Energy intake from total core and non-core foods***

In adjusted models total daily energy intake did not differ between children with ow/ob mothers and those with hw/uw mothers either at 18 months ( $p = 0.176$ ) or at 43 months ( $p = 0.564$ ). In addition, total daily core food energy intake did not differ between the two groups of children either at 18 months ( $p = 0.105$ ) or at 43 months ( $p = 0.109$ ) (Table 2).

Children with ow/ob mothers had a greater daily mean intake of energy from non-core foods than children with hw/uw mothers at both 18 and 43 months of age. At 18 months non-core energy intake was 0.20 MJ [48 kcal]/day (10.0%) greater ( $p < 0.001$ ) and at 43 months it was 0.19 MJ [45 kcal]/day (7.2%) greater ( $p = 0.008$ ) (Table 2).

### ***Energy intakes from macronutrients***

In adjusted models, at 18 months of age children with ow/ob mothers had a 0.06 MJ [14 kcal] (9.7%) greater daily mean intake of energy from free sugars than children with hw/uw mothers ( $p = 0.010$ ), and a 0.06 MJ [14 kcal] (11.1%) lower energy intake from intrinsic sugar

( $p=0.002$ ) (Table 2). There were no differences in the other macronutrients (data not shown). By 43 months, there was no difference between the two groups of children in the daily mean energy intakes from any of the macronutrients (Table 2).

#### ***Energy intake from individual core and non-core food groups***

In adjusted models for core foods at 18 months, only two food groups differed between the two maternal groups: total fruit (excluding fruit juice) energy intake was 14.8% ( $p=0.012$ ) lower and the yogurt, cheese and milk group was 7.7% ( $p=0.037$ ) lower in children with ow/ob mothers compared with those with hw/uw mothers (Table 3). There were no differences at 43 months.

For non-core foods at 18 months, energy intakes from most of the food groups were greater amongst children with ow/ob mothers than children with hw/uw mothers: processed meat, fish, poultry ( $p=0.015$ ), potatoes cooked in oil and baked beans ( $p=0.007$ ), and sweetened processed foods and savoury snacks ( $p=0.012$ ) (Table 3). Energy intake from spreads, soups and sauces was marginally greater ( $p=0.050$ ) in children with ow/ob mothers (Table 3). At 43 months, the daily mean energy intake from processed meat, fish, eggs and poultry was greater amongst children with ow/ob mothers than those with hw/uw mothers ( $p=0.019$ ). There was weak evidence for greater energy intake from sweetened processed foods ( $p=0.087$ ) in children with ow/ob mothers (Table 3). There were no differences in the energy from unsweetened fruit juice or sugar sweetened drinks between the two groups at either age (Table 3), and for low calorie/diet drinks at 43 months. At 18 months children with ow/ob mothers who were consumers of low calorie/diet drinks consumed 4 kJ/day more energy from low calories/diet drinks than children of hw/uw mothers.

#### ***Change in energy intake from core and non-core foods from 18 to 43 months of age***

Across all children, the percentage of total dietary energy intake from core foods decreased by 10.6% from 18 to 43 months ( $p<0.001$ ), whilst the percentage of total dietary energy from non-core foods increased by 10.9% ( $p<0.001$ ) (Supplementary Table 2). **The percentage of energy from non-core foods in children with ow/ob mothers was 14.6% greater at 18 months and 9.3% greater at 43 months than in those with hw/uw mothers (Fig. 1).** However, these changes were of the same magnitude in children with ow/ob mothers as in those with hw/uw mothers in adjusted models **(core foods: beta coefficient 1.41 (95% CI -1.10, 3.93),  $p=0.271$ ,  $n=687$ ; non-core: foods beta coefficient -1.20 (95% CI -3.63, 1.23),  $p=0.332$ ,  $n=687$ ).**

### *Odds of obesity at 43 months*

Children with ow/ob mothers had an OR of 1.74 (1.17, 2.58) of being ow/ob at 43 months compared with children with hw/uw mothers in adjusted analyses (p=0.006).

## **DISCUSSION**

This is the first study of diet in young children to show that maternal ow/ob is associated with poor diet quality in preschool children. Overall, children with ow/ob mothers were found to consume greater amounts of all of the non-core food groups, and lower amounts of some core food groups, than children with hw/uw mothers. Furthermore, children in both groups had a lower quality diet at 43 months than at 18 months with a move towards the consumption of more non-core and less core foods.

Non-core foods are considered excess to requirements for maintaining a healthy- childhood diet and are often consumed between meals as snacks<sup>(15)</sup>. The consumption of such low quality foods has been associated with overweight in childhood<sup>(19)</sup>. Furthermore, the consumption of non-core food has been increasing amongst children over recent decades, mirroring the increase in childhood obesity prevalence. In this study children of ow/ob mothers consumed, on average, 10.0% (18 months) and 7.2% (43 months) more energy daily from non-core foods than children with hw/uw mothers. This difference suggests that maternal ow/ob is a risk factor for poor diet quality during early childhood.

A US study comparing past and present food intake amongst children using nationally representative large surveys shows that children have considerably increased their consumption of unhealthy snack foods since the 1970s. In 2006, 27% of children's daily energy intake was from snacks<sup>(20)</sup>, which is a similar percentage to that found in this study. However, the US sample of children were of an older mean age and were more ethnically diverse than children in this study. The US investigation also found that children aged 2–6 years snack more than any other child age group and increased the amount they snacked the most from 1977 to 2006<sup>(20)</sup>. The results of the present study are the first to show that maternal ow/ob is a risk factor for the consumption of such unhealthy snack foods in young children.

Regular consumption of core food groups is highly important to maintain a healthy diet and provide essential nutrients<sup>(15)</sup>. In this study the 18-month-old children with hw/uw mothers consumed higher amounts of fresh fruit than those with ow/ob mothers. Although there was

no difference in total core-food consumption between the groups. In England, children consume considerably less than the recommended amounts of fruits and vegetables<sup>(5)</sup>. The results of this study suggest that having an ow/ob mother is a risk factor for low fruit consumption at 18 months of age. In addition, at 18 months, children with ow/ob mothers consumed less of the yogurt, cheese and milk group than children with hw/uw mothers. An Australian cohort study found that dairy consumption at 18 months was negatively associated with BMI at age 8 years<sup>(21)</sup>, suggesting that dairy consumption at this age may be protective against childhood obesity. The Australian study design was similar to that of this study as both are cohort-based and used 3-day structured food records to collect dietary information. However, a smaller sample size was used in the Australian study (n=364)<sup>(21)</sup>. In children from the US, there has been a decrease in the consumption of fresh fruit, milk and dairy products from 1976 to 2006, whilst the prevalence of obesity has increased<sup>(20)</sup>.

Less intrinsic sugars (found in foods such as fruit, vegetables and milk) was consumed by the 18-month-old children with ow/ob mothers than children with hw/uw mothers in this study. At the same time these children consumed greater amounts of free sugars. Free sugars are found in sugar-sweetened beverages, sweetened processed foods, confectionary and chocolate. It has been found that the consumption of sugar-sweetened beverages, in particular, can increase the risk of childhood obesity<sup>(22)</sup>. However, in this study there was no difference in the percentage of consumers of sugar-sweetened beverages between the two groups of children at either age although the overall percentage increased with age. In some studies high free sugar consumption has been associated with low intakes of fruits and vegetables<sup>(23)</sup> <sup>(24)</sup>, thus impacting negatively on core food consumption. A large Swedish cohort study (n=10,762) found a positive association between maternal ow/ob and the early introduction of sweets/pastries and sugar-sweetened drinks in infancy<sup>(25)</sup>.thus supporting the findings of the present study that maternal ow/ob is associated with increased free sugars consumption at a very early age.

The results of this study support other findings that maternal ow/ob significantly increases the odds of childhood obesity<sup>(5)</sup>. It has also provided evidence that ow/ob mothers provide their children with a poorer quality diet than hw/uw mothers. This may therefore be a mediating factor in the association between maternal ow/ob and child ow/ob at 43 months.

This study has confirmed what has been found previously: that there is a large decrease in core-food consumption from age 18 to 43 months, whilst non-core food and soft drink

consumption increases<sup>(10)</sup>. Therefore, diet quality seems to undergo a deterioration from ages 18 to 43 months. Both children with ow/ob mothers and those with hw/uw mothers experienced a deterioration; however, there was limited evidence for difference in the magnitude of this deterioration between the two groups. This is because the diet quality of children with ow/ob mothers was considerably worse than children with hw/uw mothers both at both 18 months and 43 months of age. In addition, maternal ow/ob has its largest detrimental impact on the quality of diet at 18 months rather than 43 months of age.

There are several strengths of this study. (1) The fact that dietary data were derived from detailed food records, collected prospectively with the same children followed at 18 and 43 months allows the detection of changes in diet occurring across childhood. (2) Food record data were collected over a 3-day period and verified by trained staff in parental interviews. The data are therefore extensive, allowing for a comprehensive assessment of diet. (3) The large sample size provides a high power to detect associations. (4) ALSPAC provides a range of socio-economic data and health information relating to these children and mothers that facilitates the consideration of confounders.

There are also limitations for the study. (1) The food records were completed by caregivers who may have mis-recorded their child's diet. Any systematic differences in the misrepresentation of diet could bias the findings of this study. (2) Weights and energy content of foods were estimated using standard food tables, rather than directly measured. However, alternative approaches such as weighing all food items would have been more time consuming and challenging for participants, likely resulting in underreporting or alternation from normal diet. In addition, the dietary information may not be entirely representative of long-term diet. However, these dietary collection issues equally affect the food record data collected at 18 and 43 months, and therefore should not result in biases in the direct comparisons made between these two ages. (3) BMI measurements of pre-maternal BMI were derived from self-reported weight and height measurements which limit their accuracy. In contrast, child BMI measurements were made in clinics by trained professions, providing robust data for child BMI. (4) Despite generally high retention rates in ALSPAC, attrition is an issue in any longitudinal prospective study. The sample size used in the 43-month analyses was about 15% smaller than that used in the 18-month analyses. In ALSPAC, loss to follow up tends to be greater amongst mothers and children from lower socioeconomic background, which may result in bias in this study (5) The results may not be generalisable to other parts of the UK or other countries because the ALSPAC study was undertaken in one

geographically defined area. Although the cohort was similar to the UK population at recruitment, some differences existed<sup>(12)</sup>. These differences are more apparent in CiF subsample used in this study, with children more likely to have a mother with a higher educational attainment than in the general UK population<sup>(13)</sup>. (6) Dietary information was taken from children in this study in the year 1994 (when the children were aged 18 months) and again in 1996 (when the children were aged 43 months). In 1995, the prevalence of childhood obesity in England was 11% for boys and 12% for girls<sup>(26)</sup>. Presently, 15% of children in England are obese<sup>(1)</sup> and so it is likely that current childhood diet plays an even more important role in causing childhood obesity. However, it has been shown that ALSPAC dietary data is similar to current food and nutrient intakes amongst children<sup>(10)</sup>, this supports the generalisability of this study's findings to the current UK context.

Future research is needed to explore the reasons why maternal ow/ob is a risk factor for poor diet quality during early childhood. The relative contribution of genetic inheritance, learned eating behaviour, food availability and other possible transgenerational risk factors should be investigated. This will help to inform future interventions that aim to improve diet quality in young children.

### **Conclusion**

The findings of this study provide evidence that overweight or obese mothers are more likely to feed their children a poorer quality of diet than hw/uw mothers and this may contribute to the increased odds of them being obese before school age. This indicates that reducing or preventing obesity in mothers may have the secondary benefit of improving the diet quality of their children and therefore helping to tackle childhood obesity. The results show that mothers can have a significant influence over the quality of their child's diet, which should be considered when developing public health strategies to improve childhood diet quality.

Diet quality deteriorated from ages 18 to 43 months. However, these changes occurred irrespective of maternal BMI class. The findings from the study support the view that children should be encouraged to eat a diet that is high in core foods and low in non-core foods such as processed foods high in fat and sugar. This can be achieved through the promotion of three balanced meals a day, whilst encouraging the provision of 'healthy' snack foods and water to drink.

**Table 1** Demographic characteristics of overweight/obese mothers compared with those with healthy weight/underweight mothers in ALSPAC-CiF

	<b>Maternal BMI classification group</b>		<b>P value</b>
	<b>Healthy weight/underweight</b>	<b>Obese/overweight</b>	
Maternal age (years)	29.4 (4.3)	29.0 (4.4)	0.189
Smoking			0.722
Yes	87 (12.2%)	24 (11.3%)	
No	628 (87.8%)	189 (88.7%)	
Maternal education			<0.001
A level or above	341 (44.9%)	74 (33.8%)	
O level/CSE/None	419 (55.1%)	145 (66.2%)	
Family adversity index	0.98 (1.31)	1.17 (1.44)	0.061
Parity			0.183
0	96 (43.7%)	365 (48.7%)	
≥1	124 (56.3%)	48.7 (51.3%)	
Birthweight (g)	3434 (503)	3563 (521)	<0.001
Baby ever breastfed			<0.001
Yes	608 (82.9%)	147 (69.7%)	
No	125 (17.7%)	64 (30.3%)	

Values are mean (SD) or n (%)

BMI classifications: healthy-/underweight <25 kg/m<sup>2</sup>, overweight/obese ≥25 kg/m<sup>2</sup>.



**TABLE 2** The differences in energy intake (MJ/day) from all foods, core and non-core foods and macronutrients in ALSPAC-CiF children at 18 and 43 months of age with overweight/obese mothers compared with those with healthy weight/underweight mothers

	Age attending CiF clinic			
	18 months (n=908)		43 months (n=769)	
	$\beta$ coefficient (95% CI) <sup>a</sup>	P value	$\beta$ coefficient (95% CI) <sup>a</sup>	P value
Total energy intake	0.10 (-0.05, 0.25)	0.176	0.05 (-0.13, 0.23)	0.564
Energy intake from all core foods	-0.10 (-0.22, 0.02)	0.105	-0.12 (-0.27, 0.03)	0.109
Energy intake from all non-core-foods	0.20 (0.10, 0.30)	<0.001	0.19 (0.05, 0.33)	0.008
Energy intake from all soft drinks	0.10 (-0.33, 0.53)	0.656	-0.24 (-0.58, 0.10)	0.163
Protein	0.00 (-0.03, 0.03)	0.889	-0.00 (-0.03, 0.03)	0.958
Fat	0.04 (-0.03, 0.11)	0.253	0.04 (-0.05, 0.12)	0.382
Carbohydrate	0.06 (-0.02, 0.14)	0.122	0.02 (-0.08, 0.11)	0.753
Total sugar	-0.00 (-0.05, 0.05)	0.960	-0.03 (-0.10, 0.04)	0.412
Intrinsic sugar	-0.06 (-0.10, -0.02)	0.002	-0.01 (-0.06, 0.03)	0.504
Free sugar	0.06 (0.01, 0.10)	0.010	-0.01 (-0.07, 0.05)	0.647

BMI classifications: healthy-/underweight <25 kg/m<sup>2</sup>, overweight/obese ≥25 kg/m<sup>2</sup>.

<sup>a</sup>Adjusted for maternal age, maternal education, parity and Family Adversity Index.

Reference group: children with healthy-/underweight mothers.

Conversion factor: 1 MJ is equivalent to 239 kcal.

**TABLE 3** Difference in energy intake (MJ/day) from individual core and non-core food groups in ALSPAC-CiF children at 18 and 43 months of age with overweight/obese mothers compared with those with healthy-/underweight mothers

	Age attending CiF clinic			
	18 months (n=908)		43 months (n=769)	
	$\beta$ coefficient (95% CI) <sup>a</sup>	P value	$\beta$ coefficient (95% CI) <sup>a</sup>	P value
<b>Core foods (MJ/day)</b>				
All fruit	-0.04 (-0.07, -0.01)	0.012	-0.00 (-0.03, 0.03)	0.785
Meat, fish, poultry and eggs	-0.00 (-0.04, 0.04)	0.931	-0.05 (-0.09, 0.00)	0.063
Potatoes, vegetables and legumes	-0.01 (-0.03, 0.01)	0.393	-0.02 (-0.05, 0.01)	0.229
Vegetables and legumes excluding potatoes	-0.01 (-0.02, 0.01)	0.328	-0.01 (-0.03, 0.01)	0.500
Cereal foods	0.04 (-0.01, 0.09)	0.091	0.00 (-0.07, 0.07)	0.925
Yogurt, cheese and milk	-0.10 (-0.20, -0.01)	0.037	-0.05 (-0.16, 0.07)	0.410
<b>Non-core foods (MJ/day)</b>				
Processed meat, fish, poultry	0.04 (0.01, 0.07)	0.015	0.06 (0.01, 0.11)	0.019
Potatoes cooked in oil and baked beans	0.04 (0.01, 0.08)	0.007	0.02 (-0.03, 0.07)	0.512
Spreads, soups and sauces	0.02 (-0.00, 0.05)	0.050	0.02 (-0.02, 0.05)	0.333
Sweetened processed foods and savoury snacks	0.09 (0.02, 0.17)	0.012	0.10 (-0.01, 0.21)	0.087
<b>Soft drinks (kJ/day)<sup>b</sup></b>				
Fruit juice (unsweetened)	14 (-7, 36), n=342	0.190	-12 (1 (-26, 27), n=331)	0.962
Sugar sweetened	19 (-7, 45), n=296	0.157	-19 (-41, 3), n=447	0.090
Low calorie/diet	4 (0, 8), n=583	0.028	-2 (-5, 1), n=563	0.163

BMI classifications: healthy-/underweight <25 kg/m<sup>2</sup>, overweight/obese ≥25 kg/m<sup>2</sup>.

<sup>a</sup>Adjusted for maternal age, maternal education, parity and Family Adversity Index.


<sup>b</sup>Consumers only.

Reference group: children with healthy-/underweight mothers.

Conversion factor: 1 MJ is equivalent to 239 kcal.

Formatted: Width: 29.7 cm, Height: 21 cm

Formatted: Tab stops: 7.96 cm, Centered + 10.12 cm, Left



Formatted: Width: 21 cm, Height: 29.7 cm

**Figure 1** Mean daily energy intake from core and non-core food groups of ALSPAC-CiF children at 18 months and 43 months of age by maternal BMI classification. Values are means with 95% confidence intervals. Hw/Uw, healthy weight/underweight; Ow/Ob, overweight/obese. At 18 months: Hw/Uw n=706, Ow/Ob n=202. At 48 months: Hw/Uw n=601, Ow/Ob n=168.



## REFERENCES

1. Health and Social Care Information Centre (2018) Overweight and obesity in adults and children. *Health Survey for England 2018* <https://files.digital.nhs.uk/52/FD7E18/HSE18-Adult-Child-Obesity-rep.pdf> (accessed 11 February 2021)
2. Manu R. Obesity and cardiovascular risk in children and adolescents. *Indian Journal of Endocrinology and Metabolism*. doi: 10.4103/2230-8210.91176.
3. Bhadoria A, Sahoo K, Sahoo B *et al.* (2015) Childhood obesity: Causes and consequences. *Journal of Family Medicine and Primary Care* 4, 187.
4. Roberts C, Steer T, Maplethorpe N *et al.* (2018) National Diet and Nutrition Survey. Results from Years 7-8 (combined) of the Rolling Programme (2014/15 to 2015/16): Public Health England.
5. Health Survey for England 2018 (2018) Summary of key findings. <https://digital.nhs.uk/data-and-information/publications/statistical/health-survey-for-england/2018#summary> (accessed 11 February 2020)
6. Simmonds M, Llewellyn A, Owen CG *et al.* (2016) Predicting adult obesity from childhood obesity: a systematic review and meta-analysis. *Obes Rev* 17, 95-107.
7. Whitaker KL, Jarvis MJ, Beeken RJ *et al.* (2010) Comparing maternal and paternal intergenerational transmission of obesity risk in a large population-based sample. *American Journal of Clinical Nutrition* 91, 1560-1567.
8. Thompson AL (2013) Intergenerational impact of maternal obesity and postnatal feeding practices on pediatric obesity. *Nutrition Reviews* 71, S55-S61.
9. Donath S & Amir Lisa H. A systematic review of maternal obesity and breastfeeding intention, initiation and duration. *BMC Pregnancy and Childbirth*. doi: 10.1186/1471-2393-7-9.
10. Emmett PM & Jones LR (2015) Diet, growth, and obesity development throughout childhood in the Avon Longitudinal Study of Parents and Children. *Nutrition reviews* 73, 175-206.
11. Boyd A, Golding J, Macleod J *et al.* (2013) Cohort Profile: The 'Children of the 90s'—the index offspring of the Avon Longitudinal Study of Parents and Children. *International Journal of Epidemiology* 42, 111-127.
12. Fraser A, Macdonald-Wallis C, Tilling K *et al.* (2013) Cohort Profile: the Avon Longitudinal Study of Parents and Children: ALSPAC mothers cohort. *International journal of epidemiology* 42, 97-110.
13. Pauline ME, Victoria LC, David H *et al.* Sources of Vitamin A in the Diets of Pre-School Children in the Avon Longitudinal Study of Parents and Children (ALSPAC). *Nutrients*. doi: 10.3390/nu5051609.
14. World Health Organization (2019) Body mass index - BMI. <http://www.euro.who.int/en/health-topics/disease-prevention/nutrition/a-healthy-lifestyle/body-mass-index-bmi> (accessed 5 September 2019)
15. Roberts DCK (1998) The Australian Guide to Healthy Eating. *Nutrition Bulletin* 23, 224-225.
16. Crawley E, Hughes R, Northstone K *et al.* (2012) Chronic disabling fatigue at age 13 and association with family adversity. *Pediatrics* 130, X23.
17. Cole TJ, Freeman JV Preece MA (1995) Body mass index reference curves for the UK, 1990. *Archives of disease in childhood* 73, 25-29.
18. Dinsdale H, Ridler C Ells LJ (2011) A simple guide to classifying body mass index in children. Oxford: National Obesity Observatory.
19. Nicklas TA, Yang SJ, Baranowski T *et al.* (2003) Eating patterns and obesity in children. The Bogalusa Heart Study. *American journal of preventive medicine* 25, 9-16.

20. Piernas C & Popkin BM (2010) Trends in snacking among U.S. children. *Health affairs (Project Hope)* 29, 398-404.
21. Garden FL, Marks GB, Almqvist C *et al.* (2011) Infant and early childhood dietary predictors of overweight at age 8 years in the CAPS population. *European Journal of Clinical Nutrition* 65, 454-462.
22. Ludwig DS, Peterson KE Gortmaker SL (2001) Relation between consumption of sugar-sweetened drinks and childhood obesity: a prospective, observational analysis. *Lancet (London, England)* 357, 505-508.
23. Øverby NC, Lillegaard IT, Johansson L *et al.* (2004) High intake of added sugar among Norwegian children and adolescents. *Public health nutrition* 7, 285-293.
24. Forshee RA & Storey ML (2001) The role of added sugars in the diet quality of children and adolescents. *Journal of the American College of Nutrition* 20, 32-43.
25. Brekke HK, van Odijk J Ludvigsson J (2007) Predictors and dietary consequences of frequent intake of high-sugar, low-nutrient foods in 1-year-old children participating in the ABIS study. *British Journal of Nutrition* 97, 176-181.
26. Health Survey for England (2014) Health Survey for England 2014: Health, social care and lifestyles. Summary of key findings. <https://digital.nhs.uk/data-and-information/publications/statistical/health-survey-for-england/health-survey-for-england-2014> (accessed 11 February 2021)