



Mason, A. E., Saleem, M. A., & Bierzynska, A. (2021). A critical re-analysis of cases of post-transplantation recurrence in genetic nephrotic syndrome. *Pediatric Nephrology*, 36(11), 3757-3769. <https://doi.org/10.1007/s00467-021-05134-4>

Publisher's PDF, also known as Version of record

License (if available):
CC BY

Link to published version (if available):
[10.1007/s00467-021-05134-4](https://doi.org/10.1007/s00467-021-05134-4)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the final published version of the article (version of record). It first appeared online via Springer at <https://doi.org/10.1007/s00467-021-05134-4> .Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available: <http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>



A critical re-analysis of cases of post-transplantation recurrence in genetic nephrotic syndrome

Anna E. Mason¹ · Moin A. Saleem¹ · Agnieszka Bierzyńska¹

Received: 26 November 2020 / Revised: 29 April 2021 / Accepted: 12 May 2021
© The Author(s) 2021

Abstract

Background Genetic defects in podocyte proteins account for up to 30% of steroid-resistant nephrotic syndrome (SRNS) in the paediatric population. Most children with genetic SRNS are resistant to immunosuppression and at high risk of progression to stage 5 chronic kidney disease. Kidney transplantation is often the treatment of choice. The possibility of post-transplantation disease recurrence in genetic SRNS remains controversial, and poses fundamental questions about disease biology.

Methods We critically evaluated the published cases of post-transplantation recurrence in genetic patients, particularly testing ‘mutations’ against the most recent population variant databases, in order to clarify the diagnoses, and compare the clinical courses and responses to therapy.

Results Biallelic pathogenic variants in *NPHS1* leading to a complete absence of nephrin were the most commonly reported and best understood instance of nephrotic syndrome occurring post-transplantation. This is an immune-mediated process driven by antibody production against the novel nephrin protein in the allograft. We also identified a number of plausible reported cases of post-transplantation recurrence involving pathogenic variants in *NPHS2* (8 patients, biallelic), one in *WT1* (monoallelic) and one in *NUP93* (biallelic). However, the mechanism for recurrence in these cases remains unclear. Other instances of recurrence in genetic disease were difficult to interpret due to differing clinical criteria, inclusion of patients without true pathogenic variants or the influence of other factors on renal outcome.

Conclusions Overall, post-transplantation recurrence remains very rare in patients with genetic SRNS. It appears to occur later after transplantation than in other patients and usually responds well to plasmapheresis with a good renal outcome.

Keywords Nephrotic syndrome · Post-transplantation disease recurrence · Genetic

Introduction

Nephrotic syndrome (NS) has an estimated annual incidence of 2–5 per 100,000 children and is characterised by proteinuria, oedema and hypoalbuminaemia. Eighty percent of children are steroid responsive and subsequently have a good long-term prognosis. The remainder are steroid resistant, either from presentation (primary steroid resistance) or following an initial period of steroid sensitivity (secondary steroid resistance). Those with steroid-resistant nephrotic syndrome (SRNS) are at risk of extrarenal complications and 30–40%

develop stage 5 chronic kidney disease (CKD 5) within 10 years of follow-up. Focal segmental glomerulosclerosis (FSGS) is the commonest renal histology in SRNS. The podocyte cell is crucial in FSGS pathogenesis with disruption to the glomerular filtration barrier, via podocyte injury, leading to loss of permselectivity and albuminuria.

Steroid-sensitive nephrotic syndrome (SSNS) and a subset of SRNS cases are considered to be immunologically mediated and a proportion are caused by an unidentified proteinuric circulating factor(s) produced due to T cell dysfunction. Evidence for the presence of a circulating factor includes the post-transplantation recurrence of proteinuria in SRNS patients and the increased glomerular permeability to albumin caused by FSGS patient serum in rat glomeruli in vitro [1, 2]. Possibly the remainder of SRNS cases are due to genetic disorders, leading to structural defects in the slit diaphragm and podocyte cytoskeletal components, although some may be due to other as yet unclear mechanisms. Over 70 genes are

✉ Moin A. Saleem
M.saleem@bristol.ac.uk

¹ Bristol Renal, Translational Health Sciences, Bristol Medical School, University of Bristol, Dorothy Hodgkin Building, Whitson Street, Bristol BS1 3NY, UK

currently associated with NS and have been shown to account for up to 30% of SRNS cases, including the majority of those presenting within the first year of life [3–17].

Those with a genetic defect have a different pathophysiology, clinical course and response to therapy than those with immune-mediated disease. Most are resistant to immunosuppression, progress more rapidly to CKD 5 and have a lower risk of recurrence following transplantation [3, 18]. Kidney transplantation is often the treatment of choice in this group of patients and knowledge regarding the likely post-transplant clinical course is important to counsel families effectively.

Recurrence of nephrotic syndrome after transplantation is common, occurring in 30–50% of FSGS cases after the first transplant and up to 80% at subsequent transplantations [19]. The onset of proteinuria is generally early (median time 14 days) and shows some response to plasmapheresis and intensified immunosuppression, such as cyclophosphamide [19]. However, despite this treatment response, FSGS recurrence continues to have a negative impact on graft survival with significant physical, social and psychological consequences. The pathogenesis of post-transplantation recurrence remains uncertain but is thought to share a common aetiology with idiopathic FSGS, where circulating plasma factor(s) are postulated to act on the podocyte and disrupt glomerular permeability — so called circulating factor disease. Much work has focused on identifying risk factors for recurrence in order to guide transplantation decisions. Factors which influence recurrence risk include age at diagnosis, rate of progression to CKD 5 and original biopsy result [19, 20]. Paediatric patients with secondary steroid resistance have a much higher post-transplantation recurrence risk than those who were steroid resistant from presentation [21]. However, by far the most important factor to take into consideration is the underlying aetiology of disease. In general, patients with a genetic cause of NS do not recur after transplantation due to correction of the underlying defect and transplantation is therefore the treatment of choice [22]. However, there are a few exceptions to this rule which clinicians should be aware of. Most common is post-transplantation nephrotic syndrome occurring in patients with biallelic pathogenic variants in *NPHS1* leading to an absence of nephrin. This occurs due to the production of antibodies against the novel nephrin protein in the allograft so should be considered as ‘anti-nephrin’ antibody disease, rather than disease recurrence [23].

The question of whether recurrence of NS post-transplantation can occur in genetic NS, and whether this recurrence has the same characteristics as circulating factor disease remains controversial, and poses fundamental questions about the biology of recurrent disease. This report will focus on the reported cases of post-transplantation recurrence in patients with Mendelian NS with a particular focus on re-analysing the genetic variants reported, in the light of current population-level data on rare variants that allows us to assign pathogenicity far more accurately than has been historically possible.

Methods

Reported cases of post-transplantation recurrence in genetic NS were identified from the literature using a PubMed search and the authors’ knowledge of the topic. Key terms used in the PubMed search criteria are shown in Fig. 1. Further cases were then identified using citation chasing; analysing the bibliography of references for each paper (backward citation chasing) and through Google Scholar (forward citation chasing).

Each reported case was reviewed by the authors and the genetic variants re-analysed in view of the current population-level data now available for rare variants. The criteria used to identify true pathogenic variants are shown in Fig. 2. Standard clinical criteria for NS recurrence were used [24]. If further genetic or clinical information was required, the corresponding authors were contacted.

Results of a critical review of reported cases of genetic post-transplant recurrence

NPHS1 and the formation of antibodies

Biallelic pathogenic variants in *NPHS1* are the most commonly reported and best understood instance of post-transplantation disease in genetic nephrotic syndrome. *NPHS1* encodes nephrin, a type-1 transmembrane protein found at the podocyte slit diaphragm, and mutations in this gene are responsible for most cases of congenital nephrotic syndrome (CNS). Fin-major c.121_122delCT; p.(Leu41Aspfs*50) and Fin-minor c.3325C>T; p.(Arg1109*) account for the majority of Finnish-type CNS (78% and 16% respectively), but 373 other *NPHS1* likely pathogenic variants have also been identified to date (HGMD® Professional 2020.4). Homozygous Fin-major is a well-known pathogenic variant leading to the complete absence of nephrin in the native kidney. Post-transplantation disease in these patients is an immune-mediated process driven by the production of antibodies against the novel nephrin protein in the allograft. One study of patients with *NPHS1* Finnish-type CNS demonstrated post-transplantation nephrotic syndrome in 25% with a mean time to disease onset of 12 months [23]. All patients with post-transplantation disease had the homozygous Fin-major variant and anti-nephrin antibodies were detected in 60%. Immunosuppressive therapy led to remission in some cases, but loss of function occurred in almost half of allografts. Antibody production can explain the pathogenesis of post-transplantation nephrotic syndrome in *NPHS1*-associated NS, which is most accurately considered as ‘anti-nephrin’ antibody disease rather than true disease recurrence. However, the cause is less clear in the few reported cases of recurrence associated with other genetic defects.

Fig. 1 Key terms used for the PubMed literature search. All currently known genes associated with NS were also used as search terms [3–17]

Key terms used for PubMed literature search		
Nephrotic syndrome	Recurrence / Recurrent	Congenital
Post(-)transplant(ation)	Paediatric / Pediatric	Genetic
Renal / Kidney transplant(ation)	Proteinuria	Mendelian
Steroid(-)resistant / resistance / SRNS		Hereditary
Focal segmental glomerulosclerosis / glomerular sclerosis / FSGS		Familial

NPHS2

Currently, there is little definitive evidence that SRNS patients with defects in genes other than *NPHS1* will experience post-transplantation recurrence. However, there are a few reports that suggest this can occur in association with certain *NPHS2* biallelic pathogenic variants. Pathogenic variants in *NPHS2*, encoding podocin, have an autosomal recessive pattern of inheritance and account for the majority of infantile and childhood-onset SRNS. Podocin, a raft-associated membrane component located at the insertion of the slit diaphragm, maintains the stability of the slit diaphragm network through recruitment of nephrin and CD2AP to microdomains. Podocin null mice have been shown to develop extensive proteinuria, podocyte foot process effacement and interestingly lack nephrin as well as podocin [25]. This slit diaphragm disassembly in the absence of podocin was also shown by Huber et al. through cell transfection studies [26]. More than 200 pathogenic *NPHS2* variants have now been identified (HGMD@ Professional 2020.4) and are associated with a wide spectrum of clinical phenotype. For instance, the p.(Arg138Gln), which is the most common *NPHS2* pathogenic variant, manifests as very early-onset disease [27]. Conversely, the p.(Arg229Gln) non-neutral polymorphism can lead to late-onset NS when found in association with specific pathogenic *NPHS2* variants [28]. Despite the large number of pathogenic variants found in podocin, post-transplantation recurrence has been described in only a few patients with true biallelic pathogenic variants (Table 1, patients 1–8) [18, 27, 29–32, 35]. Other reported cases of recurrence involve patients with single heterozygous variants which, given the autosomal recessive inheritance of *NPHS2*, should not be considered to have genetic NS (Table 2). This will be discussed in more detail later.

Anti-podocin antibodies have never been identified [29, 31]. This may be because podocin consists of a hairpin membrane domain with two intracellular ends but no extracellular domain. It is therefore a ‘hidden’ protein which may not trigger antibody production. In addition, antibodies should only be stimulated following transplantation when the original pathogenic variant led to truncation or absence

of the protein, such as the p.(Arg138*) which results in podocin truncation. However, many pathogenic variants in *NPHS2* are missense which lead to the production of whole protein and the allograft podocin would not be expected to trigger antibody production in these cases. Post-transplantation recurrence in *NPHS2*-associated NS therefore suggests that there must be an alternative aetiology for recurrence. The possible underlying mechanisms for this will be explored in more detail later.

WT1

One case of post-transplantation recurrence in genetic NS has been reported in association with the Wilms tumour suppressor gene 1 (*WT1*) which fulfils our pathogenic variant criteria (Table 1, patient 10). *WT1* encodes a zinc finger transcription factor involved in kidney and gonadal development. *WT1* may affect podocyte structural proteins as it transcriptionally activates the *NPHS1* promoter, leading to upregulation of nephrin mRNA. Pathogenic variants in this gene (autosomal dominant mode of inheritance — one copy of the gene affected) usually result in syndromic forms of nephrotic syndrome, namely Frasier and Denys–Drash syndrome, but have also been associated with isolated SRNS. This patient developed nephrotic syndrome and FSGS on biopsy in association with Frasier syndrome (triad of glomerulopathy, male pseudohermaphroditism and gonadoblastoma) [34]. She had a pathogenic *WT1* splice site variant c.1447+4C>T:p.? (aka IVS9+4C>T), and received a deceased-donor transplant at 11 years of age. Seven days after transplantation, she developed proteinuria, which gradually increased to nephrotic levels. This responded well to plasmapheresis and allograft function remained good after three years of follow-up. There is a second report of recurrence in a patient with a de novo heterozygous c.1399C>T:p.Arg467Trp (aka p.Arg394Trp) pathogenic variant in exon 9 of the *WT1* gene and early-onset nephrotic syndrome associated with Denys–Drash syndrome [37]. However, this patient had evidence of immune-complex glomerulonephritis on biopsy indicating a different pathogenesis of disease (Table 3, patient 19).

Box 1: Autosomal Recessive Genes**Basic assumptions:**

Full age-related penetrance, defined clinical phenotype, 'pathogenic/disease causing variants' implies that an allele changes the phenotype, known genes responsible for similar phenotype have been excluded. Variants are confirmed with Sanger sequencing; either in homozygous or compound heterozygous state (in *trans* – present on the opposite chromosomes); if familial – segregate with affected family members; novel or rare (MAF<1% and not seen in homozygous state in the 'general population' both global and within sub-populations e.g. gnomAD or in the unaffected family member), if missense – amino acid continuously conserved at least up to *Danio rerio* (zebrafish), loss of function is supported by functional *in vitro/in vivo* data or predicted by at least 3 *in silico* prediction tools for missense and at least 4 for splice site variants.

Disease causing variants can be split into 3 categories depending on the supporting evidence available.

Possibly disease causing:

Novel or MAF <1% in global and sub-populations, predicted to affect the protein

Probably disease causing:

Homozygous novel nonsense or frameshift variant

Or novel and affecting the same codon as previously described pathogenic variant

Or a compound heterozygote where one of the variants is already known to be pathogenic (and the second meets criteria for possibly disease causing)

Definitely disease causing:

If meets the above criteria and was previously described as causing SRNS/related phenotype

Box 2: Autosomal Dominant Genes**Basic assumptions:**

Full age-related penetrance, defined clinical phenotype, 'pathogenic/disease causing variant' implies that an allele changes the phenotype, known genes responsible for similar phenotype have been excluded. Variants are confirmed with Sanger sequencing in heterozygous state; if familial – segregates with affected family members; novel or rare MAF of a disease-causing variant in the general population according to <http://cardiodb.org/allelefrequencyapp/>; variant not seen in unaffected family members (unless adult onset is possible and variant found in an unaffected child); if missense – amino acid continuously conserved at least up to *Danio rerio* (zebrafish), loss of function is supported by functional *in vitro/in vivo* data or predicted by at least 3 *in silico* prediction tools for missense and at least 4 for splice site variants

Disease causing/pathogenic variants can be split into 3 categories depending on the supporting evidence available.

Possibly disease causing:

Novel/MAF in 'general' population in accordance with <http://cardiodb.org/allelefrequencyapp/>, predicted to affect the protein

Probably disease causing

Novel nonsense or frameshift or stop

Or novel and affecting the same codon as previously described pathogenic variant

Definitely disease causing:

If meets the above criteria and was previously described as causing SRNS/related phenotype

Fig. 2 Criteria used to identify patients with true pathogenic variants in recessive (Box 1) or dominant (Box 2) genes. MAF, minor allele frequency

Table 1 Reported cases of post-transplantation recurrence of nephrotic syndrome in patients with true pathogenic variants (pathogenicity criteria defined in Fig. 2). Recurrence is deemed 'possible' when it occurred late (> 3 months) and/or there was insufficient information available. Accession numbers: *NPHS2* NM_014625.4, *NUP93* NM_014669.4, *WT1* NM_024426.6. Allele count and MAF (minor allele frequency; gnomAD v2.1.1 The Genome Aggregation Database, <http://gnomad.broadinstitute.org/>). Allele count = allele count/ number of homozygotes/allele number). The MAF is taken from the total/global population. Nephrotic-range proteinuria is defined in children as > 40 mg/m²/h or uPCR > 2000 mg/g (Kidney disease improving global outcomes, KDIGO guidelines). CP cyclophosphamide, CsA cyclosporin A, d days, FSGS focal segmental glomerulosclerosis, FU follow-up post-transplantation, H homozygous, m months, MP methylprednisolone, uPCR urine protein creatinine ratio, PP plasmapheresis, w weeks, y years

Patient	Gene	Pathogenic variant	Allele count	MAF	Age at onset	Age at transplant	Time to recurrence	Proteinuria at recurrence	Histology at recurrence	Likelihood of disease recurrence	Treatment	Outcome	Reference
1	<i>NPHS2</i>	c.413G>A;p.Arg138Gln (H)	163/0/282690	0.0005766	11m	9y	10d	300 mg/m ² /h	–	Probable	PP (15 cycles) CP (2mg/Kg) 60d	Good	Bertelli 2003 [29]
2	<i>NPHS2</i>	c.413G>A;p.Arg138Gln (H)	163/0/282690	0.0005766	23m	4.5y	300d	120 mg/m ² /h	FSGS	Possible	PP (6 cycles) CP (4 pulses 1g)	Good	Bertelli 2003 [29]
3	<i>NPHS2</i>	c.948delT;p.Ala317Leufs*31 (H)	4/0/282410	0.00001416	8m	4.5y	7d	uPCR 2400 mg/g	–	Probable	MP (500mg 3d) Increased CsA	Good	Billing 2004 [30]
4	<i>NPHS2</i>	c.412C>T;p.Arg138* (H)	4/0/251298	0.00001592	2m	3y	4y	uPCR 5500 mg/g	FSGS, mesangial proliferation	Possible	PP (11w) Ramipril 3.75mg/d	Slow reduction in graft function	Becker-Cohen 2007 [31]
5	<i>NPHS2</i>	c.412C>T;p.Arg138* (H)	4/0/251298	0.00001592	–	–	2y	–	FSGS	Possible	–	–	Weber 2004 [27]
6	<i>NPHS2</i>	c.378G>T;p.Lys126Asn c.948delT;p.Ala317Leufs*31 (H)	4/0/282410	0.00001416	0m	6.6y	–	–	Tacrolimus toxicity	Possible	–	–	Ruf 2004 [18]
7	<i>NPHS2</i>	c.419del;p.Gly140Aspfs*41 (H)	2/0/251358	0.000007957	19m	–	–	–	–	Possible	–	–	Caridi 2003 [32]
8	<i>NPHS2</i>	c.538G>A;p.Val180Met c.467dupT;p.Leu156Phefs*11 (H)	3/0/250804 42/0/204742	0.00001196 0.0002051	96m	–	–	–	–	Possible	–	–	Caridi 2003 [32]
9	<i>NUP93</i>	c.1772G>T;p.Gly591Val c.1916T>C;p.Leu639Pro (H)	38/0/281912	0.0001348	'Infantile'	6y	1.7y	uPCR 4460 mg/g	Subtotal fusion of podocytes, early FSGS	Possible	PP Rituximab	Good (3.1y FU)	Seeman 2018 [33]
10	<i>WT1</i>	c.1447+4C>T;p.? (aka IVS9+4C>T)	–	–	6y	11y	7d	> 40 mg/m ² /hr	NA	Probable	PP (9 cycles)	Good (3y FU)	Ghiggeri 2006 [34]

Table 2 Reported cases of post-transplantation recurrence of nephrotic syndrome in patients with single heterozygous variants (in recessive genes), and therefore unlikely true genetic cause. Accession numbers; *CD2AP* NM_012120.3, *NPHS2* NM_014625.4. Allele count and MAF (minor allele frequency; gnomAD v2.1.1 The Genome Aggregation Database, <http://gnomad.broadinstitute.org/>). Allele count = allele count/ number of homozygotes/allele number). The MAF is taken from the total/global population. Nephrotic-range proteinuria is defined in children as > 40 mg/m²/h or uPCR > 2000 mg/g (Kidney disease improving global outcomes, KDIGO guidelines). *ATN* acute tubular necrosis. *CP* cyclophosphamide. *CsA* cyclosporin A. *d* days. *CKD* stage 5 chronic kidney disease. *FSGS* focal segmental glomerulosclerosis. *FU* follow-up post-transplantation. *m* months. *MP* methylprednisolone. *PD* peritoneal dialysis. *uPCR* urine protein creatinine ratio. *PP* plasmapheresis. *w* weeks. *y* years

Patient	Gene	Variant	Allele count	MAF	Age at onset	Age at transplant	Time to recurrence	Proteinuria at recurrence	Histology at recurrence	Treatment	Outcome	Reference
11	<i>NPHS2</i>	c.622G>A;p.Ala208Thr	13/0/251248	0.00005174	3.5y	8.5y	2w	–	ATN, FSGS	PP	Pre-terminal kidney failure (1.5y FU)	Lowik 2008 [36]
	<i>CD2AP</i>	c.1488G>T;p.Met496Ile	32/0/250916	0.000003985								
12	<i>NPHS2</i>	c.59C>T;p.Pro20Leu	363/2/120298	0.003018	52m	133m	20d	240 mg/m ² /hr	FSGS	PP (18 cycles) CP (4 pulses 1g) PP (20 cycles)	CKD 5-PD after 3m Good	Bertelli 2003 Caridi 2003 [29, 32] Bertelli 2003 Caridi 2003 [29, 32] Bertelli 2003 [29]
13	<i>NPHS2</i>	c.59C>T;p.Pro20Leu	363/2/120298	0.003018	117m	180m	10d	120 mg/m ² /hr	–	PP (8 cycles)	2 nd recurrence after 1m	Caridi 2003 [29, 32] Bertelli 2003 [29]
14	<i>NPHS2</i>	c.631T>A;p.Ser211Thr	1/0/251252	0.000003980	366m	512m	25d	200 mg/m ² /hr	–	PP (8 cycles)	Loss of allograft after 4y	Weber 2004 [27]
15	<i>NPHS2</i>	c.976dupA: p.Thr326Asnfs*20	–	–	–	–	18d	–	FSGS (at 1y)	–	–	Weber 2004 [27]
16	<i>NPHS2</i>	c.59C>T;p.Pro20Leu	363/2/120298	0.003018	–	–	3m	'massive'	–	PP Tacrolimus	–	Weber 2004 [27]
17	<i>NPHS2</i>	c.709G>C;p.Glu237Gln	208/2/282220	0.0007370	–	–	18m	–	–	PP IV CsA	–	Weber 2004 [27]

Table 3 Reported cases of post-transplantation recurrence of nephrotic syndrome in genetic patients where there is insufficient clinical evidence to confirm disease recurrence, or other significant contributory factors to the clinical picture. Accession numbers: *ACTN4* NM_004923.6, *NPHS2* NM_014625.4, *WT1* NM_024426.6. Allele count and MAF (minor allele frequency; gnomAD v2.1.1) The Genome Aggregation Database, <http://gnomad.broadinstitute.org/>. Allele count = allele count/ number of homozygotes/allele number). The MAF is taken

from the total/global population. Nephrotic range proteinuria is defined in children as > 40 mg/m²/h or uPCR > 2000 mg/g (Kidney disease improving global outcomes, KDIGO guidelines). *CNI* calcineurin inhibitor. *CsA* cyclosporin A. *d* days. *FSGS* focal segmental glomerulosclerosis. *FU* follow-up post-transplantation. *GN* glomerulonephritis. *m* months. *MPGN* membranoproliferative glomerulonephritis. *uPCR* urine protein creatinine ratio. *PP* plasmapheresis. *SRL* strolimus. *w* weeks. *y* years

Patient	Gene	Pathogenic variant	Allele count	MAF	Age at onset	Age at transplant	Time to recurrence	Proteinuria at recurrence	Histology at recurrence	Treatment	Outcome	Notes	Reference
18	<i>NPHS2</i>	c.413G>A;p.Arg138Gln c.535-1G>T;p.?	163/0/282690	0.0005766	2m	7y	10y	445 mg/m ² /hr (10.7 g/m ² /d)	FSGS, chronic CNI-induced nephrotoxicity	Conversion back to CsA	Good (2.5y FU)	Recurrence followed switch from CsA to SRL	Hocker 2006 [35]
19	<i>WT1</i>	c.1399C>T;p.Arg467Trp (aka p.Arg394Trp)	–	–	4y	5y	4y	uPCR 350 mg/g (35 g/mol)	Immune complex GN with an MPGN pattern	No changes made	Stable uPCR 1840 mg/g (184 g/mol) (8y FU)	Proteinuria not nephrotic range, biopsy suggests different disease pathogene- sis	Neuhaeus 2011 [37]
20	<i>ACTN4</i>	c.175T>C;p.Trp59Arg	–	–	5y	10y	2y	uPCR 1900 mg/g	Non-specific interstitial fibrosis and tubular atrophy. FSGS not excluded	PP	NA	Proteinuria not nephrotic range	Weins 2005 [38]

NUP93

There is also one published case of post-transplantation recurrence in a patient with compound heterozygous pathogenic variant in nucleoporin 93 (*NUP93*). This is a relatively newly identified cause of genetic NS with onset in early childhood. *NUP93* is an essential component of the nuclear pore complex (NPC), which regulates the transport of protein and RNA between the cytoplasm and the nucleoplasm. Knockdown of *NUP93* has been shown to impair NPC assembly, cell migration, proliferation and SMAD signalling pathways. The reported patient had a compound heterozygous variant affecting exons 16 and 18 of *NUP93* (c.1172G>T and c.1916T>C). He developed infantile SRNS, with biopsy-proven FSGS, and received a deceased-donor transplant at 6 years of age (Table 1, patient 9) [33]. Nephrotic range proteinuria (4460 mg/g) occurred 1.7 years post-transplant, triggered by an upper respiratory tract infection, with early signs of FSGS recurrence on biopsy. The proteinuria responded well to plasmapheresis, but became plasmapheresis-dependent. Rituximab was introduced with good effect and the patient remains proteinuria negative 3.1 years after transplant.

Analysis of clinical course of recurrence

Despite aggressive treatment protocols, recurrence remains a significant cause of allograft loss and paediatric patients with FSGS have a lower allograft survival rate when compared to patients with other primary diseases. However, the clinical picture is variable as some patients with recurrent proteinuria maintain adequate kidney function for several years. Post-transplantation recurrence of NS usually occurs early, with a median time to proteinuria of 14 days [19]. It is interesting to note that four patients with genetic NS, meeting our pathogenic variant criteria, suffered post-transplantation recurrence much later than would be expected (Table 1, patients 2, 4, 5 and 9; 300 days, 4 years, 2 years and 1.7 years, respectively). The majority of patients responded quickly to plasmapheresis and/or immunosuppression and had good graft function. This may suggest that recurrence in genetic NS is a different entity to other FSGS recurrence, with a potentially better clinical course and outcome. Further evaluation of the time course, treatment response and long-term outcome will be needed from future reports of recurrence in genetic NS before this theory can be determined.

Discussion

The potential for, and characteristics of, post-transplantation recurrence in genetic NS is currently controversial and has implications both for our understanding of disease biology

and our ability to make transplantation decisions. There is now a greater degree of population-level data available on rare variants allowing us to assign pathogenicity far more accurately than before. In light of this, we have reviewed and summarised all the published cases of recurrence in genetic NS according to genetic and clinical re-appraisal (Tables 1, 2 and 3). This provides a clearer view on the risk and characteristics of recurrence in patients with true genetic NS which will help to guide clinical management.

Finn major in *NPHS1* remains the most well-established cause of post-transplantation nephrotic syndrome due to ‘anti-nephrin’ antibody disease. However, there is limited evidence that patients with other genetic causes of NS will recur following transplantation. Following re-analysis of the genetic variants, we identified cases of plausible post-transplant recurrence in eight patients with true pathogenic variants in *NPHS2*, one in *WT1* and one in *NUP93* that met our pathogenicity criteria. The mechanism for post-transplant disease is clear in *NPHS1*-associated NS, where antibodies are produced against the nephrin protein in the allograft. However, the pathogenesis of recurrence in other forms of genetic NS remains uncertain. No anti-podocin antibodies have been detected in any tested patient, even in patients with truncating variants [29, 31]. This may be due to difficulties in antibody detection techniques. However, several factors suggest that antibodies may not be the cause: (1) there is no evidence of immunoglobulin deposition in the allograft [29, 31], (2) the onset of recurrence is very early [31] and (3) discrete focal FSGS lesions are seen on biopsy whereas auto-antibodies would be expected to produce a more diffuse pattern of injury [29]. It is thought that one or more circulating plasma factors are present which alter glomerular permeability to proteins and cause proteinuria [1]. This factor(s) remains so far unidentified but its production appears to follow T cell dysfunction. Serum from patients with FSGS increases rat glomerular permeability to albumin in vitro [2]. Measuring volume changes of these rat glomeruli on exposure to FSGS serum has been used as an indirect measure of permeability activity (P_{alb}). High P_{alb} values are predictive of post-transplantation recurrence in FSGS [39]. In one study, the risk of recurrence in patients with a positive proteinuric factor was ten times greater than in those where it was undetectable [39]. Carraro et al. identified two patients with *NPHS2*-associated NS, meeting our pathogenic variant criteria, and high P_{alb} values pre-transplant and at the time of recurrence (Table 1, patients 1–2) [40]. In one patient, P_{alb} values directly correlated with proteinuria and decreased progressively with plasmapheresis. The good clinical response to plasmapheresis and/or immunosuppression in post-transplantation recurrence is additional support for the role of a systemic factor in this process. Permeability activity may be determined by the balance of plasma factors that either induce or inhibit permeability. Apolipoproteins have been shown to inhibit permeability induced by FSGS serum

in vitro [41]. Loss of inhibitory serum components in the urine during proteinuria may play an important role in the recurrence of FSGS post-transplantation [42].

Whilst post-transplantation recurrence generally occurs early (median 14 days), it is interesting that four of these 10 genetic NS patients had a much later onset of recurrence than would be expected (300 days, 1.7 years, 2 years and 4 years). The majority of patients also responded quickly to treatment and had good graft function, raising the possibility that recurrence in genetic NS is a different entity than other FSGS recurrence. However, further evaluation of the time course, treatment response and long-term outcome in future reports of recurrence in genetic NS will be needed to explore this further. Whilst late recurrence may be a clue to a different type of disease recurrence in genetic NS, it is also possible that these cases represent chronic allograft nephropathy rather than true recurrence.

Apart from these 10 cases, the other reported instances of recurrence in genetic disease were difficult to interpret due to differing clinical criteria, inclusion of patients without true pathogenic variants or the influence of other factors on renal outcome. Interpreting these as definite evidence for post-transplantation recurrence in genetic NS would overestimate the risk in this patient group. Some studies include patients with single heterozygous variants (Table 2). Pathogenic variants in genes, such as *NPHS2*, with an autosomal recessive pattern of inheritance need a variant on both alleles (biallelic) in order to be disease-causing. Single heterozygous variants (found on only one allele) should therefore not be considered pathogenic. It is possible that the development of proteinuria in carriers of single heterozygous podocin variants is due to multigenic inheritance, with the presence of an additional pathogenic variant which has not yet been discovered. However, it is more likely that these patients have non-genetic NS caused by circulating factor disease and experience post-transplantation recurrence in the same way as

other patients in this group. Clear criteria should be followed to identify and include only patients with true pathogenic variants and give a more accurate risk of post-transplantation recurrence in this group (Fig. 2).

There is also variability in the clinical criteria used to define post-transplantation recurrence. The reported case of post-transplantation recurrence in a child with α -actinin-4 gene (*ACTN4*)-associated NS (autosomal dominant) is one example where the clinical details are unconvincing of true recurrence (Table 3, patient 20) [38]. The affected child had a *de novo* p.(Trp59Arg) heterozygous pathogenic variant and received a living-related donor transplant from his father (normal *ACTN4* genotype) at 10 years of age. He developed recurrence of proteinuria post-transplantation but this was of a low level (urine protein creatinine ratio 1900 mg/g). Two years later, his graft function declined and a biopsy was taken. Although recurrent FSGS could not be excluded, the biopsy was not clearly indicative of recurrence showing non-specific interstitial fibrosis with tubular atrophy. The clinical evidence to confirm true recurrence in this case is limited. No other cases of recurrence in *ACTN4*-associated NS have been reported to date and treating this clinical scenario as evidence that it can occur may be misleading when counselling other families with the same condition.

Some episodes of recurrence reported in the literature may actually represent secondary FSGS due to post-transplant morbidity, such as rejection, reflux nephropathy, uncontrolled hypertension or immune-mediated glomerulonephritis [31]. One example of this is a case of post-transplantation recurrence reported in a patient with a heterozygous pathogenic variant in an autosomal dominant gene – *INF2* (inverted formin 2) [43]. We contacted the lead author who subsequently feels this was secondary FSGS due to renal scarring as a result of multiple post-transplant kidney injuries. Side-effects of immunosuppression may also contribute to the development of proteinuria following transplantation. One case is

Fig. 3 Key information and practical advice for clinicians

Practical Summary for Clinicians

- Post-transplantation disease recurrence is extremely rare in genetic nephrotic syndrome
- If suspected:
 - Consider other causes for disease (rejection, medications, immune-mediated glomerulonephritis, uncontrolled hypertension, chronic allograft nephropathy)
 - Confirm the pathogenicity of the gene variant with a geneticist. A variant previously deemed pathogenic may no longer be considered so as more population-level data becomes available or further investigations are carried out.
 - Only then consider recurrence in genetic disease
- Variants of genes with autosomal recessive inheritance must have a pathogenic variant in both alleles. Single heterozygous variants (found on a single allele) are not pathogenic.
- Disease recurrence requires the *de novo* development of nephrotic range proteinuria without a plausible alternative clinical scenario. Foot process effacement on biopsy and prompt response to intensified immunosuppression (including plasma exchange) are strongly supportive features.

reported of a patient with a pathogenic compound heterozygous *NPHS2* variant (p.Arg138Gln + c.535-1G>T) who developed biopsy-proven FSGS recurrence ten years following transplantation (Table 3, patient 18) [35]. This coincided with a medication change from cyclosporin A to sirolimus. Conversion back to the original regimen led to significant improvement in proteinuria and graft function. It may be that sirolimus was insufficient to suppress the immune activation, thereby inducing proteinuria [44]. Cyclosporin A has been shown to reduce proteinuria in non-immune diseases through several mechanisms and, of particular interest in NS, stabilises podocyte foot processes through increased expression of synaptopodin [45]. An overview of the reported cases of genetic NS with insufficient clinical evidence to support true disease recurrence, or other significant contributing factors, can be seen in Table 3. Including cases such as these, which may not signify true recurrence, may lead to an overestimate of the risk. Clear clinical criteria to define post-transplantation disease recurrence should be used consistently throughout the literature. As a minimum, this should include the de novo development of nephrotic range proteinuria without a plausible alternative clinical scenario. In those cases where there is doubt, foot process effacement on biopsy and prompt response to intensified immunosuppression (including plasma exchange) are strongly supportive features.

In addition to the effect of recipient genotype on the risk of post-transplantation recurrence, several donor-dependent genetic factors have been identified which may influence long-term allograft survival. A form of post-transplant recurrence, not related to circulating factors, is of a living donor (LD) carrying a genetic risk allele in the donor kidney. There has already been much discussion in the literature about the use of LD kidney transplants in patients with genetic NS. In summary, in cases of autosomal recessive inheritance, a parent who was a heterozygous carrier would be accepted as a LD due to the negligible recurrence risk, except for carriers of pathogenic variants in *COL4A* genes [24]. Another potential exception is the podocin p.(Arg229Gln) non-neutral polymorphism which encodes a protein with altered nephrin-binding capability when compared to wild-type podocin [28]. This polymorphism increases the risk of adult-onset FSGS when associated with another pathogenic *NPHS2* variant. In contrast, if there is an identified pathogenic variant in an autosomal dominant gene, or a family history suggestive of dominant inheritance, living-related donation should not be used. The exception would be if donor genetic testing for the identified causative variant could be carried out prior to transplantation. Autosomal dominant pathogenic variants can present with variable penetrance and phenotype, including adult-onset NS. Therefore, even if the donor does not have evidence of disease, LD transplant may increase the risk of NS in the recipient and the donor.

A practical summary for clinicians can be seen in Fig. 3. In conclusion, post-transplantation recurrence remains very rare in patients with genetic NS. Whilst some convincing cases do exist, the wider literature should be interpreted with caution due to the differing genetic or clinical criteria, or the influence of other factors on renal outcome. Clear criteria should be followed to identify and include only patients with true pathogenic variants and give a more accurate risk of post-transplantation recurrence in this group. Post-transplantation recurrence in genetic NS appears to occur later, and have better graft outcomes, than in other patients and this warrants further investigation.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s00467-021-05134-4>.

Author contribution The literature search was performed by Anna E Mason and Agnieszka Bierzynska. Genetic analysis was performed by Agnieszka Bierzynska. Clinical expertise was provided by Moin A Saleem. The first draft of the manuscript was written by Anna E Mason and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

Funding This research was supported by the Elizabeth Blackwell Institute, University of Bristol, and funded in part by the Wellcome Trust. For the purpose of Open Access, the author has applied a CC BY public copyright licence to any Author Accepted Manuscript version arising from this submission. A.B is funded by Kidney Research UK (Post-doctoral Fellowship). This work was also supported by the Wellcome Trust Institutional Strategic Support Fund.

Data availability Data sharing is not applicable to this article as no datasets were generated during the current study.

Declarations

Ethics approval Ethics approval is not applicable to this study as it is a re-analysis of previously published literature.

Consent to participate Participant consent is not applicable to this article as all data was taken from previously published literature.

Conflict of interest The authors declare no competing interests.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s00467-021-05134-4>.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Savin VJ, Sharma R, Lovell HB, Welling DJ (1992) Measurement of albumin reflection coefficient with isolated rat glomeruli. *J Am Soc Nephrol* 3:1260–1269
- Savin VJ, Sharma R, Sharma M, McCarthy ET, Swan SK, Ellis E, Lovell H, Warady B, Gunwar S, Chonko AM, Artero M, Vincenti F (1996) Circulating factor associated with increased glomerular permeability to albumin in recurrent focal segmental glomerulosclerosis. *N Engl J Med* 334:878–883. <https://doi.org/10.1056/nejm199604043341402>
- Bierzynska A, McCarthy HJ, Soderquest K, Sen ES, Colby E, Ding WY, Nabhan MM, Kerecuk L, Hegde S, Hughes D, Marks S, Feather S, Jones C, Webb NJ, Ognjanovic M, Christian M, Gilbert RD, Sinha MD, Lord GM, Simpson M, Koziell AB, Welsh GI, Saleem MA (2017) Genomic and clinical profiling of a national nephrotic syndrome cohort advocates a precision medicine approach to disease management. *Kidney Int* 91:937–947. <https://doi.org/10.1016/j.kint.2016.10.013>
- Bierzynska A, Soderquest K, Dean P, Colby E, Rollason R, Jones C, Inward CD, McCarthy HJ, Simpson MA, Lord GM, Williams M, Welsh GI, Koziell AB, Saleem MA (2017) MAGI2 mutations cause congenital nephrotic syndrome. *J Am Soc Nephrol* 28:1614–1621. <https://doi.org/10.1681/asn.2016040387>
- Gee HY, Sadowski CE, Aggarwal PK, Porath JD, Yakulov TA, Schueler M, Lovric S, Ashraf S, Braun DA, Halbritter J, Fang H, Airik R, Vega-Warner V, Cho KJ, Chan TA, Morris LG, French-Constant C, Allen N, McNeill H, Buscher R, Kyrieleis H, Wallot M, Gaspert A, Kistler T, Milford DV, Saleem MA, Keng WT, Alexander SI, Valentini RP, Licht C, Teh JC, Bogdanovic R, Koziell A, Bierzynska A, Soliman NA, Otto EA, Lifton RP, Holzman LB, Sibinga NE, Walz G, Tufro A, Hildebrandt F (2016) FAT1 mutations cause a glomerulotubular nephropathy. *Nat Commun* 7:10822. <https://doi.org/10.1038/ncomms10822>
- Prasad R, Hadjidemetriou I, Maharaj A, Meimaridou E, Buonocore F, Saleem M, Hurcombe J, Bierzynska A, Barbagelata E, Bergada I, Cassinelli H, Das U, Krone R, Hacihamdioglu B, Sari E, Yesilkaya E, Storr HL, Clemente M, Fernandez-Cancio M, Camats N, Ram N, Achermann JC, Van Veldhoven PP, Guasti L, Braslavsky D, Guran T, Metherell LA (2017) Sphingosine-1-phosphate lyase mutations cause primary adrenal insufficiency and steroid-resistant nephrotic syndrome. *J Clin Invest* 127:942–953. <https://doi.org/10.1172/jci90171>
- Rao J, Ashraf S, Tan W, van der Ven AT, Gee HY, Braun DA, Feher K, George SP, Esmailniakooshkghazi A, Choi WI, Jobst-Schwan T, Schneider R, Schmidt JM, Widmeier E, Warejko JK, Hermle T, Schapiro D, Lovric S, Shril S, Daga A, Nayir A, Shenoy M, Tse Y, Bald M, Helmchen U, Mir S, Berdeli A, Kari JA, El Desoky S, Soliman NA, Bagga A, Mane S, Jairajpuri MA, Lifton RP, Khurana S, Martins JC, Hildebrandt F (2017) Advillin acts upstream of phospholipase C 1 in steroid-resistant nephrotic syndrome. *J Clin Invest* 127:4257–4269. <https://doi.org/10.1172/jci94138>
- Ashraf S, Kudo H, Rao J, Kikuchi A, Widmeier E, Lawson JA, Tan W, Hermle T, Warejko JK, Shril S, Airik M, Jobst-Schwan T, Lovric S, Braun DA, Gee HY, Schapiro D, Majmundar AJ, Sadowski CE, Pabst WL, Daga A, van der Ven AT, Schmidt JM, Low BC, Gupta AB, Tripathi BK, Wong J, Campbell K, Metcalfe K, Schanze D, Niihori T, Kaito H, Nozu K, Tsukaguchi H, Tanaka R, Hamahira K, Kobayashi Y, Takizawa T, Funayama R, Nakayama K, Aoki Y, Kumagai N, Iijima K, Fehrenbach H, Kari JA, El Desoky S, Jalalah S, Bogdanovic R, Stajic N, Zappel H, Rakhmetova A, Wassmer SR, Jungraithmayr T, Strehlau J, Kumar AS, Bagga A, Soliman NA, Mane SM, Kaufman L, Lowy DR, Jairajpuri MA, Lifton RP, Pei Y, Zenker M, Kure S, Hildebrandt F (2018) Mutations in six nephrosis genes delineate a pathogenic pathway amenable to treatment. *Nat Commun* 9:1960. <https://doi.org/10.1038/s41467-018-04193-w>
- Braun DA, Lovric S, Schapiro D, Schneider R, Marquez J, Asif M, Hussain MS, Daga A, Widmeier E, Rao J, Ashraf S, Tan W, Lusk CP, Kolb A, Jobst-Schwan T, Schmidt JM, Hoogstraten CA, Eddy K, Kitzler TM, Shril S, Moawia A, Schrage K, Khayyat AIA, Lawson JA, Gee HY, Warejko JK, Hermle T, Majmundar AJ, Hugo H, Budde B, Motameny S, Altmuller J, Noegel AA, Fathy HM, Gale DP, Waseem SS, Khan A, Kerecuk L, Hashmi S, Mohebbi N, Ettenger R, Serdaroglu E, Alhasan KA, Hashem M, Goncalves S, Ariceta G, Ubetagoyena M, Antonin W, Baig SM, Alkuraya FS, Shen Q, Xu H, Antignac C, Lifton RP, Mane S, Nurnberg P, Khokha MK, Hildebrandt F (2018) Mutations in multiple components of the nuclear pore complex cause nephrotic syndrome. *J Clin Invest* 128:4313–4328. <https://doi.org/10.1172/jci98688>
- Braun DA, Rao J, Mollet G, Schapiro D, Dageron MC, Tan W, Gribouval O, Boyer O, Revy P, Jobst-Schwan T, Schmidt JM, Lawson JA, Schanze D, Ashraf S, Ullmann JFP, Hoogstraten CA, Boddart N, Collinet B, Martin G, Liger D, Lovric S, Furlano M, Guerrero IC, Sanchez-Ferraz O, Hu JF, Boschac AC, Sanquer S, Menten B, Vergult S, De Rocker N, Airik M, Hermle T, Shril S, Widmeier E, Gee HY, Choi WI, Sadowski CE, Pabst WL, Warejko JK, Daga A, Basta T, Matejas V, Scharmann K, Kienast SD, Behnam B, Beeson B, Begtrup A, Bruce M, Ch'ng GS, Lin SP, Chang JH, Chen CH, Cho MT, Gaffney PM, Gipson PE, Hsu CH, Kari JA, Ke YY, Kiraly-Borri C, Lai WM, Lemyre E, Littlejohn RO, Masri A, Moghtaderi M, Nakamura K, Ozaltin F, Praet M, Prasad C, Prytulka A, Roeder ER, Rump P, Schnur RE, Shihara T, Sinha MD, Soliman NA, Soullami K, Sweetser DA, Tsai WH, Tsai JD, Topaloglu R, Vester U, Viskochil DH, Vatanavicharn N, Waxler JL, Wierenga KJ, Wolf MTF, Wong SN, Leidel SA, Truglio G, Dedon PC, Poduri A, Mane S, Lifton RP, Bouchard M, Kannu P, Chitayat D, Magen D, Callewaert B, van Tilbeurgh H, Zenker M, Antignac C, Hildebrandt F (2017) Mutations in KEOPS-complex genes cause nephrotic syndrome with primary microcephaly. *Nat Genet* 49:1529–1538. <https://doi.org/10.1038/ng.3933>
- Braun DA, Warejko JK, Ashraf S, Tan W, Daga A, Schneider R, Hermle T, Jobst-Schwan T, Widmeier E, Majmundar AJ, Nakayama M, Schapiro D, Rao J, Schmidt JM, Hoogstraten CA, Hugo H, Bakkaloglu SA, Kari JA, El Desoky S, Daouk G, Mane S, Lifton RP, Shril S, Hildebrandt F (2019) Genetic variants in the LAMA5 gene in pediatric nephrotic syndrome. *Nephrol Dial Transplant* 34:485–493. <https://doi.org/10.1093/ndt/gfy028>
- Hermle T, Schneider R, Schapiro D, Braun DA, van der Ven AT, Warejko JK, Daga A, Widmeier E, Nakayama M, Jobst-Schwan T, Majmundar AJ, Ashraf S, Rao J, Finn LS, Tasic V, Hernandez JD, Bagga A, Jalalah SM, El Desoky S, Kari JA, Laricchia KM, Lek M, Rehm HL, MacArthur DG, Mane S, Lifton RP, Shril S, Hildebrandt F (2018) GAPVD1 and ANKFY1 mutations implicate RAB5

- regulation in nephrotic syndrome. *J Am Soc Nephrol* 29:2123–2138. <https://doi.org/10.1681/asn.2017121312>
13. Dorval G, Kuzmuk V, Gribouval O, Welsh GI, Bierzynska A, Schmitt A, Miserey-Lenkei S, Koziell A, Haq S, Benmerah A, Mollet G, Boyer O, Saleem MA, Antignac C (2019) TBC1D8B loss-of-function mutations lead to X-linked nephrotic syndrome via defective trafficking pathways. *Am J Hum Genet* 104:348–355. <https://doi.org/10.1016/j.ajhg.2018.12.016>
 14. Maroofian R, Schuele I, Najafi M, Bakey Z, Rad A, Antony D, Habibi H, Schmidts M (2018) Parental whole-exome sequencing enables sialidosis type II diagnosis due to an NEU1 missense mutation as an underlying cause of nephrotic syndrome in the child. *Kidney Int Rep* 3:1454–1463. <https://doi.org/10.1016/j.ekir.2018.07.015>
 15. Mistry K, Ireland JH, Ng RC, Henderson JM, Pollak MR (2007) Novel mutations in NPHP4 in a consanguineous family with histological findings of focal segmental glomerulosclerosis. *Am J Kidney Dis* 50:855–864. <https://doi.org/10.1053/j.ajkd.2007.08.009>
 16. Sato Y, Tsukaguchi H, Morita H, Higasa K, Tran MTN, Hamada M, Usui T, Morito N, Horita S, Hayashi T, Takagi J, Yamaguchi I, Nguyen HT, Harada M, Inui K, Maruta Y, Inoue Y, Koiwa F, Sato H, Matsuda F, Ayabe S, Mizuno S, Sugiyama F, Takahashi S, Yoshimura A (2018) A mutation in transcription factor MAFB causes focal segmental glomerulosclerosis with duane retraction syndrome. *Kidney Int* 94:396–407. <https://doi.org/10.1016/j.kint.2018.02.025>
 17. Solanki AK, Widmeier E, Arif E, Sharma S, Daga A, Srivastava P, Kwon SH, Hugo H, Nakayama M, Mann N, Majmundar AJ, Tan W, Gee HY, Sadowski CE, Rinat C, Becker-Cohen R, Bergmann C, Rosen S, Somers M, Shril S, Huber TB, Mane S, Hildebrandt F, Nihalani D (2019) Mutations in KIRREL1, a slit diaphragm component, cause steroid-resistant nephrotic syndrome. *Kidney Int* 96:883–889. <https://doi.org/10.1016/j.kint.2019.06.016>
 18. Ruf RG, Lichtenberger A, Karle SM, Haas JP, Anacleto FE, Schultheiss M, Zalewski I, Imm A, Ruf EM, Mucha B, Bagga A, Neuhaus T, Fuchshuber A, Bakkaloglu A, Hildebrandt F (2004) Patients with mutations in NPHS2 (podocin) do not respond to standard steroid treatment of nephrotic syndrome. *J Am Soc Nephrol* 15:722–732. <https://doi.org/10.1097/01.asn.0000113552.59155.72>
 19. Weber S, Tonshoff B (2005) Recurrence of focal-segmental glomerulosclerosis in children after renal transplantation: clinical and genetic aspects. *Transplantation* 80(1 Suppl):S128–S134. <https://doi.org/10.1097/01.tp.0000187110.25512.82>
 20. Ponticelli C (2010) Recurrence of focal segmental glomerular sclerosis (FSGS) after renal transplantation. *Nephrol Dial Transplant* 25:25–31. <https://doi.org/10.1093/ndt/gfp538>
 21. Ding WY, Koziell A, McCarthy HJ, Bierzynska A, Bhagavatula MK, Dudley JA, Inward CD, Coward RJ, Tizard J, Reid C, Antignac C, Boyer O, Saleem MA (2014) Initial steroid sensitivity in children with steroid-resistant nephrotic syndrome predicts post-transplant recurrence. *J Am Soc Nephrol* 25:1342–1348. <https://doi.org/10.1681/asn.2013080852>
 22. Morello W, Puvinathan S, Puccio G, Ghiggeri GM, Dello Strologo L, Peruzzi L, Murer L, Cioni M, Guzzo I, Cocchi E, Benetti E, Testa S, Ghio L, Caridi G, Cardillo M, Torelli R, Montini G (2020) Post-transplant recurrence of steroid resistant nephrotic syndrome in children: the Italian experience. *J Nephrol* 33:849–857. <https://doi.org/10.1007/s40620-019-00660-9>
 23. Patrakka J, Ruotsalainen V, Reponen P, Qvist E, Laine J, Holmberg C, Tryggvason K, Jalanko H (2002) Recurrence of nephrotic syndrome in kidney grafts of patients with congenital nephrotic syndrome of the Finnish type: role of nephrin. *Transplantation* 73:394–403. <https://doi.org/10.1097/00007890-200202150-00013>
 24. Trautmann A, Vivarelli M, Samuel S, Gipson D, Sinha A, Schaefer F, Hui NK, Boyer O, Saleem MA, Feltran L, Müller-Deile J, Becker JU, Cano F, Xu H, Lim YN, Smoyer W, Anochie I, Nakanishi K, Hodson E, Haffner D (2020) IPNA clinical practice recommendations for the diagnosis and management of children with steroid-resistant nephrotic syndrome. *Pediatr Nephrol* 35:1529–1561. <https://doi.org/10.1007/s00467-020-04519-1>
 25. Roselli S, Heidet L, Sich M, Henger A, Kretzler M, Gubler MC, Antignac C (2004) Early glomerular filtration defect and severe renal disease in podocin-deficient mice. *Mol Cell Biol* 24:550–560. <https://doi.org/10.1128/mcb.24.2.550-560.2004>
 26. Huber TB, Simons M, Hartleben B, Sernetz L, Schmidts M, Gundlach E, Saleem MA, Walz G, Benzing T (2003) Molecular basis of the functional podocin-nephrin complex: mutations in the NPHS2 gene disrupt nephrin targeting to lipid raft microdomains. *Hum Mol Genet* 12:3397–3405. <https://doi.org/10.1093/hmg/ddg360>
 27. Weber S, Gribouval O, Esquivel EL, Moriniere V, Tete MJ, Legendre C, Niaudet P, Antignac C (2004) NPHS2 mutation analysis shows genetic heterogeneity of steroid-resistant nephrotic syndrome and low post-transplant recurrence. *Kidney Int* 66:571–579. <https://doi.org/10.1111/j.1523-1755.2004.00776.x>
 28. Tsukaguchi H, Sudhakar A, Le TC, Nguyen T, Yao J, Schwimmer JA, Schachter AD, Poch E, Abreu PF, Appel GB, Pereira AB, Kalluri R, Pollak MR (2002) NPHS2 mutations in late-onset focal segmental glomerulosclerosis: R229Q is a common disease-associated allele. *J Clin Invest* 110:1659–1666. <https://doi.org/10.1172/jci16242>
 29. Bertelli R, Ginevri F, Caridi G, Dagnino M, Sandrini S, Di Duca M, Emma F, Sanna-Cherchi S, Scolari F, Neri TM, Murer L, Massella L, Basile G, Rizzoni G, Perfumo F, Ghiggeri GM (2003) Recurrence of focal segmental glomerulosclerosis after renal transplantation in patients with mutations of podocin. *Am J Kidney Dis* 41:1314–1321. [https://doi.org/10.1016/s0272-6386\(03\)00364-0](https://doi.org/10.1016/s0272-6386(03)00364-0)
 30. Billing H, Muller D, Ruf R, Lichtenberger A, Hildebrandt F, August C, Querfeld U, Haffner D (2004) NPHS2 mutation associated with recurrence of proteinuria after transplantation. *Pediatr Nephrol* 19:561–564. <https://doi.org/10.1007/s00467-003-1408-6>
 31. Becker-Cohen R, Bruschi M, Rinat C, Feinstein S, Zennaro C, Ghiggeri GM, Frishberg Y (2007) Recurrent nephrotic syndrome in homozygous truncating NPHS2 mutation is not due to anti-podocin antibodies. *Am J Transplant* 7:256–260. <https://doi.org/10.1111/j.1600-6143.2006.01605.x>
 32. Caridi G, Bertelli R, Di Duca M, Dagnino M, Emma F, Onetti Muda A, Scolari F, Miglietti N, Mazzucco G, Murer L, Carrea A, Massella L, Rizzoni G, Perfumo F, Ghiggeri GM (2003) Broadening the spectrum of diseases related to podocin mutations. *J Am Soc Nephrol* 14:1278–1286. <https://doi.org/10.1097/01.asn.0000060578.79050.e0>
 33. Seeman T, Vondrak K (2018) First report of recurrent nephrotic syndrome after kidney transplantation in a patient with NUP93 gene mutations: a case report. *Transplant Proc* 50:3954–3956. <https://doi.org/10.1016/j.transproceed.2018.07.010>
 34. Ghiggeri GM, Aucella F, Caridi G, Bisceglia L, Ghio L, Gigante M, Perfumo F, Carraro M, Gesualdo L (2006) Posttransplant recurrence of proteinuria in a case of focal segmental glomerulosclerosis associated with WT1 mutation. *Am J Transplant* 6:2208–2211. <https://doi.org/10.1111/j.1600-6143.2006.01418.x>
 35. Hocker B, Knuppel T, Waldherr R, Schaefer F, Weber S, Tonshoff B (2006) Recurrence of proteinuria 10 years post-transplant in NPHS2-associated focal segmental glomerulosclerosis after conversion from cyclosporin A to sirolimus. *Pediatr Nephrol* 21:1476–1479. <https://doi.org/10.1007/s00467-006-0148-9>
 36. Lowik M, Levtschenko E, Westra D, Groenen P, Steenbergen E, Weening J, Lilien M, Monnens L, van den Heuvel L (2008) Bigenic heterozygosity and the development of steroid-resistant

- focal segmental glomerulosclerosis. *Nephrol Dial Transplant* 23: 3146–3151. <https://doi.org/10.1093/ndt/gfn208>
37. Neuhaus TJ, Arnold W, Gaspert A, Hopfer H, Fischer A (2011) Recurrence of membranoproliferative glomerulonephritis after renal transplantation in Denys-Drash. *Pediatr Nephrol* 26:317–322. <https://doi.org/10.1007/s00467-010-1669-9>
 38. Weins A, Kenlan P, Herbert S, Le TC, Villegas I, Kaplan BS, Appel GB, Pollak MR (2005) Mutational and biological analysis of alpha-actinin-4 in focal segmental glomerulosclerosis. *J Am Soc Nephrol* 16:3694–3701. <https://doi.org/10.1681/asn.2005070706>
 39. Dall'Amico R, Ghiggeri G, Carraro M, Artero M, Ghio L, Zamorani E, Zennaro C, Basile G, Montini G, Rivabella L, Cardillo M, Scalamogna M, Ginevri F (1999) Prediction and treatment of recurrent focal segmental glomerulosclerosis after renal transplantation in children. *Am J Kidney Dis* 34:1048–1055. [https://doi.org/10.1016/s0272-6386\(99\)70010-7](https://doi.org/10.1016/s0272-6386(99)70010-7)
 40. Carraro M, Caridi G, Bruschi M, Artero M, Bertelli R, Zennaro C, Musante L, Candiano G, Perfumo F, Ghiggeri GM (2002) Serum glomerular permeability activity in patients with podocin mutations (NPHS2) and steroid-resistant nephrotic syndrome. *J Am Soc Nephrol* 13:1946–1952. <https://doi.org/10.1097/01.asn.0000016445.29513.ab>
 41. Candiano G, Musante L, Carraro M, Faccini L, Campanacci L, Zennaro C, Artero M, Ginevri F, Perfumo F, Gusmano R, Ghiggeri GM (2001) Apolipoproteins prevent glomerular albumin permeability induced in vitro by serum from patients with focal segmental glomerulosclerosis. *J Am Soc Nephrol* 12:143–150
 42. Coward RJ, Foster RR, Patton D, Ni L, Lennon R, Bates DO, Harper SJ, Mathieson PW, Saleem MA (2005) Nephrotic plasma alters slit diaphragm-dependent signaling and translocates nephrin, Podocin, and CD2 associated protein in cultured human podocytes. *J Am Soc Nephrol* 16:629–637. <https://doi.org/10.1681/asn.2004030172>
 43. Gbadegesin RA, Lavin PJ, Hall G, Bartkowiak B, Homstad A, Jiang R, Wu G, Byrd A, Lynn K, Wolfish N, Ottati C, Stevens P, Howell D, Conlon P, Winn MP (2012) Inverted formin 2 mutations with variable expression in patients with sporadic and hereditary focal and segmental glomerulosclerosis. *Kidney Int* 81:94–99. <https://doi.org/10.1038/ki.2011.297>
 44. Skhiri H, Morelon E, Noel LH, Mamzer-Bruneel MF, Legendre C, Peraldi MN, Kreis H (2005) Unusual post-transplantation recurrence of focal segmental glomerulosclerosis which resolved with cyclosporine but not with sirolimus. *Transpl Int* 18:458–460. <https://doi.org/10.1111/j.1432-2277.2004.00054.x>
 45. Franz S, Faul C, Mundel P (2005) Cyclosporin protects podocyte stress fibers through stabilization of synaptopodin protein expression. *J Am Soc Nephrol* 16:109A

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.