



Williams, A., Lampasona, V., Michael, S., Mueller, P., Pittman, D., Winter, W., Akolkar, B., Wyatt, R., Brigatti, C., Krause, S., & Achenbach, P. (2015). Detection of antibodies directed to the N-terminal region of GAD is dependent on assay format and contributes to differences in the specificity of GAD autoantibody assays for type 1 diabetes. *Diabetes*, 64(9), 3239-3246. <https://doi.org/10.2337/db14-1693>

Peer reviewed version

Link to published version (if available):  
[10.2337/db14-1693](https://doi.org/10.2337/db14-1693)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

## Participating Laboratories and contacts for the DASP 2009 GADA workshop

P. Achenbach, A.-G. Ziegler, Diabetes Research Group of the TU Munich, Munich, Germany; J. Furmaniak, FIRS Laboratories RSR Limited, Cardiff, UK; Vinod Gaur, S.M. Marcovina, Northwest Lipid Metabolism and Diabetes Research Laboratory, University of Washington, Seattle, WA, USA; S. Gellert, Endocrinology Lab, Centre for Medical Research, Royal Melbourne Hospital, Melbourne, Australia; F. Gorus, J. De Grijse, P. Goubert, Dept of Clinical Chemistry and Radioimmunology, University of Brussels, Brussels, Belgium; W. A. Hagopian, Pacific North West Research Institute, Seattle, WA, USA; D.M. Harlan, J. Lee, Clinical Research Center, NIH, Bethesda, MD, USA; M. I. Hawa, Center for Diabetes and Metabolic Medicine, London, UK; E. Kawasaki, Department of Metabolism/Diabetes & Clinical Nutrition, Nagasaki University, Hospital of Medicine and Dentistry, Nagasaki, Japan; M. Knip, Hospital for Children and Adolescents, University of Helsinki, Helsinki, Finland; V. Lampasona, Laboratorio Autoimmunita-LIDEA, Diagnostica e Ricerca San Raffaele, Milan, Italy; P.W. Mueller, National Diabetes Laboratory, Centers for Disease Control and Prevention, Atlanta, GA, USA; A. Nilsson, F. Vaziri Sani, Diabetes and Celiac Disease Unit, Lund University, Malmö, Sweden; M. Schlosser, H. Kenk, Department of Medical Biochemistry and Molecular Biology, University of Greifswald, Karlsburg, Germany; C. Tiberti, F. Dotta, Department of Clinical Science, Immunoendocrinology, University of Rome "La Sapienza", Rome, Italy; P.A. Torjesen, Hormone Laboratory, Aker University Hospital, Oslo, Norway; M. Probst, T. Pfeiffer, S. Torkler, Euroimmun, Lübeck, Germany; R. Uibo, K. Reimand, Department of Immunology, University of Tartu, Tartu, Estonia; A.J.K. Williams, P.J. Bingley, Diabetes and Metabolism, University of Bristol, Bristol, UK; A. Xu, Z. Zhou, G. Huang, Diabetes Center, Institute of Metabolism and Endocrinology, 2nd Xiangya Hospital, Central South University, The University of Hong Kong, Hongkong, China; L. Yu, Barbara Davies Center for Childhood Diabetes, University of Colorado, Aurora, CO, USA; E. Poskus, S. Valdez, A. Cardoso, LIE, School of Pharmacy and Biochemistry, University of Buenos Aires, Buenos Aires, Argentina; M. Atkinson, C. Wasserfall, University of Florida, Gainesville, USA; N. Schloot, F. Zivehe, German Diabetes Center, Institute for Clinical Diabetology, Dusseldorf, Germany; M.R. Batstra, T. Cieremans, Diagnostic Center SSDZ, Medical Laboratories Department of Medical Immunology, Delft, The Netherlands; R. Veijola, Department of Pediatrics, University of Oulu, Oulu, Finland; A. Pollard, J. Couper, Dept. of Paediatrics, Women's & Children's Hospital, Adelaide, Australia; Michael Eckhard, Diabetes and Endocrinology, UKGM, Giessen, Germany; L. Chatenoud, Immunologie Biologique, Hôpital Necker, University of Paris Descartes, Paris, France; T. Kobayashi, Third Department of Internal Medicine, University of Yamanashi, Yamanashi, Japan; I. Johansson, Division of Pediatrics, Department of Clinical and Experimental Medicine, Linköping University, Linköping, Sweden; D. Pittman, Department of Pathology, University of Florida, Gainesville, FL, USA; G. Houen, Department of Clinical Biochemistry and Immunology, Statens Serum Institute, Copenhagen, Denmark; C. Hampe, Jared Radtke, Metabolism, Endocrinology and Nutrition, Department of Medicine, University of Washington, Seattle, WA, USA; A. Esquerda, Clinical Laboratory, Arnau de Vilanova University Hospital, Lleida, Spain; A. Miettinen, Department of Immunology, HUSLAB, University of Helsinki, Helsinki, Finland; M.G. Baroni, Department of Experimental Medicine, University of Rome "La Sapienza", Rome, Italy; A. Caston-Balderrama, A. Fraser, M. French, Quest Diagnostics, Nichols Institute, San Juan Capistrano, CA, USA; L. Park, Department of Bioengineering, College of Engineering, Hanyang University, Seoul, South Korea; A. Pugliese, Immunogenetics Laboratory, Diabetes Research Institute, University of Miami, Miami, FL,

USA; R. Casamitjana, Biochemistry and Molecular Genetics Department, University of Barcelona, Barcelona, Spain; W. Mlynarski, K. Wyka, Laboratory of Immunopathology and Genetics, Department of Pediatrics, Oncology, Hematology and Diabetology, UMED Medical University of Lodz, Lodz, Poland.

#### Participating Laboratories and contacts for the DASP 2010 GADA workshop

P. Achenbach, A.-G. Ziegler, Diabetes Research Group of the TU Munich, Munich, Germany; J. Furmaniak, FIRS Laboratories RSR Limited, Cardiff, UK; Vinod Gaur, S.M. Marcovina, Northwest Lipid Metabolism and Diabetes Research Laboratory, University of Washington, Seattle, WA, USA; F. Gorus, K. Verhaeghen, P. Goubert, Department of Clinical Chemistry and Radioimmunology, University of Brussels, Brussels, Belgium; M. I. Hawa, Center for Diabetes and Metabolic Medicine, London, UK; W. A. Hagopian, Pacific North West Research Institute, Seattle, WA, USA; M. Knip, Hospital for Children and Adolescents, University of Helsinki, Helsinki, Finland; V. Lampasona, Laboratorio Autoimmunita-LIDEA, Diagnostica e Ricerca San Raffaele, Milan, Italy; P.W. Mueller, C. Mapango, National Diabetes Laboratory, Centers for Disease Control and Prevention, Atlanta, GA, USA; A. Nilsson, Diabetes and Celiac Disease Unit, Lund University, Malmö, Sweden; M. Schlosser, H. Kenk, Department of Medical Biochemistry and Molecular Biology, University of Greifswald, Karlsburg, Germany; C. Tiberti, F. Dotta, Department of Clinical Science, Immunoendocrinology, University of Rome "La Sapienza", Rome, Italy; P.A. Torjesen, Hormone Laboratory, Aker University Hospital, Oslo, Norway; R. Uibo, K. Reimand, Department of Immunology, University of Tartu, Tartu, Estonia; A.J.K. Williams, P.J. Bingley, Diabetes and Metabolism, University of Bristol, Bristol, UK; L. Yu, Barbara Davies Center for Childhood Diabetes, University of Colorado, Aurora, CO, USA; M.R. Batstra, T. Cieremans, Diagnostic Center SSDZ, Medical Laboratories Department of Medical Immunology, Delft, The Netherlands; R. Veijola, M. Knip, Department of Pediatrics, University of Oulu, Oulu, Finland; A. Pollard, J. Couper, Dept. of Paediatrics, Women's & Children's Hospital, Adelaide, Australia; L. Chatenoud, Immunologie Biologique, Hôpital Necker, University of Paris Descartes, Paris, France; T. Kobayashi, Third Department of Internal Medicine, University of Yamanashi, Yamanashi, Japan; I. Johansson, Division of Pediatrics, Department of Clinical and Experimental Medicine, Linköping University, Linköping, Sweden; D. Pittman, Department of Pathology, University of Florida, Gainesville, FL, USA; G. Houen, Y.M. Moreno, K. Lunau-Christensen, Department of Clinical Biochemistry and Immunology, Statens Serum Institute, Copenhagen, Denmark; C. Hampe, Jared Radtke, Metabolism, Endocrinology and Nutrition, Department of Medicine, University of Washington, Seattle, WA, USA; Aureli Esquerda, Clinical Laboratory, Arnau de Vilanova University Hospital, Lleida, Spain; A. Miettinen, Department of Immunology, HUSLAB, University of Helsinki, Helsinki, Finland; B Ekholm, Diabetes Research Laboratory, Department of Clinical Sciences, Lund University, Sweden; L. Castaño, R. Bilbao, Endocrinology and Diabetes Research Group, Hospital de Cruces, Barakaldo, Bizkaia, Spain; M.R. Christie, Diabetes Research Group, King's College London, London, UK; C. Giordano, Biomedical Department of Internal and Specialist Medicine, Section of Endocrinology, Diabetology, and Metabolism, University of Palermo, Palermo, Italy; R. Pujol-Borrell, E. Martínez Cáceres, Immunology Lab, LIRAD-BST-HUGTP, Germans Trias Pujol University Hospital, UAB, Badalona, Spain; W. Mlynarski, K. Wyka, Laboratory of Immunopathology and Genetics, Department of Pediatrics, Oncology, Hematology and

Diabetology, UMED Medical University of Lodz, Lodz, Poland; J. Siftancova, Laboratory of Autoimmune Diseases, Paediatric Department, University Hospital Motol and 2nd Faculty of Medicine, Charles University in Prague, Prague, Czech Republic; T. McDonald, Clinical Chemistry and Immunology, Royal Devon and Exeter Hospital, UK; J. Knezevic Cuca, Immunology of Diabetes Unit, Institute of Clinical Chemistry and Laboratory Medicine, Merkur University Hospital, Zagreb, Croatia; R. Casamitjana, Biochemistry and Molecular Genetics Department, University of Barcelona, Barcelona, Spain; H.-J. Thiesen, Institute for Immunology, STZ für Proteom-Analyse, Rostock, Germany; S. Scheucher, LKH-University Klinikum, Graz, Austria.

#### Participating Laboratories and contacts for the IASP2012 GADA substudy

P. Achenbach, A.-G. Ziegler, Diabetes Research Group of the TU Munich, Munich, Germany; F. Gorus, K. Verhaeghen, P. Goubert, Department of Clinical Chemistry and Radioimmunology, University of Brussels, Brussels, Belgium; M. Knip, Hospital for Children and Adolescents, University of Helsinki, Helsinki, Finland; M. Schlosser, H. Kenk, Department of Medical Biochemistry and Molecular Biology, University of Greifswald, Karlsburg, Germany; R. Uibo, K. Reimand, Department of Immunology, University of Tartu, Tartu, Estonia; A.J.K. Williams, P.J. Bingley, Diabetes and Metabolism, University of Bristol, Bristol, UK; M.R. Christie, Diabetes Research Group, King's College London, London, UK; A. Xu, Z. Zhou, G. Huang, Diabetes Center, Institute of Metabolism and Endocrinology, 2nd Xiangya Hospital, Central South University, The University of Hong Kong, Hongkong, China; T Yang, Department of Endocrinology, First Affiliated Hospital of Nanjing Medical University, Nanjing, China; E. Kawasaki, Department of Metabolism/Diabetes & Clinical Nutrition, Nagasaki University, Hospital of Medicine and Dentistry, Nagasaki, Japan.