



Liu, A. G. S. C., Matthews, J., Menon, L., Mcllroy, D., & Brasier, M. (2015). The arrangement of possible muscle fibres in the Ediacaran taxon *Hootia quadriformis*. *Proceedings of the Royal Society B: Biological Sciences*, 282(1803).
<https://doi.org/10.1098/rspb.2014.2949>

Peer reviewed version

Link to published version (if available):
[10.1098/rspb.2014.2949](https://doi.org/10.1098/rspb.2014.2949)

[Link to publication record in Explore Bristol Research](#)
PDF-document

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

PROCEEDINGS B

The arrangement of possible muscle fibres in the Ediacaran taxon *Haootia quadriformis*

Journal:	<i>Proceedings B</i>
Manuscript ID:	RSPB-2014-2949.R1
Article Type:	Invited Reply
Date Submitted by the Author:	n/a
Complete List of Authors:	Liu, Alexander; University of Bristol, School of Earth Sciences Matthews, Jack; University of Oxford, Department of Earth Sciences Menon, Latha; University of Oxford, Department of Earth Sciences McIlroy, Duncan; Memorial University of Newfoundland, Department of Earth Sciences Brasier, Martin; University of Oxford, Department of Earth Sciences; Memorial University of Newfoundland, Earth Sciences
Subject:	Evolution < BIOLOGY, Developmental biology < BIOLOGY, Palaeontology < BIOLOGY
Keywords:	Ediacaran, Cnidarian, Muscle, Newfoundland
Proceedings B category:	Palaeontology

SCHOLARONE™
Manuscripts

1 **Title: The arrangement of possible muscle fibres in the**
2 **Ediacaran taxon *Hootia quadriformis*.**

3

4 **Authors:** Alexander G. Liu^{1, *}, Jack J. Matthews², Latha R. Menon², Duncan McIlroy³ and
5 Martin D. Brasier^{2, 3, §}

6 ¹*School of Earth Sciences, University of Bristol, Life Sciences Building, 24 Tyndall Avenue,*
7 *Bristol, BS8 1TQ, U.K.*

8 ²*Department of Earth Sciences, University of Oxford, South Parks Road, Oxford, OX1 3AN,*
9 *U.K.*

10 ³*Department of Earth Sciences, Memorial University of Newfoundland, 300 Prince Philip*
11 *Drive, St. John's, NL, A1B 3X5, Canada*

12 § *Deceased*

13 *Corresponding Author. Email: alex.liu@bristol.ac.uk

14

15 **Keywords:** Ediacaran; cnidarian; muscle; Newfoundland

16

17 *Hootia quadriformis* from Newfoundland, Canada, is one of the most unusual impressions
18 of a soft-bodied macro-organism yet described from the late Ediacaran Period. Interpreted as
19 a metazoan of cnidarian grade [1], the body impression of *H. quadriformis* possesses features
20 interpreted as fibrous structures that represent possible evidence for muscular tissue.

21 Evidence both in support of and against a relationship between *H. quadriformis* and the
22 Staurozoa, one of the cnidarian groups to which *Hootia* was compared in Liu et al. [1], is

23 outlined by Miranda et al. [2]. Our intention in our original paper was to illustrate the
24 staurozoan body plan for comparative purposes, rather than suggest homology or direct
25 ancestry. Nevertheless, fresh insights from workers with expertise in the biology of extant
26 cnidarians are welcomed.

27 We are pleased that the main points of our paper find support from the biological
28 community: that recently discovered *Haootia quadriformis* likely preserves the impressions
29 of muscle fibres in a macrofossil characterized by tetra-radial symmetry, and that these
30 structures are consistent with a cnidarian body plan [2]. Histological images provided by
31 Miranda et al. [2], plus their accompanying discussion, present a clear picture of muscular
32 arrangements within modern stauromedusae. We concur that *Haootia* shows both similarities
33 and differences with respect to stauromedusans.

34 Miranda et al. [2] suggest that the organisation of musculature described by us (see
35 [1], fig. 3b) within the body of *Haootia* appears inconsistent with that observed in modern
36 staurozoan taxa. While the argument for the presence of a radial (longitudinal) muscle
37 arrangement is well reasoned (see [2], fig 1s), a revisiting of the type material to determine
38 the orientation of fibrous structures in *Haootia* leads us to conclude that they are essentially
39 as published in our original paper. The paratype specimen (see [1], fig. 1f) has fibres from
40 potential coronal muscles that extend in arcs almost to the base of the calyx, with no clear
41 preservation of radial fibres running perpendicular to these. We would argue that this
42 phenomenon contradicts the suggestion that coronal muscle may have been restricted to the
43 margins of the *Haootia* body [2].

44 The arrangement of musculature inferred by Liu et al. [1] does not preclude the
45 presence of additional radial muscle bundles within the body of the organism. The holotype
46 of *H. quadriformis* contains evidence for superposition of fibrous structures. We can clearly
47 discern that the body sheet/calyx drapes parallel fibres observed in the stalk/peduncle, and

48 that subsidiary branches appear to extend beneath the body (Fig. 1). In such cases, the fibres
49 beneath the calyx are not expressed in the fossilised impression, and are inferred to lie
50 beyond the plane of preservation. It is therefore possible that both coronal and radial
51 musculature were present in the calyx of *Haootia* (just as both are present in the arms of
52 modern stauromedusae [2]), but only the outermost set is recorded in the cast of the body
53 tissues so beautifully preserved in the *Haootia* impression.

54 If one were to adopt a stauromedusan analogue for *Haootia*, the inferred coronal
55 musculature would be in the ‘calyx’ rather than the radial disc. This proposed muscular
56 arrangement (see [1], fig. 3b) would argue against a free-swimming mode of life, and adds
57 support to a benthic mode of life for *Haootia*. We therefore take great interest in the
58 suggestion that this arrangement of muscles is consistent with an organism capable of
59 producing powerful body contractions, potentially involved in a pulsing feeding strategy [2].
60 We must keep in mind that some, or maybe most, Ediacaran body plans and feeding
61 strategies may have been specifically adapted to Ediacaran conditions. Possible behavioural
62 ecologies such as surface detritus feeding, which would be feasible for an organism with so
63 flexible a body, still await analysis.

64 The hypothetical models of muscle fibre arrangement proposed by ourselves [1] and
65 Miranda et al. [2] both require further testing against the fossil material. The remarkably
66 preserved holotype specimen of *Haootia* has now been collected (under permits issued by the
67 Department of Tourism, Culture and Recreation, Government of Newfoundland and
68 Labrador), and is housed at The Rooms Provincial Museum, St. John’s, Newfoundland
69 (specimen NFM F-994). While the fidelity of preservation restricts elements of the
70 interpretation of this key fossil, the opportunity to apply controlled lighting conditions to the
71 specimen following its removal from the outcrop offers the potential for further analysis (e.g.
72 Fig. 1). We are confident that well-informed discussion of analogue taxa such as that offered

73 by Miranda et al. will lead to a more complete understanding of *Haootia*, and its evolutionary
74 significance. Similarly detailed consideration of the physiology of other extant cnidarian
75 clades would be greatly beneficial for palaeontologists working on candidate early cnidarians.

76 We would like to add in support of our discussion of *Haootia* that strata of similar age
77 in Newfoundland also contain rare fossilized surface locomotion trails that are considered to
78 have formed by contractile activities associated with a cnidarian body plan [3–6]. We have
79 hitherto found no evidence that the basal disc of *Haootia* could contract or move, but having
80 two independent lines of evidence for the presence of organisms with contractile tissues in
81 the Ediacaran makes a more compelling case for the presence of cnidarian-grade organisms
82 [1]. While we recognise the potential importance of a fossil cnidarian ancestor from
83 somewhat well-dated strata for understanding the phylogeny of the Cnidaria, we would
84 caution against the uncritical extension of such interpretations to other Ediacaran taxa. The
85 preservation and morphology of many late Ediacaran macrofossils makes them difficult to
86 interpret, and further work is required before we can be confident in our understanding of
87 these ancient and evolutionarily important organisms.

88

89 **Acknowledgement**

90 We are grateful to two anonymous reviewers, and to NSERC (to D.M.) and the Natural
91 Environment Research Council (grant no. NE/L011409/1 to A.G.L., and NE/J5000045/1 to
92 J.J.M.) for financial support. The staff of The Rooms Provincial Museum and the
93 Newfoundland and Labrador Department of Tourism, Culture and Recreation are thanked for
94 their role in the successful salvage and curation of the holotype specimen. Additional images
95 can be found here: doi:10.5061/dryad.8m2q8. We dedicate this manuscript to our co-author,
96 mentor and friend Martin Brasier, who contributed so much to this field.

97

98 **References**

- 99 1. Liu AG, Matthews JJ, Menon LR, McIlroy D, Brasier MD. 2014 *Haootia quadriformis*
100 n. gen., n. sp., interpreted as a muscular cnidarian impression from the late Ediacaran
101 Period (~560 Ma). *Proc. R. Soc. B* **281**. (doi: 10.1098/rspb.2014.1202).
- 102 2. Miranda LS, Collins AG, Marques AC. 2014 Is *Haootia quadriformis* related to extant
103 Staurozoa (Cnidaria)? Evidence from the muscular system reconsidered. *Proc. R. Soc.*
104 *B. (This volume)*
- 105 3. Liu AG, McIlroy D. 2015. Horizontal surface traces from the Fermeuse Formation,
106 Ferryland (Newfoundland, Canada), and their place within the late Ediacaran
107 ichnological revolution. In: McIlroy D. (Ed.) ICHNOLOGY: Publications arising from
108 ICHNIA III. *Geological Association of Canada, Miscellaneous Publication* **9**, 141–156.
- 109 4. Liu AG, McIlroy D, Brasier MD. 2010 First evidence for locomotion in the Ediacara
110 biota from the 565Ma Mistaken Point Formation, Newfoundland. *Geology* **38**, 123–
111 126. (doi: 10.1130/G30368.1).
- 112 5. Liu AG, McIlroy D, Brasier MD, Matthews JJ. 2014 Confirming the metazoan
113 character of a 565 Ma trace-fossil assemblage from Mistaken Point, Newfoundland.
114 *Palaios* **29**, 420–430. (doi: 10.2110/palo.2014.011).
- 115 6. Menon LR, McIlroy D, Brasier MD. 2013 Evidence for Cnidaria-like behavior in ca.
116 560 Ma Ediacaran *Aspidella*. *Geology* **41**, 895–898. (doi:10.1130/G34424.1).

117

118 **Figure Caption**

119 **Figure 1.** The disc, stalk and body of the *Haootia quadriformis* holotype. Image taken at The
120 Rooms Provincial Museum, St. John's. Note that although the outline of the stalk/peduncle
121 can be seen beneath the body/calyx, the longitudinal fibres running along its length cannot be

122 easily traced (white arrow). Similarly, fibres radiating from the smallest subsidiary branches
123 are typically not seen to continue beneath the body (e.g. black arrow). It is thus plausible that
124 additional muscle bundles, perhaps in different arrangements to those preserved, were present
125 in the original *Haootia* organism, but do not lie in the plane of preservation. Scale bar = 10
126 mm.



The disc, stalk and body of the *Hootia quadriformis* holotype. Image taken at The Rooms Provincial Museum, St. John's. Note that although the outline of the stalk/peduncle can be seen beneath the body/calyx, the longitudinal fibres running along its length cannot be easily traced (white arrow). Similarly, fibres radiating from the smallest subsidiary branches are typically not seen to continue beneath the body (e.g. black arrow). It is thus plausible that additional muscle bundles, perhaps in different arrangements to those preserved, were present in the original *Hootia* organism, but do not lie in the plane of preservation.

Scale bar = 10 mm.

133x88mm (600 x 600 DPI)