



Collin, S. M., Leclercq, B., Twungubumwe, N., Andréoletti, L., Richardier, F-C., & Bertin, E. (2016). Leaf concentrate compared with skimmed milk as nutritional supplementation for HIV-infected children: a randomized controlled trial in Burundi. *Public Health Nutrition*, 19(10), 1904-1912. <https://doi.org/10.1017/S1368980015003456>

Peer reviewed version

Link to published version (if available):
[10.1017/S1368980015003456](https://doi.org/10.1017/S1368980015003456)

[Link to publication record in Explore Bristol Research](#)
PDF-document

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Abstract

Objective: The effectiveness of leaf concentrate powder (LCP) as a nutritional supplement was established in trials conducted among adolescent girls and pregnant women in India. Here we evaluate LCP, compared with skimmed milk powder (SMP), as a supplement for antiretroviral-naïve children living with HIV in a sub-Saharan African country.

Design: Randomized controlled two-arm 6-month trial comparing effects of isoproteic (5g) LCP (10g daily) and SMP (15g daily) on HIV-1 viral load, CD4+ cell count/percentage, weight/height-for-age, general blood parameters, diarrhea, respiratory and HIV-related opportunistic infections.

Setting: Bujumbura and Kirundo, Burundi.

Subjects: 83 HIV+ ART-naïve children age 5-14 years: median (range) CD4 count 716 (361-1690) cells/mm³; log₁₀ HIV-1 viral load 4.39 (1.79-6.00).

Results: LCP was equivalent to SMP in relation to HIV-specific blood parameters, and did not demonstrate superiority over SMP in relation to Hb. Three children in each arm (LCP 7.1% (3/42); SMP 7.3% (3/41)) proceeded to ART because their CD4 counts fell below 350 cells/mm³. Children in the LCP group reported higher levels of appetite and overall health at 6 months. There were no differences in clinical events or any other outcome measures. LCP was less palatable than SMP to the children in this population, but there were few negative perceptions of appearance, texture and taste.

Conclusions: LCP appears to be equivalent to SMP as a nutritional supplement in this population, despite slightly lower palatability. In relation to viral load and CD4 count, equivalence may indicate no effect in either group. Effectiveness relative to no supplementation remains to be determined.

Introduction

Maintaining good nutritional status among HIV-infected children is clearly of paramount importance⁽¹⁾. WHO guidelines for incorporating nutritional support into care programs for HIV-infected children are available for local adaptation⁽²⁾, and nutritional support services appear to be generally available at HIV care and treatment sites in sub-Saharan Africa⁽³⁾. However, the evidence-base for the effectiveness of specific nutritional interventions in HIV-infected children is weak⁽⁴⁾, and nutritional interventions impose additional costs on HIV care programs⁽⁵⁾.

These costs could be reduced by developing locally-produced food-based supplements, which are also likely to be more palatable and culturally acceptable. Leaf concentrate is a food-based nutritional supplement⁽⁶⁾ which has been shown to be effective at preventing anemia in adolescent girls⁽⁷⁾ and pregnant women⁽⁸⁾ in India. Although anemia may actually be less prevalent among HIV-infected than among uninfected children⁽⁹⁾, it remains an important comorbidity and, together with other nutritional deficiencies and diarrheal disease, is a strong predictor of poor outcomes among children after initiation of antiretroviral therapy (ART)^(10; 11; 12).

Leaf concentrate was discovered in France in the 18th century, and was developed as a foodstuff in England between 1940-1970⁽¹³⁾. It has since been promoted by several non-governmental organizations, including Leaf for Life in the USA⁽¹⁴⁾ and the Association pour la Promotion des Extraits Foliaires en Nutrition (APEF) in France⁽¹⁵⁾, as a sustainable form of protein and micronutrient supplementation for low-income communities⁽⁶⁾. In this study, we investigate the potential benefits of dried lucerne (alfalfa) leaf concentrate as a nutritional supplement for HIV-infected ART-naïve children living in two cities in Burundi. The aim of the study was to determine the acceptability and palatability of leaf concentrate, and to measure its effectiveness (compared with skimmed milk powder) in relation to HIV-specific (viral load, CD4+ count) and general (haematological, anthropometric) parameters, and overall health.

Methods

Study population and recruitment

Our target population was antiretroviral-naïve HIV-positive children age 5-14 years. Children were recruited from two clinical centres (in Bujumbura and Kirundo) run by the Association Nationale de Soutien aux Séropositifs et malades du Sida (ANSS, National Association of Support for People Living with HIV and AIDS). Founded in 1993, ANSS' mission is to promote the prevention of HIV transmission and to improve the well-being of people living with and affected by HIV. The ANSS was the first Burundian civil society organization to provide HIV services to people living with HIV, including the distribution of antiretroviral therapy. At the end 2013, the ANSS was providing treatment to more than 6,000 people in Burundi. From 2007 to 2013, the association provided HIV counselling and testing to more than 56,000 people as part of its HIV prevention efforts. Recruitment was conducted by medical staff at the two centres, who identified eligible children at routine clinic appointments to which HIV-positive children are invited for medical and psychosocial monitoring purposes. The same staff explained the purpose of the study to children and their parent(s)/guardian(s), who were provided with a study information sheet and consent form in the local language (Kirundi).

Ethical approval, trial registration and consent

Ethical approval for the study was granted on 16/06/2012 by the "Comité National d'Ethique pour la protection des êtres humains participants à la recherche biomédicale et comportementale" of Burundi. Written informed consent was obtained from the parent(s)/guardian(s) of eligible children, with original consent forms being held by ANSS.

Inclusion criteria

Children were eligible if they were ≥ 5 and ≤ 14 years old, had tested positive for HIV-1 (HIV 1/2 STAT-PAK® DIPSTICK Assay, Chembio Diagnostic Systems, Inc., 3661 Horseblock Road, Medford, NY 11763, USA), and had a CD4+ cell count ≥ 350 cells/mm³(16). Children with a CD4+ cell count < 500 cells/mm³ were required to receive prophylactic sulfamethoxazole/trimethoprim⁽²⁾. All children were dewormed before the study began using an appropriate dose of albendazole.

Exclusion criteria

Children were ineligible if they had a major concurrent illness (e.g. cancer, tuberculosis or diabetes), a BMI <12 kg/m², were liable to move out of the study area or were unlikely to be able to receive the weekly supplements, or had participated in a clinical trial in the preceding 3 months.

Primary and secondary outcomes

Primary outcomes were percentage of CD4+ T lymphocytes, HIV viral load, and Hb measured after 6 months of follow-up. Secondary outcomes (also measured at 6 months) included: anthropometric indices (height-for-age, weight-for-age, handgrip strength); episodes of illness (diarrhea, respiratory); overall health (anemia, inflammation (CRP); fatigue, appetite); and palatability of the supplements (appearance, texture, taste).

Sample size

We aimed to recruit 42 children into each arm of the trial. This sample size was calculated to yield 90% power at a 5% level of significance to detect: a) equivalence between LCP and SMP in percentage of CD4+ T lymphocytes at 6 months, by excluding a difference in means of 5% (assuming that the standard deviation (SD) in the outcome = 7%); b) equivalence between LCP and SMP in log₁₀ HIV viral load, by excluding a difference in means of 0.5 (assuming SD=0.7 and 15% of participants lost to follow-up); c) superiority of LCP over SMP in increasing mean Hb from 115g/L (SD 14g/L) to 125g/L (SD 14g/L), compared with no change in SMP arm (because LCP provides 5.4mg Fe compared with 0.1mg Fe in SMP).

Randomization, allocation and adherence

Eligible children with consent to participate were allocated to LCP or SMP using a computer-generated random number sequence and consecutively-numbered sealed envelopes.

Envelopes were held and opened by a member of ANSS administrative staff not otherwise involved in the study, i.e. allocation was concealed from the ANSS medical staff who were running the trial. Delivery of supplements to all participants were made by ANSS health workers at an interval of 2 weeks and were recorded for each child. Observance of supplement consumption was monitored and discussed collectively (with all children and their parents) at each monthly monitoring visit in order to highlight difficulties and elaborate solutions. For the few parents who were not present at a supplement delivery time or reported

difficulties in supplement consumption by their child, ANSS health workers visited them at home to see how supplements were consumed.

Leaf concentrate powder

The LCP group received 10g of lucerne (*Medicago sativa*) leaf concentrate powder (France-Luzerne Agricultural Co-operative, 51130 Aulnay-aux-Planches, France). The leaf concentrate was contained in sachets, and could be taken orally by mixing with water or tea. The composition of the LCP is shown in **Table 1**.

Skimmed milk powder

The SMP group received 15g of skimmed milk powder (Société Industrielle Laitière du Léon, Le Raden, 29860 Plouvien, France). The skimmed milk powder was given in a sachet which could be taken orally by mixing with water or tea. The composition of the SMP is shown in **Table 1**.

HIV-specific parameters and non-specific blood parameters

HIV-1 viral load, absolute count and percentage of CD4+ T lymphocytes (CD4%), and general blood parameters (Hb, WBC, lymphocytes, platelets, and CRP) were measured at baseline, 3 and 6 months in blood samples which were collected at ANSS clinics and transferred to the National Institute of Public Health (INSP), Rivubu Avenue, Bujumbura and to a laboratory at the New Hope Centre, Bujumbura. HIV viral load was measured at INSP using real-time PCR (Abbott RealTime HIV-1 assay, Abbott Laboratories, Abbott Park, Illinois, USA). CD4 absolute count and percentage was measured at the New Hope Centre using BD FACS Count (BD Biosciences, 2350 Qume Drive, San Jose, CA, USA). Non-specific blood parameters were measured at INSP. Anemia was defined as Hb <115g/L, CRP-positivity as CRP \geq 6mg/L.

Anthropometric parameters

Weight and height were measured using electronic scales and standing measurement boards during clinic visits at baseline, 3 and 6 months. Height-for-age and weight-for-age z-scores were estimated against British 1990 (UK90) growth charts^(17; 18). The UK90 BMI reference provides centile curves for children from birth to 23 years, based on a sample of 32,222 measurements from 12 distinct surveys in British children collected between 1978 and 1994.

Handgrip strength was measured, using a Jamar® hydraulic hand dynamometer (Patterson Medical, Nottinghamshire, UK), by two successive trials on the dominant side with the elbow at 90°, the maximal score of the two trials being recorded.

Diarrhea, respiratory infection, appetite, fatigue and overall health levels

Diarrhea, respiratory infection and other illnesses were recorded by ANSS clinicians monthly and during clinic visits at baseline, 3 and 6 months. Disease occurrences and corresponding treatments were recorded for each child in order to detect the medical events during each three months period (0-3months, 3-6months). Appetite, fatigue and overall health were assessed by questioning parents/carers at baseline, 3 and 6 months, with the following possible responses: fatigue (none, moderate, severe); appetite (weak, satisfactory, good, very good); overall health (very poor, poor, satisfactory, good, excellent).

Palatability

Three aspects of palatability (appearance, texture and taste) were recorded by ANSS health workers during clinic visits at 3 and 6 months, with the following possible responses: appearance (unappealing, acceptable, appealing, very appealing); texture (unappealing, acceptable, appealing, very appealing); taste (very bad, bad, acceptable, good, very good).

Statistical analyses

Data were entered into a computer using Microsoft Excel, and were analyzed using Stata (StataCorp. 2013. Stata Statistical Software: Release 13. College Station, TX: StataCorp LP). Between-group differences in continuous outcome measures were estimated using linear regression models. Between-group differences in binary outcome measures were tested using Fisher's exact tests. We fitted linear regression models using generalized estimating equations for each continuous outcome measure, including an interaction term between study arm and time-point (0, 3 or 6 months) to estimate within- and between-group changes at each time-point. We used an exchangeable within-child correlation structure (which assumes that the observations within a subject are equally correlated), and we verified by inspection that other correlation structures (ar1 and ar2) did not produce different results.

Results

Characteristics of trial participants

The median age of the participants was 10 years (range 5 to 14 years) and 39.8% (33/83) were male (**Table 2**). 42 children were randomized to receive SMP, and 41 to LCP (**Figure 1**).

Loss to follow-up and adherence

One child in the SMP arm and two children in the LCP arm did not receive their allocated intervention and dropped out at the beginning of the trial (**Figure 1**). One child in the LCP arm dropped out just after the 3-month follow-up. Three children in each arm had an HIV viral load $<350\text{cells/mm}^3$ measured at the 3-month follow-up and proceeded to receive antiretroviral therapy. At 6 months, primary and secondary outcome measures were available for 90.5% (38/42) and 85.4% (35/41) of participants in the SMP and LCP groups, respectively. The final analysis was performed on the 38 SMP and 35 LCP children who had data on one or more outcome measures at the 6 months follow-up.

Blood parameters

The 95% confidence intervals of the between-group differences in percentage of CD4+ T lymphocytes and HIV-1 viral load at 6 months suggested non-equivalence between SMP and LCP (**Table 3**). For CD4+ percentage, equivalence had been defined as a difference not greater than 5%, but the 95% CI for the between-group difference in this parameter at 6 months was -1.17% to 6.07%, i.e. the upper bound exceeded 5% (in favour of LCP). Similarly, equivalence for HIV viral load had been defined as a difference not greater than 0.5 logs, but the 95% CI of the observed difference was -0.19 to 0.67, i.e. the upper bound exceeded 0.5 (in favour of SMP). However, the 95% CI for the baseline values of these two HIV-related measures were also consistent with non-equivalence (suggesting the role of chance). Adjusted for baseline, the between-group differences at 6 months for CD4+ percentage and HIV viral load were 0.59% (95% CI -1.42% to 2.61%) and 0.18 (-0.08 to 0.44), respectively. These differences and their 95% confidence intervals are consistent with equivalence between SMP and LCP.

The estimated additional effects of LCP (compared with SMP) on the changes in CD4+ percentage and HIV viral load over 6 months indicate no additional effect of LCP: for CD4+ percentage, 0.97% (95% CI -0.86%, 2.80%), $P=0.30$; for \log_{10} HIV viral load, 0.20 (-0.03,

0.43), $P=0.09$. LCP was not superior to SMP in improving Hb, with similar increases over the 6 months in both arms, and no additional effect of LCP compared to SMP (-1.71 (95% CI -6.49 to 3.07) g/L, $P=0.48$). LCP did not have any additional effect over SMP on 0-6 month changes in any of the other blood parameters (**Supplementary Table 1**).

We found only slight changes in the interaction (adjusted 0-6 month change) estimates and their confidence intervals when we used ar1 and ar2 autoregression correlation structures, instead of an exchangeable (exch) structure: percentage CD4+ T lymphocytes, exch 0.97% (-0.86%, 2.80%), ar1 0.98% (-0.93%, 2.88%), ar2 0.90% (-1.03%, 2.83%); \log_{10} HIV-1 viral load, exch 0.20 (-0.03, 0.43), ar1 0.19 (-0.04, 0.43), ar2 0.23 (-0.01, 0.46); Hb exch -1.71 (-6.49, 3.07) g/L, ar1 -2.14 (-7.20, 2.91) g/L, ar2 -1.51 (-6.47, 3.45) g/L.

Anthropometric parameters

Height-for-age and weight-for-age Z-scores increased to the same degree in both groups of children over the 6 months, and there was no additional effect of LCP over SMP (**Table 4**). There were no between-group differences in median handgrip strength scores at 3 or 6 months, but children in the LCP group increased their (geometric mean) handgrip strength score 2.4-fold over the 6 months, compared with no change in the SMP group (**Table 4**).

Diarrhea, respiratory infection and other illnesses

There were few reported cases of diarrhea during the study in either group: in the LCP group, 4 cases at baseline, then 1 case during 0-3 months and 1 case during 3-6 months follow-up; in the SMP group, 3 cases at baseline, no cases during 0-3 or 3-6 months follow-up. Pneumonia was reported to be present only once in each group, in both cases, during 3-6 months follow-up. ENT infections (angina, ear infection) occurred in 2 LCP group children and in 5 SMP group children. Malaria was diagnosed during an acute clinic access by 2 children of the LCP group and 3 children of the SMP group.

Anemia, inflammation, fatigue, appetite and overall health

Overall prevalence of anemia ($Hb < 115$ g/L) fell from 39.7% (29/73, 95% CI 28.5-51.9%) at baseline to 24.7% (18/73, 15.3-36.1%) at 3 months and 20.6% (15/73, 12.0-31.6%) at 6 months, with similar prevalence in both groups at each time-point. CRP-positivity fell from 23.3% (17/73, 14.2-34.6%) at baseline to 15.1% (11/73, 7.8-25.4%) and 9.6% (7/73, 3.9-18.8%) at 3 and 6 months, respectively, again with no detectable between-group differences

(**Supplementary Table 2**). Appetite and overall health appeared to be better at 6 months in the LCP group, although there were no differences if the responses “very good” and “good” to the appetite question were merged (92.1% (35/38, 78.6-98.3%) in the SMP group compared with 97.1% (34/35, 85.1-99.9%) in the LCP group, $P=1.00$) and if the “excellent” and “good” overall health responses were merged (73.7% (28/38, 56.9-86.6%) in the SMP group compared with 85.7% (30/35, 69.7-95.2%) in the LCP group, $P=0.31$).

Palatability

The appearance, texture and taste of SMP were rated more favourably than for LCP. All (38/38) of participants in the SMP arm rated appearance and texture as appealing/very appealing at 3 and 6 months, and 97.4% (37/38, 95% CI 86.2-99.9%) and 97.3% (36/37, 85.5-99.9%) rated taste as good/very good at 3 and 6 months, respectively (**Supplementary Table 3**). In contrast, 80.0% (28/35, 63.1-91.6%) and 82.9% (29/35, 66.4-93.4%) of participants in the LCP arm rated appearance and texture as appealing/very appealing, and 77.1% (27/35, 59.9-89.6%) rated taste as good/very good at 3 months. At 6 months, 85.7% (30/35, 69.7-95.2%) and 80.0% (28/35, 63.1-91.6%) rated the appearance and texture of LCP as appealing/very appealing, and 68.6% (24/35, 50.7-83.1%) the taste as good/very good. There were fewer negative ratings of the palatability of LCP at 6 compared with 3 month, but with insufficient numbers to provide statistical evidence of change: unappealing appearance 11.4% (4/35, 3.2-26.7%) to 0.0% (0/35); unappealing texture 5.7% (2/35, 0.7-19.2%) to 2.9% (1/35, 0.1-14.9%); and bad taste 17.1% (6/35, 6.6-33.6%) to 11.4% (4/35, 3.2-26.7%).

Discussion

During 6 months of follow-up in this population of HIV-positive ART-naïve children, LCP was equivalent to SMP, after adjustment for baseline values, in relation to HIV-specific blood parameters (which showed little or no change in either group), and did not demonstrate superiority over SMP in relation to Hb levels (which increased in both groups). Higher levels of appetite and overall health were reported for children in the LCP group, and they also had higher handgrip strength scores. There were no differences in any of the other outcome measures. In terms of palatability, LCP was less acceptable than SMP to the children in this population, but with few negative ratings of appearance, texture and taste.

Our study was powered to detect equivalence in relation to HIV-specific blood parameters (CD4% and HIV viral load), and superiority in relation to Hb levels (because of the higher

iron content of LCP). Differences between the two groups at baseline (which could only have arisen by chance) exceeded the thresholds for equivalence, suggesting that a wider *a priori* equivalence interval was required. However, the fully-adjusted analysis using longitudinal linear regression models gave estimates which were consistent with equivalence in the primary outcome measures. No differences were detected in any of the other outcome measures, although assessment of 'appetite', 'overall health' and handgrip strength at 6 months appeared to favour LCP. Several participants were lost to follow-up from each arm of the study, which would have reduced the power of the study, and the study was not sized to detect pre-specified differences in the secondary outcome measures. Allocation to the supplementation arms was concealed, but children and their parents/carers and field workers were not blinded to allocation.

In the fifteen or so years since paediatric HIV came to be regarded as a chronic disease⁽¹⁹⁾, very little research has been conducted into the nutritional aspects of HIV infection in children age ≥ 5 years. Even for children under 5 years old, the evidence-based for nutritional interventions remains weak. The most recent Cochrane review reported some evidence for beneficial effects of vitamin A, zinc and micronutrient supplements, based on seven trials in sub-Saharan African countries⁽⁴⁾. This dearth of evidence may have arisen because children and adolescents fall into a gap between research into HIV infection during pregnancy and infancy (because of the focus on vertical transmission) and during adulthood (wherein lies the main 'burden' of HIV infection, in terms of numbers of people infected), or it may have arisen because nutritional interventions have proven to be less effective than expected, given observational evidence for the importance of nutrition among people infected with HIV⁽²⁰⁾.

In policy terms, it has been acknowledged that children, together with adolescents (girls in particular), are being "left behind" in global efforts to end the AIDS epidemic⁽²¹⁾. This may be attributable in part to success in reducing the incidence of new infections in children, which have fallen by 58% from a peak of 580,000 in 2002. However, in 2013 there were an estimated 240,000 new infections, and 3.2 million children under 15 were living with HIV, 91% of whom were in sub-Saharan Africa. Given that these children are more likely to live in food-insecure households⁽¹⁾, that HIV infection is associated with faltering growth to at least 10 years of age⁽²²⁾, and that micronutrient sufficiency might be an important adjunct to anti-retroviral therapy⁽²³⁾, effective and sustainable approaches to maintaining adequate nutrition for HIV-infected children in resource-poor settings should be a priority area for research.

In the absence of comparable studies of nutritional interventions among ART-naïve children age 5-14 years, and because our study did not have an unsupplemented control group, we can only conclude that children receiving leaf concentrate and dairy supplements had equivalent outcomes in terms of CD4 percentage and HIV viral load. We cannot conclude that either supplement had any effect on these two outcomes, i.e. equivalence may indicate lack of effect. The higher iron content of the leaf concentrate supplement did not have any additional effect on Hb levels, for which there was evidence of improvement in both groups. This result may be attributable to the relatively normal baseline hemoglobin levels and/or to the consumption of LCP within tea (an inhibitor of iron absorption). Similar improvements in inflammation and fatigue were detected in both arms. Although both supplements contained zinc, the amounts are unlikely to have had an effect on diarrhea, for which some evidence of a protective effect of zinc supplementation has been reported⁽⁴⁾. The leaf concentrate supplement contained much higher levels of β -carotene than did the skimmed milk powder, but our study was not designed to detect differential effects of leaf concentrate on outcomes such as diarrhea and cough, for which evidence of beneficial effects of vitamin A supplementation is reasonably consistent⁽⁴⁾.

That growth indices improved in both groups is reassuring, given the accuracy with which these outcomes can be measured and the 6-month duration of our study. Classical anthropometric indices improved to the same degree in both groups, whereas handgrip strength score increased at a higher level in the LCP group. These data, combined with impact on appetite and overall health, suggest that LCP may increase the nutritional status more efficiently than SMP, because a relationship has been highlighted between muscle mass and function in children^(24; 25). These differences need to be confirmed by a larger study, ideally comparing LCP to no supplement (or to a supplement with known effects), and perhaps providing larger amounts of supplement or in conjunction with other nutrients.

HIV-infected adults have increased energy requirements⁽²⁶⁾, and additional calories are recommended by WHO for HIV-positive children⁽²⁷⁾, particularly in children experiencing weight loss. In our study, even greater gains in height and weight might have been achieved had the supplements been used to fortify a calorific snack, something which should be considered for future trials in this population. Incorporating the supplements into isocaloric biscuits could also address the issue of blinding (if the non-LCP biscuits were dyed green),

and might improve adherence and acceptability⁽²⁸⁾. The calcium content of both supplements might be expected to ameliorate low bone mass, an effect of HIV infection that has been reported for children and adolescents⁽²⁹⁾, although longer follow-up would probably be required to detect any such effects. Trials which aim to test novel dietary interventions would benefit from information obtained from qualitative research into local tastes and preferences⁽³⁰⁾. Such research would, ideally, be conducted before the trial, but could also be conducted during the trial.

Leaf concentrate might be considered as a candidate ingredient for the food multimix (FMM) concept⁽³¹⁾. The aim of FMM is to combine locally-available and commonly-consumed food ingredients into a culturally-acceptable end product which makes the best use of each ingredient's nutritional 'strengths'. Although our study compared LCP *against* SMP, parallel production of the two supplements is rendered eminently feasible by the unique capacity of the leaf fractionation concept to provide direct access to the edible fraction of the leaves and to provide feed for livestock. For example, lucerne grown on 1 hectare can generate the production of 2,000 kg of SMP if consumed directly by cattle; the same lucerne, fractionated into LCP for human consumption, fibre cake and whey for cattle can still produce 1,800 kg of SMP, with the added production of 1,200 kg of LCP⁽⁶⁾. In this context, our demonstration that the nutritional qualities of SMP and LCP are comparable is an important finding.

The acceptability of leaf concentrate, and any economic advantage that it may have over industrially-sourced skimmed milk powder, remains to be determined. Much as local and sustainable production of leaf concentrate is espoused as a key selling point⁽⁶⁾, this remains somewhat theoretical, and the development of the leaf fractionation concept is an ongoing process. In the case of Burundi, as in many tropical and sub-tropical countries, vegetables constitute a far greater component of the traditional diet than dairy products⁽³²⁾, and vegetable crops (including those with leaves which are suitable for leaf concentrate production, e.g. beans) are one of the main agricultural products⁽³³⁾.

These uncertainties should not deter further pursuit of alternative food-based solutions, including leaf fractionation, to nutritional problems among impoverished and vulnerable communities, particularly where local resilience might be needed in the face of endemic food insecurity and episodic civil instability^(34; 35). In the meantime, our small study in a

particularly vulnerable group has demonstrated that a leaf-derived food supplement could meet this need as effectively as a conventional milk-based supplement.

Table 1: Composition (mean \pm SD) of leaf concentrate and skimmed milk powders

Nutrient	Leaf concentrate powder (per 10g sachet) ⁽³⁶⁾	Skimmed milk powder* (per 15g sachet)
Protein	5.1 \pm 0.46 g	5.2 g
Fat	1.0 \pm 0.1 g	0.1 g
Linoleic acid (omega-6) [†]	12.2 \pm 0.8 %	
α -linolenic acid (omega-3) [†]	34.1 \pm 4.5 %	
Carbohydrates	0.1 \pm 0.02 g	7.8 g
Iron	5.4 \pm 0.86 mg	0.072 \pm 0.02 mg
Folic acid	13.4 \pm 7.3 μ g	10.3 \pm 4.6 μ g
β -carotene [‡]	767 \pm 382 μ g	1.4 \pm 0.6 μ g
Vitamin E	9.9 \pm 1.9 mg	3.9 \pm 2.0 μ g
Vitamin C	6mg [§]	1.24 \pm 0.4 mg
Calcium	338 \pm 43 mg	185 \pm 29 mg
Copper	76 \pm 10 μ g	9.0 \pm 3 mg
Manganese	0.6 \pm 0.06 mg	6.6 \pm 3 μ g
Zinc	0.2 \pm 0.08 mg	0.63 \pm 0.05 mg

* French Agency for Food, Environmental and Occupational Health and Safety (<https://pro.anses.fr/TableCIQUAL/>, accessed 03/07/2015)

[†]percentage of fat content

[‡]retinol activity equivalents (RAE)

[§]corresponding to ascorbic acid added during manufacturing to prevent oxidation

Table 2: Characteristics of children in each arm at the beginning of the study (baseline)

		Skimmed milk (n=42) [†]	Leaf concentrate (n=41) [†]
Site	Bujumbura	25 (59.5%)	24 (58.4%)
	Kirundo	17 (40.5%)	17 (41.5%)
Sex (male)		15 (35.7%)	18 (43.9%)
Age (years)		10 (8 to 11)	10 (8 to 12)
Height-for-age Z-score		-1.10 (-2.05 to -0.39)	-1.47 (-1.95 to -0.74)
Weight-for-age Z-score		-1.33 (-2.36 to -0.54)	-2.20 (-1.58 to -0.81)
Handgrip strength score (kg)		12 (8 to 15)	13 (10 to 16)
Log ₁₀ HIV-1 viral load		4.39 (3.74 to 5.13)	4.39 (4.03 to 4.74)
Number of CD4+ T lymphocytes (cells/ μ L)		723 (606 to 966)	678 (523 to 895)
Percentage of CD4+ T lymphocytes (%)		27.0 (21.5 to 30.8)	27.9 (24.5 to 32.6)
Haemoglobin (g/L)		117 (110 to 127)	119 (111 to 125)
Presence of diarrhea		4 (9.5%)	3 (7.3%)
Presence of respiratory infection		0 (0.0%)	0 (0.0%)
CRP positive (\geq 6mg/L)		11 (26.2%)	10 (24.4%)

[†] Values are frequency (%) or median (IQR).

Table 3: Percentage of CD4+ T lymphocytes, HIV-1 viral load and Hb at baseline, 3 and 6 months*

	Skimmed milk powder group (n=38)	Leaf concentrate powder group (n=35)	Between-group difference [†]
Percentage of CD4+ T lymphocytes (%)			
Baseline (n=37, 35)	26.6 (24.3, 29.0)	28.6 (26.3, 31.0)	1.96 (-1.29, 5.22)
3 months (n=38, 33)	25.1 (22.8, 27.4)	27.9 (25.1, 30.7)	2.76 (-0.75, 6.27)
6 months (n=38, 34)	26.4 (23.9, 28.8)	28.8 (26.1, 31.6)	2.45 (-1.17, 6.07)
0-6 month change	-0.24 (-1.36, 0.88)	0.29 (-1.30, 1.89)	0.53 (-1.41, 2.48)
Adjusted 0-6 month change [‡]	-0.01 (-1.32, 1.30)	0.96 (-0.72, 2.65)	0.97 (-0.86, 2.80)
Log ₁₀ HIV-1 viral load			
Baseline (n=38, 35)	4.23 (3.89, 4.57)	4.33 (4.11, 4.56)	0.10 (-0.30, 0.51)
3 months (n=38, 35)	4.23 (3.91, 4.54)	4.45 (4.20, 4.70)	0.22 (-0.18, 0.62)
6 months (n=36, 35)	4.33 (4.02, 4.65)	4.57 (4.28, 4.87)	0.24 (-0.19, 0.67)
0-6 month change	0.06 (-0.11, 0.23)	0.23 (0.05, 0.41)	0.17 (-0.08, 0.42)
Adjusted 0-6 month change [‡]	0.14 (-0.07, 0.36)	0.34 (0.12, 0.57)	0.20 (-0.03, 0.43)
Haemoglobin (g/L) [‡]			
Baseline (n=38, 35)	116 (112, 121)	116 (111, 120)	-0.82 (-6.90, 5.26)
3 months (n=38, 35)	121 (115, 127)	125 (119, 130)	3.75 (-3.66, 11.2)
6 months (n=38, 35)	130 (125, 135)	126 (121, 131)	-3.53 (-10.4, 3.37)
0-6 month change	13.6 (10.6, 16.5)	10.5 (6.32, 14.6)	-3.13 (-8.21, 1.96)
Adjusted 0-6 month change [‡]	11.4 (8.44, 14.3)	9.66 (5.13, 14.2)	-1.71 (-6.49, 3.07)

* Values shown are mean (95% CI).

[†] The between-group difference corresponding to the “0-6 month change” shows the estimated additional effect of leaf concentrate powder compared with milk powder, i.e. the interaction term between the intervention and 6-month time-point.

[‡] Adjusted for age, sex, all blood parameters, CRP positivity, and height-for-age Z-score.

Table 4: Anthropometric parameters at baseline, 3 and 6 months

	Skimmed milk powder group	Leaf concentrate powder group	Between-group difference*
Height-for-age Z-score			
Baseline (n=38, 35)	-1.17 (-1.56, -0.78)	-1.47 (-1.82, -1.11)	-0.30 (-0.82, 0.22)
3 months (n=38, 35)	-0.82 (-1.21, -0.44)	-1.14 (-1.52, -0.76)	-0.32 (-0.85, 0.22)
6 months (n=38, 35)	-0.63 (-1.03, -0.23)	-0.96 (-1.34, -0.58)	-0.33 (-0.88, 0.21)
0-6 month change	0.54 (0.45, 0.62)	0.51 (0.42, 0.60)	-0.03 (-0.15, 0.10)
Adjusted 0-6 month change [†]	0.53 (0.44, 0.62)	0.48 (0.37, 0.59)	-0.05 (-0.18, 0.08)
Weight-for-age Z-score			
Baseline (n=38, 35)	-1.36 (-1.80, -0.92)	-1.61 (-1.94, -1.29)	-0.25 (-0.80, 0.29)
3 months (n=38, 35)	-1.20 (-1.64, -0.75)	-1.49 (-1.81, -1.16)	-0.29 (-0.84, 0.26)
6 months (n=38, 35)	-1.03 (-1.50, -0.56)	-1.35 (-1.76, -0.95)	-0.32 (-0.94, 0.29)
0-6 month change	0.31 (0.19, 0.42)	0.25 (0.02, 0.49)	-0.05 (-0.31, 0.21)
Adjusted 0-6 month change [†]	0.27 (0.14, 0.39)	0.24 (-0.01, 0.50)	-0.02 (-0.26, 0.22)
Handgrip strength (kg)[‡]			
Baseline (n=38, 35)	10.8 (9.2, 12.7)	11.4 (9.7, 13.5)	1.02 (0.93, 1.13)
3 months (n=38, 35)	11.1 (9.5, 13.0)	11.7 (9.8, 13.9)	1.02 (0.93, 1.13)
6 months (n=38, 35)	11.6 (10.2, 13.3)	12.5 (10.7, 14.5)	1.03 (0.95, 1.12)
0-6 month change	1.73 (0.60, 4.98)	2.59 (1.37, 4.86)	1.49 (0.44, 5.10)
Adjusted 0-6 month change [†]	1.76 (0.59, 5.27)	2.40 (1.18, 4.91)	1.36 (0.38, 4.94)

* The between-group difference corresponding to the “0-6 month change” shows the estimated additional effect of leaf concentrate powder compared with milk powder, i.e. the interaction term between the intervention and 6-month time-point.

[†] Adjusted for age, sex and all blood parameters.

[‡] Values shown for baseline, 3- and 6-month levels of handgrip strength are geometric mean (95% CI). Between-group difference and 0-6 month changes are ratios of geometric means from linear regression of log-transformed handgrip strength values, e.g. coefficient = 1.10 indicates that strength is 10% higher in LCP group compared with SMP group, coefficient = 0.90 indicates that strength is 10% lower in LCP group compared with SMP group, coefficient = 1.00 indicates no difference/change.

References

1. Rose AM, Hall CS, Martinez-Alier N (2014) Aetiology and management of malnutrition in HIV-positive children. *Arch Dis Child* **99**, 546-551.
2. World Health Organisation (2009) *Guidelines for an integrated approach to the nutritional care of HIV-infected children (6 months - 14 years)*. Geneva.
3. Anema A, Zhang W, Wu Y *et al.* (2012) Availability of nutritional support services in HIV care and treatment sites in sub-Saharan African countries. *Public Health Nutr* **15**, 938-947.
4. Irlam JH, Siegfried N, Visser ME *et al.* (2013) Micronutrient supplementation for children with HIV infection. *Cochrane Database Syst Rev* **10**, Cd010666.
5. Cobb G, Bland RM (2013) Nutritional supplementation: the additional costs of managing children infected with HIV in resource-constrained settings. *Trop Med Int Health* **18**, 45-52.
6. Davys MNG, Richardier FC, Kennedy D *et al.* (2011) Leaf Concentrate and Other Benefits of Leaf Fractionation. In *Combating Micronutrient Deficiencies: Food-based Approaches*, pp. 338-365 [B Thompson and L Amoroso, editors]. Rome: FAO.
7. Vyas S, Collin SM, Bertin E *et al.* (2010) Leaf concentrate as an alternative to iron and folic acid supplements for anaemic adolescent girls: a randomised controlled trial in India. *Public Health Nutr* **13**, 418-423.
8. Magon A, Collin SM, Joshi P *et al.* (2014) Leaf concentrate fortification of antenatal protein-calorie snacks improves pregnancy outcomes. *J Health Popul Nutr* **32**, 430-440.
9. Esan MO, Jonker FA, Hensbroek MB *et al.* (2012) Iron deficiency in children with HIV-associated anaemia: a systematic review and meta-analysis. *T Roy Soc Trop Med H* **106**, 579-587.
10. Auld AF, Tuho MZ, Ekra KA *et al.* (2014) Temporal trends in mortality and loss to follow-up among children enrolled in Cote d'Ivoire's national antiretroviral therapy program. *Pediatr Infect Dis J* **33**, 1134-1140.
11. Zanoni BC, Phungula T, Zanoni HM *et al.* (2012) Predictors of poor CD4 and weight recovery in HIV-infected children initiating ART in South Africa. *PloS one* **7**, e33611.
12. Zanoni BC, Phungula T, Zanoni HM *et al.* (2011) Risk factors associated with increased mortality among HIV infected children initiating antiretroviral therapy (ART) in South Africa. *PloS one* **6**, e22706.
13. Pirie NW (1987) *Leaf Protein and Its By-Products in Human and Animal Nutrition*. 2nd ed. Cambridge: Cambridge University Press.
14. Kennedy D (1993) Leaf Concentrate: A Field Guide for Small Scale Programs. *Leaf for Life, Kentucky, USA*.
15. (2008). *Association pour la Promotion des Extraits Foliaires en Nutrition (APEF)*. <http://www.nutrition-luzerne.org/> (accessed 13/07/2015)
16. Conseil National de la Lutte Contre le VIH/SIDA (2010) *Schemas de traitement par les arv au Burundi*. Bujumbura, Burundi.

17. Cole TJ, Freeman JV, Preece MA (1998) British 1990 growth reference centiles for weight, height, body mass index and head circumference fitted by maximum penalized likelihood. *Stat Med* **17**, 407-429.
18. Cole TJ, Williams AF, Wright CM (2011) Revised birth centiles for weight, length and head circumference in the UK-WHO growth charts. *Ann Hum Biol* **38**, 7-11.
19. Miller TL (2000) Nutrition in paediatric human immunodeficiency virus infection. *Proc Nutr Soc* **59**, 155-162.
20. Salomon J, De TP, Melchior JC (2002) Nutrition and HIV infection. *Br J Nutr* **87 Suppl 1**, S111-119.
21. UNAIDS (2014) *The Gap Report*. Geneva: Joint United Nations Programme on HIV/AIDS (UNAIDS).
22. Newell ML, Borja MC, Peckham C (2003) Height, weight, and growth in children born to mothers with HIV-1 infection in Europe. *Pediatrics* **111**, e52-60.
23. Drain PK, Kupka R, Mugusi F *et al.* (2007) Micronutrients in HIV-positive persons receiving highly active antiretroviral therapy. *Am J Clin Nutr* **85**, 333-345.
24. Marrodan Serrano MD, Romero Collazos JF, Moreno Romero S *et al.* (2009) [Handgrip strength in children and teenagers aged from 6 to 18 years: reference values and relationship with size and body composition]. *An Pediatr (Barc)* **70**, 340-348.
25. Schweizer R, Martin DD, Schonau E *et al.* (2008) Muscle function improves during growth hormone therapy in short children born small for gestational age: results of a peripheral quantitative computed tomography study on body composition. *J Clin Endocrinol Metab* **93**, 2978-2983.
26. Kosmiski L (2011) Energy expenditure in HIV infection. *Am J Clin Nutr* **94**, 1677s-1682s.
27. World Health Organisation (2003) *Nutrient requirements for people living with HIV/AIDS*. Geneva.
28. Ernst J, Etyyang G, Neumann CG (2014) High-nutrition biscuits to increase animal protein in diets of HIV-infected Kenyan women and their children: a study in progress. *Food Nutr Bull* **35**, S198-204.
29. Lima LR, Silva RC, Giuliano Ide C *et al.* (2013) Bone mass in children and adolescents infected with human immunodeficiency virus. *J Pediatr (Rio J)* **89**, 91-99.
30. Hong SY, Hendricks KM, Wanke C *et al.* (2013) Development of a nutrient-dense food supplement for HIV-infected women in rural Kenya using qualitative and quantitative research methods. *Public Health Nutr* **16**, 721-729.
31. Zotor FB, Amuna P (2008) The food multimix concept: new innovative approach to meeting nutritional challenges in Sub-Saharan Africa. *Proc Nutr Soc* **67**, 98-104.
32. Burundi Income and Daily Diet: Open Data for Africa (<http://opendataforafrica.org/> accessed 13/07/2015).
33. Ministère des Finances et de la Planification du Développement Economique (2014) *Enquête nationale agricole du Burundi 2012-2013*. Bujumbura, Burundi.

34. Rossi L, Verna D, Villeneuve SL (2008) The humanitarian emergency in Burundi: evaluation of the operational strategy for management of nutritional crisis. *Public Health Nutr* **11**, 699-705.
35. Verwimp P (2012) Undernutrition, subsequent risk of mortality and civil war in Burundi. *Econ Hum Biol* **10**, 221-231.
36. European Food Safety Authority (EFSA) Panel on Dietetic Products, Nutrition and Allergies (NDA) (2009) Opinion on the safety of 'Alfalfa protein concentrate' as food. *EFSA Journal* **7**, Article 997.