



Castaneda-Gameros, D., Redwood, S., & Thompson, J. (2017). Low Nutrient Intake and Frailty Among Overweight and Obese Migrant Women From Ethnically Diverse Backgrounds Ages 60 Years and Older: A Mixed-Methods Study. *Journal of Nutrition Education and Behavior*, 49(1), 3-10.e1. <https://doi.org/10.1016/j.jneb.2016.08.006>

Peer reviewed version

Link to published version (if available):
[10.1016/j.jneb.2016.08.006](https://doi.org/10.1016/j.jneb.2016.08.006)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the accepted author manuscript (AAM). The final published version (version of record) is available online via Elsevier at <http://dx.doi.org/10.1016/j.jneb.2016.08.006>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available: <http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Low Nutrient Intake And Frailty Among Overweight And Obese Migrant Women From Ethnically Diverse Backgrounds Aged 60+ Years: A Mixed-Methods Study

Research Article

Diana Castaneda-Gameros, MSc¹; Sabi Redwood, EdD MA RGN/RSCN^{2,3}; Janice L. Thompson, PhD, FACSM¹.

¹School of Sport, Exercise & Rehabilitation Sciences, University of Birmingham B15 2TT, UK

²School of Social & Community Medicine, University of Bristol BS8 1TH, UK.

³ NIHR CLAHRC West, 9th Floor, Whitefriars BS1 2NT, UK

Address for correspondence:

Diana Castaneda-Gameros, MSc.,

School of Sport, Exercise & Rehabilitation Sciences, University of Birmingham
B15 2TT, UK.

Phone: +44 (0)121 414 8745

Email: dxc242@student.bham.ac.uk

Acknowledgements:

The authors thank the study participants who participated in this study and the organizations that helped with recruitment. This study and DCG were partially supported by The National Council on Science and Technology (Mexico) and by seed-corn funding from the Institute for Research into Superdiversity (University of Birmingham). SR's time is supported by the National Institute for Health Research (NIHR) Collaboration for Leadership in Applied Health Research and Care West (CLAHRC West) at University Hospitals Bristol NHS Foundation Trust.

ABSTRACT

Objective: To examine associations between energy/nutrient intakes and frailty in older migrant women, and to explore perceptions of body weight, dietary intake and physical function.

Design: Cross-sectional mixed-methods study.

Setting: Birmingham, UK.

Participants: Seventy-six first generation migrant women ≥ 60 years.

Main Outcome Measures: Energy/nutrient intakes (assessed by 24-hr dietary recall), frailty (using the frailty phenotype), and links between perceptions of body weight, dietary intake, and physical function (via semi-structured interviews).

Analysis: Bivariate and logistic regression analyses examined associations between frailty and low energy/nutrient intakes. Interviews were analyzed using thematic analysis.

Results: Seventy-six women completed a 24-hr dietary recall; 46 participated in a semi-structured interview. Low energy intake was associated with frailty (OR: 11.71, 95% CI: 2.36-57.97). After adjusting for energy and other confounders, a low intake of >3 nutrients was associated with frailty (OR: 6.58, 95% CI: 1.01-43.08).

Qualitative data suggest that dietary intake was influenced by concerns about body weight and perceptions that unhealthy foods reduce mobility.

Conclusions and Implications: Among older migrant women with high prevalence of overweight/obesity, an inadequate dietary intake may be a stronger predictor of frailty than weight loss. Dietary interventions should focus on healthy weight maintenance and optimization of nutritional adequacy and physical function.

Key words: Frailty; nutrient inadequacy; overweight/obesity; older women;
ethnically diverse

INTRODUCTION

1

2

3

4 Frailty has become the focus of extensive research due to the ever-increasing aging of
5 the global population. Frailty is characterized as a disorder of multiple physiological
6 systems in which homeostatic mechanisms start failing, increasing the risk of declines
7 in cognitive and physical function.¹⁻³ Furthermore, longitudinal studies have
8 demonstrated a greater prevalence of cardiovascular disease and diabetes among frail
9 older people,⁴ and a greater frailty burden for women in comparison to men.^{5,6}

10 Therefore, identifying and treating individuals at risk of frailty may help delay its
11 negative consequences and reduce the financial, social, and personal burdens these
12 consequences place upon individuals, families and societies.^{7,8}

13

14 One of the most widely used definitions of frailty is the frailty phenotype proposed
15 and validated by Fried and colleagues.² This battery of tests identifies people as frail
16 when they meet three or more of five criteria: relatively weak grip strength,
17 unintentional weight loss, self-reported exhaustion, slow walking speed and low
18 levels of physical activity (PA). The inclusion of unintentional weight loss is used as a
19 proxy measure of dietary inadequacy, which is congruent with the conceptualization
20 of frailty as a wasting disorder.²⁻³ However, obesity can also be linked with frailty, as
21 indicated by the greater risk of physical function decline and pro-inflammatory state
22 commonly found among older adults who are obese.⁹

23

24 In older adults, the use of unintentional weight loss in the definition of frailty is
25 problematic as this measure may not be sensitive enough to reflect reduced energy

26 and nutrient intakes.⁸ Weight loss will not occur if energy intake matches energy
27 expenditure, however a diet that is adequate in energy can still be deficient in certain
28 nutrients, increasing a person's risk for frailty. Therefore, we hypothesize that a low
29 intake of energy and selected nutrients is a stronger predictor of frailty in
30 overweight/obese older women from diverse ethnic backgrounds than unintentional
31 weight loss. There is limited evidence examining the association between frailty as a
32 syndrome and nutrient intakes, and this has been conducted in predominantly White
33 older adults.⁸ Thus, very little is known about these associations in older adults from
34 diverse ethnic backgrounds. In addition, to develop interventions that can effectively
35 delay or prevent frailty in older women from diverse ethnic backgrounds, more
36 information is needed to explore if there are links between perceptions of body
37 weight, dietary intake and physical function in a population with disproportionately
38 higher rates of overweight and obesity.

39

40 Therefore, the aims of this study were to: 1) examine the associations between
41 dietary/nutrient intake and frailty in a sample of older women (≥ 60 years) from
42 diverse ethnic backgrounds living in the UK; 2) to gain a greater understanding of the
43 potential links between women's perceptions of body weight, dietary intake and
44 physical function.

45

46

47

METHODS

48

49

50 **Study Design**

51

52 A cross-sectional, mixed-methods design was employed, using 24-hr dietary recall
53 interviews that were enhanced with the addition of a qualitative semi-structured
54 interview. These methods allowed for the quantitative estimate of energy/nutrient
55 intake and its association with frailty, as well as providing insights into women's
56 perceptions of their body weight, dietary intake and physical function.

57

58 **Recruitment and Participants**

59

60 A convenience sample of first generation migrant women from Ireland, Jamaica,
61 Montserrat, St Kitts and Nevis, India, Pakistan, Bangladesh, Yemen, Sierra Leone,
62 Somalia, and Eritrea were recruited to participate in the study. Inclusion criteria
63 included being at least 60 years of age, with no medical conditions affecting memory
64 (e.g., dementia), and the ability to walk 15ft with no or minimal assistance (i.e., use of
65 a walking stick). Community-dwelling women living on their own or with family
66 members were recruited using maximum variation sampling¹⁰ to achieve our goal of
67 recruiting a sample across the ranges of age, migration backgrounds, socio-economic
68 status, and main ethnic groups living in the geographic region. This was achieved by
69 using the most recent Birmingham census data to identify the most representative
70 migrant groups.¹¹ Community centres serving specific migrant and older adult groups
71 were contacted and informed about the study. Those in leadership roles at these

72 centers facilitated access to potential participants so they could be approached and
73 informed of the purpose of the study. Participants were recruited via word-of-mouth
74 and snowballing.¹² Ethics approval was granted by The University of Birmingham
75 Ethics Committee (reference No. ERN_13-0557). All participants provided written
76 informed consent.

77

78 **Data Collection**

79

80 Data were collected at the participants' time and location of choice (e.g., homes or
81 community centers). For participants not fluent in English, trained interpreters fluent
82 in Punjabi, Bengali, Arabic and Somali provided simultaneous translation during
83 recruitment and data collection. Socio-demographic information was gathered via a
84 researcher-administrated questionnaire.

85 ***Dietary Intake.***

86 A multiple-pass 24-hr dietary recall interview was conducted with all participants to
87 gather data on the types and amounts of foods consumed on the previous day via a
88 standard protocol.¹³ Information was also obtained on nutrient supplement use. A
89 photographic food atlas assisted with the estimation of portion sizes.¹⁴ The first author
90 (DCG), a dietitian, trained in dietary assessment conducted all 24-hr dietary recalls.
91 Data coding and processing was conducted by DCG, with oversight from JLT who
92 has extensive expertise in dietary assessment. These procedures enabled a
93 standardized data entry and analysis process. The dietary recall interview was audio-
94 recorded to ensure accuracy of quantitative data entry and to facilitate the collection
95 of additional qualitative information. When participants stated that the previous day
96 did not reflect their habitual diet (e.g., they had engaged in fasting practices), the 24-

97 hr dietary recall was repeated later in the same week on a day that was identified by
98 participants as being representative of their habitual intake. This occurred in 5
99 participants. Data were not gathered during periods of major religious observances
100 (e.g., Ramadan, Diwali). All recalls were conducted during weekdays, excluding
101 Monday. Nutrient analysis was completed using DietPlan 6.0 software (Forestfield
102 software Ltd 2006, Horsham, UK), which included standard and supplemental food
103 composition databases that covered the range and ethnic diversity of foods consumed
104 in the UK.

105

106 Similarly to methods reported by Bartali and colleagues,⁸ low intake was defined as
107 the lowest quintile of the distribution of energy (<13 kcal/kg) and specific nutrients:
108 protein <30 g, vitamin D <0.5 µg, vitamin E <2.5 mg, retinol <101 µg, vitamin C <32
109 mg, folate <127 µg, iron <5.6 mg, calcium <349 mg, and zinc <3.6 mg. A nutritional
110 score was obtained by summing the number of nutrients categorized as low intake.
111 This nutritional score was subsequently categorized into a low intake of 0, 1-3, or >3
112 nutrients. A low intake of >3 nutrients was classified as poor nutritional status.

113

114 *Anthropometric measures and assessment of frailty.*

115 Anthropometric measures included height measured to the nearest mm (SECA 213
116 portable stadiometer), weight to the nearest 0.1 kg (SECA 899 digital scale), and hip
117 and waist circumference (WC) measured to the nearest cm using an extractable tape
118 measure. All anthropometric measurements were taken with the participant wearing
119 light clothing and no shoes. Body mass index (BMI) was calculated as weight divided
120 by height squared (kg/m^2), and waist-to hip-ratio (WHR) as waist circumference
121 divided by hip circumference (cm).

122 Frailty status was assessed using a modified version of the original frailty definition
123 developed by Fried and colleagues.² This included: 1) Exhaustion, defined using self-
124 reported fatigue from two questions from the Center for Epidemiological Studies-
125 Depression (CES-D) depression scale (“I felt that everything I did was an effort,” and
126 “I could not get going.”) Participants who reported having these feelings for ≥ 3 days
127 over the previous week to either or both questions received positive scores for
128 exhaustion; 2) Slow walking speed, with the highest quintile of the time needed to
129 walk a distance of 15 feet, adjusted by height (>14.5 seconds for height ≤ 157.7 cm
130 and >9.7 seconds for height > 157.7 cm); 3) Weak grip strength was defined as the
131 lowest quintile for adjusted grip strength using a JAMAR hand-held dynamometer
132 (Sammons Preston Rolyan, Bolingbrook, Illinois, USA), adjusted by BMI.
133 Participants met the criteria for weak grip strength if their BMI and grip strength were
134 ≤ 25.8 kg/m² and ≤ 12 kg; >25.9 - 29.6 kg/m² and ≤ 11 kg; >29.7 - 31.6 kg/m² and ≤ 12
135 kg; and ≥ 31.7 kg/m² and ≤ 14 kg. A low level of PA was defined as the lowest quintile
136 of caloric expenditure (< 60 kcal/week) using the International Physical Activity
137 Questionnaire short-form modified for the elderly (IPAQ-E).¹⁵ This version of the
138 IPAQ provides examples of activities that are more common among older adults and
139 has shown a moderate correlation ($r=0.347$, $p<0.01$), and moderate agreement κ
140 ($95\%CI= 0.448$ (0.18 - 0.72 , $p < 0.001$) with accelerometry.¹⁶
141
142 Since the purpose of this study was to examine the association between dietary intake
143 and frailty, similar to Bartali’s study, unintentional weight loss (>10 pounds in the last
144 year) was excluded from the original frailty definition.⁸ Therefore, participants with
145 >2 positive criteria were categorized as frail, while those with ≤ 1 positive criteria
146 were categorized as not frail.

147

148 ***Semi-structured interviews.***

149 A purposive sub-sample (n=46) across the range of age, ethnic groups and socio-
150 economic status was invited to participate in an interview that was guided by a list of
151 topics related to migration histories, dietary intake and eating behaviors, and
152 engagement in PA (migration histories and PA data not reported here). For the
153 purpose of this study, dietary topics centered on participants' perceptions of their diets
154 in relation to their body weight and frailty status (referred to as physical function
155 during the interviews). The interview guide was pilot-tested prior to the study and was
156 further revised via an iterative process throughout the data collection period. All
157 interviews were audio-taped and transcribed verbatim, with the interviews conducted
158 with participants who were not fluent in English being translated from their native
159 language into English by a trained interpreter during the interview process (n=16).

160

161 **Data Analysis**

162

163 ***Quantitative data analysis.***

164 Descriptive characteristics (means, SDs, and percentages) were calculated for socio-
165 demographic variables. To identify potential confounding factors, independent t-tests
166 or Mann-Whitney U tests (for non-parametric data) were conducted to examine any
167 significant differences in continuous variables between those classified as frail or not
168 frail, with Chi-squared or Fisher's exact tests conducted for categorical variables.
169 Point-biserial correlations (r_{pb}) were used to determine the association between frailty
170 status (dichotomous variable), weight loss, and indices of overweight/obesity (e.g.,

171 BMI, WC and WHR). Multiple logistic regressions were used to evaluate the
172 association between frailty status and each of its components with low energy intake
173 and poor nutritional status. Separate models were conducted to test the association
174 between nutrient intakes with frailty adjusting for confounding factors and energy
175 intake. All statistical analyses were performed using SPSS version 21.0 (SPSS INC.,
176 Chicago, IL); alpha was set at $p < 0.05$.

177

178 *Qualitative data analysis.*

179 An inductive thematic analysis of the interview transcripts was conducted, allowing
180 for the identification of themes being driven by participants' perspectives of their
181 diets in the context of their body weight and physical function/frailty status rather
182 than fitting the data into a pre-existing theoretical framework.¹⁷ Initially, a subset of
183 transcripts were read several times by the first author and two independent researchers
184 to identify predominant topics across the data. An initial coding frame using
185 qualitative analysis software (QSR NVivo, version 10) was developed which formed
186 the basis of broad coding and analysis. All of the transcripts were then coded by the
187 first author. The coding frame was discussed and refined by all authors until
188 consensus was reached. Data saturation was considered to have been achieved when
189 no new or relevant information emerged from each of the various ethnic groups
190 included in the study.¹⁸

191

192

RESULTS

193

194

195 Table 1 includes the demographic characteristics of participants. On average,
196 participants (mean age= 70.5 ± 7.6 years) reported having 2.3 ± 1.5 diseases
197 previously diagnosed by a doctor, with hypertension, arthritis and type 2 diabetes the
198 most common. Over 88% of the sample was classified as overweight or obese. BMI
199 cut-points for overweight and obesity among the Arab, Indian, Pakistani and
200 Bangladeshi participants were those recommended by the World Health Organization
201 for Asian populations.¹⁹ Although participants came from all socioeconomic levels,
202 79% were categorized as being in the two most socio-economically deprived quintiles
203 based on the English indices of deprivation.²⁰ Seventeen participants (22.4%) were
204 classified as frail, while 23 (30.3%) and 36 (47.4%) were classified as pre-frail and
205 non-frail, respectively. Frail participants were older and had a higher number of
206 diagnosed diseases; these were the only demographic variables that were statistically
207 different between frail and non-frail participants.

208 **Frailty and low nutrient intake**

209

210 Among frail participants, 82.3% had a low nutrient intake of at least one selected
211 nutrient (Table 2). The percentage of women with frailty increased with the greater
212 number of nutrients classified as low intake. Logistic regression analyses indicated
213 that low energy intake was independently associated with frailty (odds ratio [OR]:
214 11.71, 95% confidence interval [CI]: 2.36-57.97). After adjusting for energy, age and
215 number of diseases, poor nutritional status (>3 low nutrient intakes) was significantly
216 associated with frailty (OR: 6.58, 95% CI: 1.01-43.08) in comparison to those women
217 who did not have a low intake of any nutrients. After adjusting for energy and other
218 confounding variables, only slow walking speed was significantly associated with
219 poor nutritional status (OR: 1.86, 95% CI: 1.31-3.07).

220

221 In addition, a low intake of retinol (OR: 10.33, 95% CI: 1.55- 68.94) and zinc (OR:
222 8.47, 95% CI: 1.04-68.80) were significantly associated with frailty after adjustment
223 for energy intake and other confounding variables (Table 3). Self-reported weight loss
224 ($p=0.3$ for Fisher's exact test), BMI ($r_{bp}= 0.09$, $p=0.4$), waist circumference ($r_{bp}= 0.2$,
225 $p=0.1$), and WHR ($r_{bp}= 0.03$, $p=0.8$) were not associated with frailty.

226

227 **Qualitative Interview Results**

228

229 Two main themes which linked women's perceptions of body weight, dietary intake
230 and physical function were identified. They were: 1) concerns about weight and body
231 image; and 2) perceptions about negative effects of unhealthy foods on physical
232 function and health. Specific quotes from participants have been used to demonstrate
233 the themes outlined above.

234

235 ***Weight and body image concerns.***

236 Weight and body image emerged as two issues that were particularly important to
237 participants. Data suggest that these women have become more aware of their weight
238 as they have aged. Furthermore, some participants emphasized that their weight status
239 worried them more than getting older or other health problems as the excerpts below
240 indicate:

241

242 *'I am very careful that I don't eat too much, though I am very hungry but I will leave*
243 *[the food uneaten] ... I never say I want to eat more, no! ... I do not want to put on*

244 *weight, that is in the back of my mind, I never think of the heart [problem], I think of*
245 *my weight’ (Indian, 73y).*

246

247 *‘It doesn't bother me [the age], but when somebody says you are fat, then it hurts me!’*
248 *(Indian, 62y).*

249

250 Participants’ narratives also highlighted a difficult relationship between their diets and
251 body weight, leading to feelings of frustration and shame:

252

253 *‘My thinking was always eating healthy, but...I don't know how I put on so much*
254 *weight so quickly and I've been trying [to lose weight] for many years now, it's not*
255 *going down. I don't know what happened... I have gained so much I can't even get rid*
256 *of it... since I've put on weight and I am out of size as well, I think ‘Oh God people,*
257 *don't see me!’ ...That stops me from going out, dressing up as well, meeting people or*
258 *going into places’ (Pakistani, 62y)*

259

260 Given pervasive concerns about weight gain, many participants described modifying
261 their diets in an effort to lose weight. However, adopting more restrictive diets have
262 led some women to link these changes with a negative impact on their strength:

263

264 *‘When you are getting older is hard to lose weight ...well, I used to cut down my food*
265 *and then I think I was falling apart, I was getting weak... so I just said, “I'll just*
266 *continue [as normal] ”’ (African-Caribbean, 79y)*

267 Other participants who have also tried to reduce their food intake mentioned that they
268 occasionally complement their '*light diets*' with certain food items in order to meet
269 their perceived dietary requirements:

270

271 *'When I feel I haven't had enough protein... and need to rebuild some of the cells,*
272 *dying cells, ...then I would consciously have fish or chicken and try to eat a large*
273 *portion to try to convince myself that I'm eating enough protein...but no, I do a lot of*
274 *light days [of decreased consumption of fat and animal products]'* (African-
275 Caribbean, 68y).

276

277 ***Perceptions about negative effects of unhealthy foods on physical function and***
278 ***health.***

279 Participants' perceptions about the link between diet, physical function and general
280 health were mainly driven by their beliefs about the negative effects unhealthy foods
281 have on their mobility. For instance, some participants mentioned that eating
282 '*fattening food*' decreases their ability to be more active:

283

284 *'If I had fried food and I walk, I feel breathless yeah, so I keep in line what I am*
285 *eating'* (Indian, 71y)

286

287 *'Like...when you eat chips [French fries] you feel so heavy and you don't feel like*
288 *moving, you don't feel like running you know'* (Indian, 74y)

289

290 Overall, women felt that the quality of the food they eat is associated with their
291 general health, and that a healthy diet is an important component of healthy aging:

292 *'Health is related to what you put in your body, you are what you eat and if you put*
293 *healthy food in your body, you can expect to be healthy at this age' (African-*
294 *Caribbean, 69y)*

295

296

DISCUSSION

297

298

299 The present study examined the association between dietary intake and frailty in a
300 group of free-living first generation migrant older women using a mixed-methods
301 approach. Findings from this study indicated that having a low energy intake was
302 associated with frailty, and a poor nutritional status was significantly associated with
303 frailty after adjusting for energy and other confounding factors. Poor nutritional status
304 was also associated with slow walking speed, one of the criteria of the frailty
305 syndrome. The findings also provided rich insight into participants' perceptions about
306 the links between their body weight, dietary intake, and physical function.

307 Our findings support existing evidence associating frailty and its components to
308 nutrition at the nutrient level.⁸ Poor nutritional status and low serum levels of several
309 nutrient biomarkers (serum carotenoids, α -tocopherol, 25-hydroxyvitamin D, and
310 vitamin B6) have been found to be related to an increased risk of frailty among
311 predominantly White older adults.²¹⁻²³ These data, in addition to the findings from the
312 present study, suggest that an inadequate diet plays a crucial role in the physical
313 function of older adults. This is of particular importance due to the body composition
314 changes associated with old age leading to loss of muscle mass (sarcopenia) that can
315 contribute to morbidity and decreased quality of life.³

316 There are multiple pathways in which micronutrient deficiencies can increase the risk
317 of frailty in older adults by promoting conditions commonly associated with older age
318 such as oxidative stress, impaired immunity, muscle and bone metabolism, and
319 inflammation.²⁴ In our study, only retinol and zinc were independently associated
320 with frailty, suggesting that these two nutrients may be of particular concern in this
321 sample. Retinol is suggested to protect cell membranes from oxidative damage related
322 to aging,²⁵ while both retinol and zinc play an important role in maintaining the
323 integrity of the immune system.²⁶ Although malnourishment is typically associated
324 with underweight, this study confirms that overweight/obese individuals can also be
325 malnourished due to consuming a poor quality diet.²⁷ Thus, an individual can be frail
326 and not necessarily experience weight loss.

327 Among this sample, body weight concerns emerged as a key factor influencing energy
328 and nutrient intake. Therefore, the majority of participants were more conscious about
329 eating in moderation in order to lose weight, and did not identify being concerned
330 with how their dietary intake would affect nutrient adequacy. Although it is well
331 known that body dissatisfaction is highly associated with dietary intake in younger
332 adults, it is only recently that this has been reported in older adults, especially in
333 women.²⁸ Among women from minority ethnic groups, body weight perceptions have
334 been reported to be more positive and accepting of larger figures and a body weight
335 consistent with medically defined overweight or obesity.²⁹ However, our findings
336 indicate that the women in this ethnically diverse sample are concerned about their
337 body weight and the negative consequences associated with overweight/obesity.
338 These concerns may potentially lead them to adopt restrictive eating practices that
339 may cause more harm than good. Although body dissatisfaction has been previously
340 reported in younger migrant women,³⁰ to our knowledge, this is the first time that this

341 has been found in a sample of older migrant women with high rates of
342 overweight/obesity.

343 Regarding the negative effects of unhealthy foods on physical function and health, a
344 few studies have found that an unhealthy diet (i.e., poor consumption of fruits and
345 vegetables, low adherence to a Mediterranean-type diet) is associated with mobility
346 limitations and disability in older adults, particularly in women.³¹⁻³³ Although this
347 association has been found to be stronger in non-obese individuals,³² in our study
348 women felt that unhealthy foods, particularly fatty foods, were negatively related to
349 their mobility. Thus, in overweight/obese older women, healthier diets may be
350 perceived as a means of ameliorating mobility loss and further physical decline.

351 Given pervasive concerns about weight gain, findings from this study suggest that
352 older women from ethnically diverse backgrounds with a high prevalence of
353 overweight/obesity need dietary advice that promotes both the maintenance of a
354 healthy body weight and nutrient adequacy. Particularly, because both excess weight
355 and nutritionally inadequate diets are important determinants of morbidity and
356 premature mortality.³⁴

357 The major strength of the present study is the inclusion of a population commonly
358 under-represented in research,³⁵ and little is known about dietary intake, eating
359 behaviors, and frailty in older migrant women. The mixed-methods methodology is
360 also a strength, as it allowed for the examination of dietary intake and its association
361 with frailty as well as providing important insights into women's perceptions of their
362 dietary intake and its link with body weight and physical function. In addition, the
363 interview sample size was relatively large for a mixed-methods study, and data
364 saturation was reached in all participants across the range of age and ethnic groups.

365 Finally, some limitations of the study need to be considered. Due to the cross-
366 sectional study design and a relatively small sample size for the quantitative data,
367 causal inferences cannot be made and findings may not be generalizable to the wider
368 population of first generation older migrant women living in the UK. In addition,
369 almost 90% of the sample was overweight or obese. Although this could be
370 considered a strength as the sample reflects the higher prevalence of
371 overweight/obesity in ethnic groups in the UK,³⁶ the findings do not include data from
372 participants who were underweight. This could have limited the potential of finding
373 an association between frailty, protein and other micronutrients consistently found in
374 previous studies.^{8, 23, 37} In addition, BMI was used as a measure of weight status. This
375 is problematic as BMI does not distinguish between lean tissue and fat mass, and
376 cannot take into account the height loss that occurs with older age.³⁴ Studies including
377 a larger sample of older women from ethnically diverse backgrounds using an
378 accurate measure of body composition and nutritional biomarkers are needed to
379 confirm our findings. A larger sample will also allow for the examination of
380 significant differences between ethnic groups.

381 Another important limitation was the use of a single 24-hr dietary recall, a limitation
382 shared with other studies conducted with older adults and ‘hard to reach’
383 populations.³⁸ This method was considered the most appropriate as it minimized
384 participant burden and allowed participants with limited English literacy to fully
385 participate in both the quantitative and qualitative aspects of the study. Limitations in
386 willingness of participants to participate in a second 24-hr dietary recall interview, in
387 addition to budgetary constraints, prevented the use of repeated 24-hr dietary recalls.
388 In the present study, energy intake was relatively low and as such, under-reporting
389 cannot be ruled out. Under-reporting has been found to be associated with female

390 gender, higher age, lower socio-economic status, and overweight/obesity.³⁹ Because
391 of the day-to-day variability in dietary intake, the single 24-dietary recall provided
392 data for the sample rather than an estimate of an individual's dietary intake. The
393 interviews were conducted by a trained nutritionist, and when necessary with the aid
394 of interpreters with the same ethno-cultural background who were familiar with the
395 participants' dietary habits. In addition, we enhanced the 24-hr dietary recall with an
396 in-depth probing interview that allowed for a rich exploration of habitual dietary
397 behaviors not possible with a standard 24-hr dietary recall. Low dietary and nutrient
398 intakes in older adults are not uncommon given important changes in body
399 composition, intestinal absorption and decreased levels of PA.⁴⁰ In our study, women
400 were highly sedentary, which could have also influenced their energy intake.
401 Nevertheless, misreporting may have occurred and as such, our results should be
402 interpreted in light of this limitation.

403

404 **IMPLICATIONS FOR RESEARCH AND PRACTICE**

405

406

407 Findings from this study indicate that among a group of mainly overweight/obese
408 migrant women from ethnically diverse backgrounds, poor nutritional status is an
409 independent predictor of frailty. Given that weight loss may not necessarily be present
410 in community-dwelling older women, low energy and nutrient intakes make important
411 contributions to the development of frailty. Therefore, assessing dietary intake may
412 assist with screening for, and treating, frailty. Moreover, the mismatch found between
413 body weight and dietary inadequacy may potentially cause older women to engage in
414 self-imposed dietary restrictions that could cause further health problems. Future

415 strategies to prevent and detect frailty in this sub-group of the population should focus
416 on maintenance of a healthy body weight as well as in the overall nutritional quality
417 of the diet.

418

419

REFERENCES

420

421

422 1. Clegg A, Young J, Iliffe, S, et al. Frailty in elderly people. *Lancet*. 2013;381:752-
423 762.

424

425 2. Fried LP, Tangen CM, Walston J., et al. Frailty in older adults: evidence for a
426 phenotype. *J Gerontol A Biol Sci Med Sci*. 2001;56:M146-56.

427

428 3. Calvani R, Marini F, Cesari M, et al. Biomarkers for physical frailty and
429 sarcopenia: state of the science and future developments. *J Cachexia Sarcopenia*
430 *Muscle*. 2015;6:278-86.

431

432 4. Gill TM, Gahbauer EA, Han L, et al. Trajectories of disability in the last year
433 of life. *N Engl J Med*. 2010;362:1173-80.

434

435 5. Collard RM, Boter H, Schoevers RA, et al. Prevalence of frailty in community-
436 dwelling older persons: a systematic review. *J Am Geriatr Soc*. 2012;60:1487-92.

437

438 6. Hubbard RE, Rockwood K. Frailty in older women. *Maturitas*. 2011;69:203-207.

439

- 440 7. Yang Z, Hall AG. The financial burden of overweight and obesity among elderly
441 Americans: the dynamics of weight, longevity, and health care cost. *Health Serv Res.*
442 2008;43: 849-868.
443
- 444 8. Bartali B, Frongillo EA, Bandinelli S, et al. Low nutrient intake is an essential
445 component of frailty in older persons. *J Gerontol A Biol Sci Med Sci.* 2006;61:589-93.
446
- 447 9. Jensen GL, Hsiao PY. Obesity in older adults: relationship to functional
448 limitation. *Curr Opin Clin Nutr Metab Care.* 2010;13:46-51.
449
- 450 10. Teddlie C, Yu F. Mixed methods sampling a typology with examples.
451 *J Mix Methods Res.* 2007; 1:77-100.
452
- 453 11. Birmingham City Council. Population & Census.
454 <http://www.birmingham.gov.uk/census>. Accessed March 5, 2016.
455
- 456 12. Penrod J, Preston DB, Cain RE, et al. A discussion of chain referral as a method
457 of sampling hard-to-reach populations. *Journal Transcult Nur.* 2003;14:100-7.
458
- 459 13. Adamson AJ, Collerton J, Davies K, et al. Nutrition in advanced age: dietary
460 assessment in the Newcastle 85+ study. *Eur J Clin Nutr.* 2009;63:S6-18.
461
- 462 14. Nelson M, Atkinson M, Meyer J. *A photographic atlas of food portion sizes.*
463 MAFF publications.1997.
464

- 465 15. International Physical Activity Questionnaire website. [http://www.ipaq.ki.se/](http://www.ipaq.ki.se/ipaq.htm)
466 ipaq.htm. Accessed March, 2015.
467
- 468 16. Hurtig-Wennlöf A, Hagströmer M, Olsson LA. The nternational Physical
469 Activity Questionnaire modified for the elderly: aspects of validity and feasibility.
470 *Public Health Nutr.* 2010;13:1847-54.
471
- 472 17. Braun V, Clarke V. Using thematic analysis in psychology. *Qual Health*
473 *Res.* 2006;3:77-101.
474
- 475 18. Bunce GG, Johnson L. How many interviews are enough? An experiment
476 with data saturation and variability. *Field method.* 2006;18:59-82. doi:
477 10.1177/1525822X05279903
478
- 479 19. WHO, Expert Consultation. 2004. Appropriate body-mass index for Asian
480 populations and its implications for policy and intervention strategies. *Lancet.*
481 2004;363:57.
482
- 483 20. Department for Communities and Local Government. 2011. English indices of
484 deprivation. [https://www.gov.uk/government/collections/english-indices-of](https://www.gov.uk/government/collections/english-indices-of-deprivation)
485 deprivation. Accessed March, 2015.
486
- 487 21. Wilhelm-Leen ER, Hall YN, Deboer IH, et al. Vitamin D deficiency and frailty in
488 older Americans. *J Intern Med.* 2010;268:171-80.
489

- 490 22. Michelon E, Blaum C, Semba RD, et al. Vitamin and carotenoid status in older
491 women: associations with the frailty syndrome. *J Gerontol A Biol Sci Med Sci*.
492 2006;61:600-7.
493
- 494 23. Semba RD, Bartali B, Zhou J, et al. Low serum micronutrient concentrations
495 predict frailty among older women living in the community. *J Gerontol A Biol Sci*
496 *Med Sci*. 2006;61:594-599.
497
- 498 24. Bonnefoy M, Berrut G, Lesourd B, et al. Frailty and nutrition: searching for
499 evidence. *J Nutr Health Aging*. 2015;19:250-7.
500
- 501 25. Denke MA. Dietary retinol--a double-edged sword. *JAMA*. 2002. 287(1):102-4.
502
- 503 26. Haase H, Rink L. The immune system and the impact of zinc during aging. *Immun*
504 *Ageing*. 2009;6:9.
505
- 506 27. Popkin BM. The nutrition transition and obesity in the developing world. *J Nutr*.
507 2001;131: 871S-873S.
508
- 509 28. Roy M, Shatenstein B, Gaudreau P, et al. Seniors' Body Weight Dissatisfaction
510 and Longitudinal Associations with weight changes, anorexia of aging, and obesity.
511 Results from the NuAge Study. *J Aging Health*. 2015;27:220-238.
512
- 513 29. Lynch EB, Kane J. Body size perception among African American women. *J Nutr*
514 *Educ Behav*. 2014;46:412-417.

515

516 30. Greenhalgh T, Chowdhury MM, Wood GW. Big is beautiful? A survey of body
517 image perception and its relation to health in British Bangladeshis with diabetes.

518 *Psychol Health Med.* 2005;10:126-138.

519

520 31. Xu B, Houston DK, Locher JL, et al. Higher Healthy Eating Index-2005 scores
521 are associated with better physical performance. *J Gerontol A Biol Sci Med Sci.*

522 2012;67:93-99.

523

524 32. Koster A, Penninx BW, Newman AB, et al. Lifestyle Factors and Incident
525 Mobility Limitation in Obese and Non-obese Older Adults. *Obesity.* 2001;15:3122-

526 32.

527

528 33. Telegawkar SA, Bandinelli S, Bandeen-Roche, K, et al. A higher adherence to a
529 Mediterranean-style diet is inversely associated with the development of frailty in

530 community-dwelling elderly men and women. *J Nutr.* 2012;142:2161-2166.

531

532 34. Villareal DT, Apovian CM, Kushner RF, et al. Obesity in older adults: technical
533 review and position statement of the American Society for Nutrition and NAASO,

534 The Obesity Society. *Am J Clin Nutr.* 2005;82:923-34.

535

536 35. Redwood S, Gill PS. Under-representation of minority ethnic groups in research—
537 call for action. *Br J Gen Pract.* 2013; 63:342-343.

538

539 36. Landman J, Cruickshank, JK. A review of ethnicity, health and nutrition-related

- 540 diseases in relation to migration in the United Kingdom. *Public Health Nutr.*
541 2001;4:647-657.
- 542
- 543 37. Beasley JM, LaCroix AZ, Neuhaus ML, et al. Protein intake and incident frailty
544 in the Women's Health Initiative observational study. *J Gerontol A Biol Sci Med Sci.*
545 2010;58:1063-71.
- 546
- 547 38. Jonnalagadda SS, Diwan S. Nutrient intake of First Generation Gujarati Asian
548 Indian Immigrants in the U.S. *J Am Coll Nutr.* 2002; 21:372380.
- 549
- 550 39. Poslusna K, Ruprich J, de Vries JH, Jakubikova M, van't Veer P. Misreporting of
551 energy and micronutrient intake estimated by food records and 24 hour recalls,
552 control and adjustment methods in practice. *Br J Nutr.* 2009;101:S73-85.
- 553
- 554 40. Payette H, Shatenstein B. Determinants of healthy eating in community-dwelling
555 elderly people. *Can J Public Health.* 2005;1:S27-31.
- 556

Table 1. Participant Demographic Characteristics

Variable	Mean \pm SD or % Total n=76	Mean \pm SD or % Non-frail n=59	Mean \pm SD or % Frail n=17	p values
Age (y)	70.5 \pm 7.6	69.9 \pm 6.5	74.1 \pm 9.3	0.04
Residency in the UK (y)	38.73 \pm 17.1	37.2 \pm 17.8	44.1 \pm 13.5	0.10
No. of diseases	2.3 \pm 1.5	2.1 \pm 1.5	3.3 \pm 1.2	<0.001
Ethnicity, %				0.06
African-Caribbean	21 (27.6)	14 (23.7)	7 (41.2)	
African	10 (13.2)	10 (10.6)	0 (0)	
Arab	8 (10.5)	5 (8.5)	3 (17.6)	
Indian	20 (26.3)	17 (28.8)	3 (17.6)	
Pakistani	7 (9.2)	6 (10.2)	1 (5.9)	
Bangladeshi	5 (6.6)	2 (3.4)	3 (17.6)	
Irish	5 (6.6)	5 (8.5)	0 (0)	
IMD quintile, %				0.20
1 (most deprived)	49 (64.5)	34 (57.6)	15 (88.2)	
2	11 (14.5)	10 (16.9)	1 (5.9)	
3	7 (9.2)	7 (11.9)	0 (0)	
4-5 (less deprived)	9 (11.8)	8 (13.6)	1 (5.9)	
Education, %				0.07
No qualifications	26 (34.2)	16 (27.1)	10 (58.8)	
Primary school	8 (10.5)	6 (10.2)	2 (11.8)	
Secondary school	18 (23.7)	15 (37.3)	3 (17.6)	
Tertiary	24 (31.6)	22 (37.3)	2 (11.8)	
Marital status, %				0.60
Married	34 (44.7)	31 (52.5)	3 (17.6)	
Widowed	30 (39.5)	21 (35.6)	9 (52.9)	
Single/ separated/divorced	12 (15.8)	7 (11.9)	5 (29.4)	
Living alone, %	26 (34.2)	20 (33.9)	6 (35.3)	0.60
BMI (kg/m ²)	29.3 \pm 4.9	29.1 \pm 4.8	30.2 \pm 5.3	0.43
Normal, %	9 (11.8)	9 (15.3)	0 (0)	
Overweight, %	23 (30.3)	16 (27.1)	7 (41.2)	
Obese, %	44 (57.9)	34 (57.6)	10 (58.8)	
WC (cm) ^a	98.8 \pm 10.8	97.8 \pm 11.1	102.0 \pm 9.3	0.15
WHR ^a	0.92 \pm 0.8	0.92 \pm 0.1	0.92 \pm 0.6	0.70
Unintentional weight loss, %	9 (11.8)	6 (10.2)	3 (17.6)	0.41
Supplement use, %	30 (39.5)	24 (40.7)	6 (35.3)	0.46
Energy intake (Kcals)	1243.5 \pm 524.4	1379.9 \pm 507.9	819.7 \pm 262.5	<0.01

Frailty score (No. of frailty components, %)				NA
0	36 (47.4)	36 (61)	0 (0)	
1	23 (30.3)	23 (39)	0 (0)	
≥2	17 (22.4)	0 (0)	17 (100)	

^a n=68, BMI= Body Mass Index, IMD= Index of Multiple Deprivation, WC= waist circumference, WHR= waist-to-hip ratio, NA=not applicable.

Table 2. Association Between Frailty Syndrome and Frailty Criteria According to the Number of Nutrients with Low Intake (n=76)

Number of nutrients with low intake				Adjusted Odds Ratios ^a	
Variables	0	1-3	>3	Low intake of 1-3 nutrients compared to 0	Low intake of >3 nutrients compared to 0
	%	%	%	OR (95% CI)	OR (95% CI)
Frailty syndrome	17.6	29.4	52.9	3.11 (0.56-17.35)	6.58 (1.01-43.08) ^b
Frailty criteria:					
Exhaustion	24.2	26.9	35.3	0.92 (0.26-3.17)	1.12 (0.17-7.20)
Low PA	9.1	19.2	47.1	2.30 (0.46-11.33)	5.26 (0.72- 38.10)
Weak grip strength	18.2	26.9	17.6	0.57 (0.15- 2.16)	1.23 (0.14-10.26)
Slow walking speed	6.1	15.4	47.1	0.85 (0.11-6.79)	1.86 (1.13-3.07) ^b

^a Adjusted for low energy intake, age and number of diseases; ^b p< 0.05

Table 3. Frailty Syndrome Associated with Specific Low Nutrient Intakes (n=76)

Nutrient intake ^a	<u>Frailty syndrome ^b</u> OR (95% CI)
Protein (g/day)	0.76 (0.09-5.99)
Retinol (µg/day)	10.33 (1.55- 68.94) ^c
Vitamin D (µg/day)	0.96 (0.18-5.19)
Vitamin E (mg/day)	0.98 (0.17-5.68)
Vitamin C (mg/day)	3.82 (0.67-21.64)
Folate (µg/day)	0.78 (0.12- 5.06)
Calcium (mg/day)	3.87 (0.65-22.85)
Iron (mg/day)	0.94 (0.17- 5.19)
Zinc (mg/day)	8.47 (1.04-68.8) ^c

^a Defined as the lowest quintile of each selected nutrient, ^b Adjusted for low energy intake, age and number of diagnosed diseases, ^c p<0.05